

## CORRECTION

# Correction: Worldwide increased prevalence of human adenovirus type 3 (HAdV-3) respiratory infections is well correlated with heterogeneous hypervariable regions (HVRs) of hexon

Ezazul Haque, Urmila Banik, Tahmina Monowar, Leela Anthony, Arun Kumar Adhikary

The third author's name is spelled incorrectly. The correct name is: Tahmina Monowar. The correct citation is: Haque E, Banik U, Monowar T, Anthony L, Adhikary AK (2018) Worldwide increased prevalence of human adenovirus type 3 (HAdV-3) respiratory infections is well correlated with heterogeneous hypervariable regions (HVRs) of hexon. PLoS ONE 13(3): e0194516. <https://doi.org/10.1371/journal.pone.0194516>

## Reference

1. Haque E, Banik U, Monowar T, Anthony L, Adhikary AK (2018) Worldwide increased prevalence of human adenovirus type 3 (HAdV-3) respiratory infections is well correlated with heterogeneous hypervariable regions (HVRs) of hexon. PLoS ONE 13(3): e0194516. <https://doi.org/10.1371/journal.pone.0194516> PMID: 29590206



## OPEN ACCESS

**Citation:** Haque E, Banik U, Monowar T, Anthony L, Adhikary AK (2018) Correction: Worldwide increased prevalence of human adenovirus type 3 (HAdV-3) respiratory infections is well correlated with heterogeneous hypervariable regions (HVRs) of hexon. PLoS ONE 13(4): e0196263. <https://doi.org/10.1371/journal.pone.0196263>

**Published:** April 18, 2018

**Copyright:** © 2018 Haque et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.