


CORRECTION

Correction: The cultivable autochthonous microbiota of the critically endangered Northern bald ibis (*Geronticus eremita*)

Joachim Spergser, Igor Loncaric, Alexander Tichy, Johannes Fritz, Alexandra Scope

The labels indicating significance are missing in Fig 5. Please see the corrected Fig 5 here.



OPEN ACCESS

Citation: Spergser J, Loncaric I, Tichy A, Fritz J, Scope A (2018) Correction: The cultivable autochthonous microbiota of the critically endangered Northern bald ibis (*Geronticus eremita*). PLoS ONE 13(5): e0197236. <https://doi.org/10.1371/journal.pone.0197236>

Published: May 8, 2018

Copyright: © 2018 Spergser et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Fig 5. Seasonal influence in the distribution of microbial taxa. Seasonal influence in the distribution of microbial taxa (bacterial taxa separated from fungal taxa by /) isolated from adult Northern bald ibis (only different-level phylotypes with at least 10% frequency in a season are presented).
a = significantly more common, b = significantly less frequent ($p < 0.05$).

<https://doi.org/10.1371/journal.pone.0197236.g001>

Reference

1. Spergser J, Loncaric I, Tichy A, Fritz J, Scope A (2018) The cultivable autochthonous microbiota of the critically endangered Northern bald ibis (*Geronticus eremita*). PLoS ONE 13(4): e0195255. <https://doi.org/10.1371/journal.pone.0195255> PMID: 29617453