

RESEARCH ARTICLE

# Interferon lambda is required for interferon gamma-expressing NK cell responses but does not afford antiviral protection during acute and persistent murine cytomegalovirus infection

Silvia Gimeno Brias<sup>1</sup>, Morgan Marsden<sup>1</sup>, Jessica Forbester<sup>1,2</sup>, Mathew Clement<sup>1</sup>, Cordelia Brandt<sup>2</sup>, Katherine Harcourt<sup>2</sup>, Leanne Kane<sup>2</sup>, Lucy Chapman<sup>1</sup>, Simon Clare<sup>2</sup>, Ian R. Humphreys<sup>1,2\*</sup>

**1** Institute of Infection Immunity, School of Medicine/Systems Immunity University Research Institute, Cardiff University, Cardiff, United Kingdom, **2** Wellcome Trust Sanger Institute, Hinxton, Cambridgeshire, United Kingdom

\* [humphreysir@cardiff.ac.uk](mailto:humphreysir@cardiff.ac.uk)



**OPEN ACCESS**

**Citation:** Gimeno Brias S, Marsden M, Forbester J, Clement M, Brandt C, Harcourt K, et al. (2018) Interferon lambda is required for interferon gamma-expressing NK cell responses but does not afford antiviral protection during acute and persistent murine cytomegalovirus infection. *PLoS ONE* 13(5): e0197596. <https://doi.org/10.1371/journal.pone.0197596>

**Editor:** Michael Nevels, University of St Andrews, UNITED KINGDOM

**Received:** November 30, 2017

**Accepted:** May 4, 2018

**Published:** May 16, 2018

**Copyright:** © 2018 Gimeno Brias et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper.

**Funding:** I.R.H. is funded by a Wellcome Trust Senior Research Fellowship in Basic Biomedical Sciences (WT098026 and 207503/Z/17/Z). S.G.B. is funded by a Medical Research Council Doctoral Training Grant (DTG). Experiments at the WTSI were also funded by the Wellcome Trust grant WT206194 to S.C. The funders had no role in study

## Abstract

Interferon lambda (IFN $\lambda$ ) is a group of cytokines that belong to the IL-10 family. They exhibit antiviral activities against certain viruses during infection of the liver and mucosal tissues. Here we report that IFN $\lambda$  restricts *in vitro* replication of the  $\beta$ -herpesvirus murine cytomegalovirus (mCMV). However, IFN $\lambda$ R1-deficient (*Ifnlr1*<sup>-/-</sup>) mice were not preferentially susceptible to mCMV infection *in vivo* during acute infection after systemic or mucosal challenge, or during virus persistence in the mucosa. Instead, our studies revealed that IFN $\lambda$  influences NK cell responses during mCMV infection. *Ifnlr1*<sup>-/-</sup> mice exhibited defective development of conventional interferon-gamma (IFN $\gamma$ )-expressing NK cells in the spleen during mCMV infection whereas accumulation of granzyme B-expressing NK cells was unaltered. *In vitro*, development of splenic IFN $\gamma$ <sup>+</sup> NK cells following stimulation with IL-12 or, to a lesser extent, IL-18 was abrogated by IFN $\lambda$ R1-deficiency. Thus, IFN $\lambda$  regulates NK cell responses during mCMV infection and restricts virus replication *in vitro* but is redundant in the control of acute and persistent mCMV replication within mucosal and non-mucosal tissues.

## Introduction

The  $\beta$ -herpesvirus human cytomegalovirus (HCMV) is typically controlled by immune-competent individuals. However, HCMV causes disease in immune-suppressed adults such as transplant recipients, and in immunologically immature children following congenital infection. HCMV infects multiple tissues within the host and is transmitted horizontally via urine, breast milk and mucosal secretions including saliva [1]. Thus, understanding how CMV is controlled in peripheral tissues and identifying the factors that regulate these responses may inform therapeutic strategies.

design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

Cytokines are important regulators of antiviral immunity during CMV infection (reviewed in [2]). The interleukin-10 (IL-10) cytokine superfamily consists of nine members that exhibit diverse functions ranging from immune regulation to tissue protection [3]. The importance of this superfamily in CMV pathogenesis is indicated by the evolutionary acquisition by HCMV [4] (and other herpesviruses [5, 6]) of IL-10 orthologues that suppress immune activation. Moreover, the rhesus CMV IL-10 orthologue restricts inflammation at the site of infection whilst suppressing long-lived virus-specific immunity *in vivo* [7]. Studies in the murine CMV (mCMV) model of infection, which recapitulates many aspects of HCMV tropism, immunity and pathogenesis [8] although notably not the acquisition of an obvious IL-10 orthologue, has highlighted the importance of cellular IL-10 in modulation of antiviral immunity during CMV infection *in vivo*. mCMV induces IL-10 [9], and the IL-10-IL-10R pathway inhibits virus-specific immunity and control of mCMV persistence in mucosal tissue [10–14] whilst limiting pathology and activation-induced NK cell death during acute infection [10, 15–17]. Additionally, the IL-10-related cytokine IL-22 is induced upon mCMV infection [18, 19] and restricts mCMV replication during acute infections of peripheral tissues via the recruitment of antiviral neutrophils [18]. Thus, IL-10 family members significantly influence CMV pathogenesis *in vivo*.

Interferon lambdas (IFN $\lambda$ 1–4, type III IFNs) are IL-10 superfamily members that exhibit functional parallels to type I IFNs with respect to induction of antiviral cellular immunity. In experimental models, IFN $\lambda$  limits replication of numerous viruses (reviewed in [20]), including herpesviruses [21, 22]. IFN $\lambda$  receptor (IFN $\lambda$ R) primarily signals in epithelial cells [23] and subsequently affords robust protection from viruses that target mucosal surfaces [24–28]. Given the importance of mucosal CMV infection in dissemination and pathogenesis, we investigated the mCMV model of infection to ascertain whether IFN $\lambda$  influences the outcome of acute and persistent cytomegalovirus infection in mucosal and other peripheral tissues *in vivo*.

## Materials and methods

### Mice, infections and ethics

Smith strain murine cytomegalovirus (mCMV) originally obtained from the ATCC was generated following *in vivo* propagation in weanling BALB/c mice and purification of salivary gland-derived virus performed by spinning organ homogenate over a sorbital gradient, as previously described [18]. *Ifnlr1<sup>tm1a(EUCOMM)Wtsi</sup>* (*Ifnlr1<sup>-/-</sup>*) mice were generated using gene targeting as part of the International Knockout Mouse Consortium ([www.knockoutmouse.org](http://www.knockoutmouse.org)) using high throughput methods as described by Skarnes *et al.*, 2011 [29]. *Ifnlr1<sup>-/-</sup>* and age/sex-matched wild type C57BL6/n mice were bred in-house at the Wellcome Trust Sanger Institute (WTSI) research support facility. Mice were infected with either  $3 \times 10^4$  or  $5 \times 10^4$  PFU mCMV (i.p.). In some experiments mice were infected with  $1 \times 10^4$  mCMV (i.n) under iso-fluothane anesthesia. Virus load in homogenized tissues was measured by plaque assay using 3T3 cells [18]. *In vitro* infections were performed in 3T3 and BNLCL2 cell lines (ATCC). Cells were treated with IFN $\lambda$ 2 (IL-28A) (Peprotech), IFN $\alpha$  or IFN $\beta$  (PBL Assay Science) for 24 hours prior to infection with Smith strain mCMV. After 4 days of infection, virus was assessed by plaque assay [18].

### Ethics statement

All mice experiments were performed under the UK Home Office-approved project Licence (Reference: PPL 80/2596) at the Wellcome Trust Sanger Institute research support facility. Iso-fluothane was used for anesthesia (for intranasal infections) and all mice were sacrificed according to UK Home Office guidelines.

## Assessment of *in vivo* immune responses

Splenocytes were isolated as previously described [16]. Liver leukocytes were isolated by passing leukocytes through a 70 $\mu$ M sieve prior to cell purification over a percoll gradient. NK cell responses were measured as previously described [16], with additional direct *ex vivo* assessment of intracellular granzyme B NK cell expression (Biolegend). Assessment of neutrophils and virus-specific T cell responses (quantified by detecting peptide-specific cytokine production) have been previously described [12, 14, 18]. IL-12 p70 (Biolegend), IL-18 (Thermo Fisher), IFN $\gamma$  (Biolegend) and IFN $\lambda$ 2/3 (IL-28A/B, R&D Systems) protein was measured using ELISAs.

## Measured for *Ifnlr1* gene expression

RNA was extracted from cells using RNAeasy mini kit (Qiagen). Genomic DNA was eliminated from the samples using the Turbo DNA-free™ kit (Ambion) prior to cDNA synthesis (Applied Biosystems). Gene expression was measured by quantitative reverse transcription PCR using a QuantStudio™ 3 thermal cycler (Thermo Fisher Scientific) and iTaq Universal SYBR® Green supermix (Bio-rad laboratories) using primers for *Ifnlr1* (Forward: 5' -GTG ACC TAT TTC GTG ACC TAC C-3', Reverse: R 5' -CTG CCT GTA CTC GTC CTT TG -3') and  $\beta$ -actin (Forward: 5' -TGC AGA TTC CTC TCC AGC AA-3', Reverse: 5' -GTC TTC ACC CCC TGA AAC CA-3').

## *In vitro* NK cell assays

For analysis of NK cell function, splenocytes were isolated from naïve *Ifnlr1*<sup>-/-</sup> mice and corresponding WT controls, plated in R10 medium supplemented with IL-12, IL-18 or IL-12/18 +/- IL-28a (10ng/ml, all Peprotech; 1 x 10<sup>6</sup> cells per condition), and incubated for 5h with the addition of monensin (BD Biosciences). Cells were then incubated with Fc block (BioLegend), surface stained with anti-NK1.1 and anti-CD3 $\epsilon$ , and then washed with DPBS (Gibco) and stained with zombie aqua dye (BioLegend). Surface-stained cells were then fixed and permeabilized using BD Cytofix/Cytoperm, and stained with anti-IFN- $\gamma$  (BioLegend). To detect IFN $\lambda$ -induced pSTAT1 activation in NK cells, splenocytes from naïve and mCMV infected mice were incubated for 1–6 hours with or without 10000U/mL IFN- $\beta$  or 50ng/mL IFN- $\lambda$ 2. Cells were then surface-stained and phosphorylated STAT1 (pSTAT1) was detected according to the manufacturer's instructions (BD Biosciences). Cells were analyzed on a Becton Dickinson FACSAriaIII using FACS Diva software (v8) or using an Attune NxT Flow Cytometer. Data were subsequently analysed using FlowJo software version 10.2.


## Statistics

Statistical significance was assessed using Mann-Whitney U for paired analysis of viral-load data whereas students T-test was used to analyze paired flow cytometry and ELISA data. 1-Way ANOVA was used to determine statistical significance where more than 2 groups were assessed concurrently (*in vitro* virus replication). For all tests performed, p values are reported as \* $\leq$ 0.05, \*\* $\leq$ 0.01, and \*\*\* $\leq$ 0.001.

## Results and discussion

### IFN $\lambda$ restricts mCMV replication *in vitro*

We first assessed whether IFN $\lambda$  directly influences mCMV replication, using murine cell lines representative of embryonic liver cells (BNLCL2) or fibroblasts (3T3) that expressed or did not express IFN $\lambda$ R, respectively (Fig 1A). Cells were pre-treated for 24 hours with or without



**Fig 1. IFN $\lambda$  can restrict mCMV replication *in vitro*.** (A) *Ifnlr1* expression by 3T3 and BNLCL2 cells was determined by qPCR. (B) 3T3 (top) and BNLCL2 (bottom) cells were incubated with/without 50U/ml IFN $\alpha$  and/or IFN $\beta$ , or 50ng/ml IFN $\lambda$ 2 (IL-28A) for 24hrs and infected with mCMV at multiplicities of infection (MOI), as stated in the figure. After 4 days, infectious virions in supernatant were quantified by plaque assay. Statistical significance of PFU in IFN $\lambda$ 2-treated versus control cells is shown. Virus load in spleen (C), liver (D) and salivary glands (E) of WT and *Ifnlr1*<sup>-/-</sup> mice was assessed 4 (D&E) and 33 (E) days p.i. (F) IFN $\lambda$ 2/3 protein in spleen (left), liver (middle) and salivary glands (right) was measured at day 0 and 2 days p.i. (spleen and liver) or 0 and 26 days p.i. (salivary glands). Results are shown as mean + SEM of 3–7 mice/group. (G) WT and *Ifnlr1*<sup>-/-</sup> mice were infected (i.n) with mCMV in a volume of 25 $\mu$ l or 50 $\mu$ l and after 4 days, lung infectious viral load was quantified by plaque assay. Statistical significance was assessed using 1-way ANOVA (B) or Mann Whitney-U (C-E, G) or students T-Test (F) and is depicted where appropriate. Panel G represents merged data from two experiments whereas all other data represent at least two biological replicates performed separately.

<https://doi.org/10.1371/journal.pone.0197596.g001>

IFN $\lambda$ 2, or as a positive control for cytokine-mediated control of mCMV replication, IFN $\alpha$  and/or IFN $\beta$ . In accordance with receptor expression data (Fig 1A), IFN $\lambda$  did not impact upon mCMV replication in 3T3 cells following infection with a range of infection inoculums (Fig 1B). However, incubation of BNLCL2 cells with IFN $\lambda$  prior to mCMV infection led to a reduction of replicative-competent virions in supernatant by  $\sim 1$  log (Fig 1B). Induction of IFN $\lambda$ R in human astrocytes enables IFN $\lambda$ -mediated control of HCMV replication *in vitro* [30]. Thus, our data is consistent with the conclusion that IFN $\lambda$  is capable of restricting human and mouse cytomegalovirus infection within IFN $\lambda$ R-expressing cells.


### IFN $\lambda$ does not impinge upon mCMV replication *in vivo*

We next studied IFN lambda receptor deficient mice (*Ifnlr1*<sup>-/-</sup>) to determine whether IFN $\lambda$  modulated mCMV replication *in vivo*. Upon systemic (i.p.) challenge, we observed no effect of IFN $\lambda$ R1 deficiency on control of acute mCMV replication in the spleen or liver (Fig 1C and 1D), in accordance with previously published observations [31]. Although IFN $\lambda$  restricted mCMV replication in a murine liver-derived cell line (Fig 1B), murine hepatocytes respond poorly to IFN $\lambda$  [32]. However, epithelial cells are responsive to IFN $\lambda$  *in vivo* [23]. Thus, we examined the impact of IFN $\lambda$ R1 deficiency on mCMV persistence in glandular epithelial cells within salivary glands. Again, *Ifnlr1*<sup>-/-</sup> mice exhibited no defect in control of mCMV (Fig 1E). The absence of an antiviral function of IFN $\lambda$  was not attributed to the lack of cytokine induction *in vivo* as significant expression was observed during acute and chronic infection (Fig 1F). Thus, we concluded that IFN $\lambda$  does not contribute to the control of mCMV *in vivo* following systemic infection.

IFN $\lambda$  restricts viral replication within the mucosa (reviewed in [20]). Using an established intranasal mucosal challenge model of mCMV infection [33–35], we investigated whether IFN $\lambda$  restricted primary mucosal mCMV infection. We challenged adult mice with mCMV either in 25 $\mu$ l or 50 $\mu$ l volumes, reasoning that 25 $\mu$ l would restrict mCMV delivery to the nose and not trachea and thus serve to specifically probe the impact of IFN $\lambda$ R signaling on control of initial mCMV replication within the nasal cavity. As shown in Fig 1G, we observed no impact of IFN $\lambda$ R1 deficiency on mCMV infection of lung tissue following mCMV infection using either volume (Fig 1G). Thus, although the caveat exists that the unresponsive nature of murine hepatocytes to IFN $\lambda$  may mask a possible antiviral role for IFN $\lambda$  in controlling CMV replication within the liver, overall our data supports the conclusion that IFN $\lambda$ R signaling plays no significant role in controlling mCMV replication *in vivo*.

### IFN $\lambda$ does not influence the development of mCMV-specific T cell immunity

IFN $\lambda$  has been implicated in shaping adaptive immunity [20]. Studies of lymphochoriomeningitis infection revealed that IFN $\lambda$  can restrict effector T cell responses and memory development following acute virus infection whilst paradoxically promoting T cell responses during virus chronicity [36]. Using intracellular staining for IFN $\gamma$  following *ex vivo* stimulation with mCMV-derived peptides, we measured mCMV-specific CD8<sup>+</sup> and CD4<sup>+</sup> T cell responses during acute (d7 p.i) and persistent (d26 p.i) infection. We detected no discernable differences in the accumulation of functional virus-specific CD4<sup>+</sup> or CD8<sup>+</sup> T cells between WT and *Ifnlr1*<sup>-/-</sup> mice at either time-point (Fig 2A–2D). Furthermore, IFN $\lambda$ R1 deficiency did not influence the frequency of mCMV-specific T cells that co-express TNF $\alpha$  following re-stimulation *ex vivo* (Fig 2E and 2F). Therefore, although these data do not preclude a role for IFN $\lambda$  in modulating T cell and humoral immunity during the chronic/latent phase of mCMV infection, these results demonstrate that IFN $\lambda$  does not alter the magnitude or quality of mCMV-specific T cell responses during acute and persistent mCMV infection.




**Fig 2. IFN $\lambda$ R signaling does not influence mCMV-specific T cell responses *in vivo*.** mCMV-derived peptide-specific CD8<sup>+</sup> (A&C) and CD4<sup>+</sup> (B&D) T cell responses in WT and *Ifnλr1*<sup>-/-</sup> mice infected for 7 (A&B) or 26 (C&D) days were quantified *ex vivo* following peptide re-stimulation. Mean + SEM of 6–7 mice/group is shown and represent 2–3 experiments. Statistical significance was tested using an unpaired student's T-test. (E) Representative bivariate FACS plots of IFN $\gamma$  and/or TNF $\alpha$  expression by CD8<sup>+</sup> T cells stimulation with media alone (M) or 2 $\mu$ g/ml MHC class I restricted peptide derived from mCMV M45 (HGIRNASFI) or M38 (SSPPMFRVP) proteins. % positive cells are shown in the plots. Data are representative of 6 mice/group. (F) Mean + SEM from 6 mice/group is shown. Results represent 2 separate experiments.

<https://doi.org/10.1371/journal.pone.0197596.g002>

### IFN $\lambda$ R1 promotes splenic IFN $\gamma$ <sup>+</sup> NK cell responses during mCMV infection

We next investigated whether IFN $\lambda$  modulates innate immune activation during mCMV infection. Neutrophils limit mCMV replication and pathogenesis [18, 37], and IFN $\lambda$  has been reported to directly modulate neutrophil responses *in vivo* [38, 39]. However, neutrophil accumulation in the spleen and liver during mCMV infection was unaltered in *Ifnλr1*<sup>-/-</sup> mice (Fig 3A). In contrast, however, studies of NK cell responses revealed a marked reduction in NK



**Fig 3. IFN $\lambda$  promotes IFN $\gamma$ <sup>+</sup> NK cell responses.** (A) Neutrophil numbers in the spleens and livers of mCMV-infected WT and *Ifnlr1*<sup>-/-</sup> mice was assessed 4 days p.i. IFN $\gamma$ <sup>+</sup> (B&D) and granzyme B<sup>+</sup> (C&E) NK cell accumulations in spleens (B-C) and livers (E&F) were quantified day 4 pi. Individual mice + mean is shown from 2 independent experiments and represent 4 (liver) or 5 (spleen) experiments in all. (D) Representative bivariate FACS plots of IFN $\gamma$  versus granzyme B expression by live NK1.1<sup>+</sup>CD3<sup>+</sup> cells 4 days post-infection. Plots show concatenated samples from 4 (WT) and 6 (*Ifnlr1*<sup>-/-</sup>) mice. Results represent 3 experiments. (G) IFN $\gamma$  protein in the serum was measured 4 days pi. Mean + SEM of 3–4 mice/group is shown. (H&I) IL-12 (H) and IL-18 (I) protein was measured in naïve spleen tissue or after 2 days of mCMV infection. Mean + SEM of 3 (naïve) or 7 (infected) mice is shown. (J) IFN $\gamma$  expression by NK cells stimulated

with IL-12, IL-18 or IL-12/IL-18 +/- IL-28A was detected by flow cytometry. Data represent the mean +/- SD for 3 mice per group. All experiments were performed at least twice.

<https://doi.org/10.1371/journal.pone.0197596.g003>

cells from spleens from *Ifnlr1*<sup>-/-</sup> mice during acute mCMV infection that spontaneously produce IFN $\gamma$  ex-vivo (Fig 3B). Defective IFN $\gamma$ <sup>+</sup> NK cell accumulation was not defined by a broad deficit in NK cell responses in *Ifnlr1*<sup>-/-</sup> mice as, although we observed a trend in reduced accumulation of cytotoxic (granzyme B<sup>+</sup>) splenic NK cells in *Ifnlr1*<sup>-/-</sup> mice, this was not statistically significant (Fig 3C and 3D). NK cell populations that are present in the liver form a distinct lineage from splenic NK cells [40]. Interestingly, we observed no statistically significant impact of IFN $\lambda$ R deficiency on the accumulation of IFN $\gamma$ <sup>+</sup> or granzyme B<sup>+</sup> hepatic NK cells (Fig 3E and 3F) nor did we detect a defect in systemic IFN $\gamma$  protein concentrations (Fig 3G), suggesting that IFN $\lambda$  preferentially promoted the development of IFN $\gamma$ <sup>+</sup> conventional splenic NK cell responses during mCMV infection rather than broadly impacting upon NK cell responses in other tissues and influencing systemic cytokine production.

We next examined the mechanism(s) through which IFN $\lambda$  regulates IFN $\gamma$ <sup>+</sup> NK cells. IL-18 and IL-12 promote these responses during mCMV infection [41–43]. However, IL-12 and IL-18 protein concentrations were comparable in spleen supernatants from WT and *Ifnlr1*<sup>-/-</sup> mice post-infection (Fig 3H and 3I), suggesting that IFN $\lambda$  did not promote IFN $\gamma$ <sup>+</sup> NK cell responses indirectly via the regulation of IFN $\gamma$ -inducing cytokines. Instead, incubating splenocytes from naïve mice with IL-12 revealed impaired development of *Ifnlr1*<sup>-/-</sup> IFN $\gamma$ <sup>+</sup> NK cells, as assessed by flow cytometry after 5hrs (Fig 3J). A similar, albeit not significantly significant, trend was observed in splenocytes incubated with IL-18 (Fig 3J). These data support the conclusion that IFN $\lambda$ R signaling promotes NK cell responsiveness to IFN $\gamma$ -inducing cytokines.

Souza-Fonseca-Guimaraes et al [31] previously reported no impact of IFN $\lambda$ R deficiency on cytokine-induced IFN $\gamma$  secretion by purified NK cells. In our experiments, we studied whole splenocytes and examined IFN $\gamma$ <sup>+</sup> NK cell generation by flow cytometry and not ELISA, and at earlier time-points than Souza-Fonseca-Guimaraes and colleagues. Thus, variations in experimental designs may explain these disparate findings. Importantly, however, we saw no influence of IFN $\lambda$ R signaling on IFN $\gamma$ <sup>+</sup> NK cell development that was induced following co-incubation of splenocytes with IL-12 and IL-18 (Fig 3H). Given that Souza-Fonseca-Guimaraes and colleagues performed assays using purified NK cells that involved co-incubation of multiple cytokines, our data suggest that the influence of IFN $\lambda$ R1 on IFN $\gamma$ <sup>+</sup> NK cell development may be redundant in situations where cells receive strong stimuli via concurrent stimulation of multiple cytokine receptors. Indeed, the observation that IL-12 and IL-18 are often co-expressed during mCMV infection [41] may explain why hepatic NK cell responses and systemic IFN $\gamma$  responses were unaltered in *Ifnlr1*<sup>-/-</sup> mice. Whether early IL-18 production in the spleen in the absence of robust IL-12 secretion following systemic mCMV infection [41] is responsible for the observation that *Ifnlr1*<sup>-/-</sup> mice mount reduced IFN $\gamma$ <sup>+</sup> NK cell responses in this tissue, is unclear.

IFN $\lambda$ R1 has been shown to directly induce IFN $\gamma$  production by NK cells *in vivo* [31], although whether direct IFN $\lambda$ R signaling occurs in human NK cells and induces IFN $\gamma$  expression remains controversial [44, 45]. In agreement with Souza-Fonseca-Guimaraes et al, we found that IFN $\lambda$  (IL-28A) protein appeared not to directly increase IFN $\gamma$ <sup>+</sup> NK cell development in our studies (Fig 3H), suggesting that IFN $\lambda$  does not act as a co-factor for IL-12 and/or IL-18 induction of IFN $\gamma$ <sup>+</sup> NK cells, at least *in vitro* in the time-frame examined. Furthermore, IFN $\lambda$  failed to induce STAT1 phosphorylation in NK cells derived from naïve or mCMV-infected mice (S1 Fig). Instead, the observation that *Ifnlr1*<sup>-/-</sup> NK cells from naïve mice are less responsive to IL-12 and IL-18 implies that although NK cell repertoires are comparable in WT



and *Ifnlr1*<sup>-/-</sup> mice [31], IFN $\lambda$  may promote NK cell responsiveness to IL-12/IL-18 prior to stimulation with these cytokines. It is feasible that this occurs during NK cell development or in either a different anatomical location or over a different timescale to that studies herein. Alternatively, our data does not preclude the possibility that dysregulated IL-10R2 expression, signaling and/or distribution that may occur in *Ifnlr1*<sup>-/-</sup> mice as a consequence of genetic deletion of this partner of IL-10R2 may impact on NK cell responses.

Overall our data suggest that although IFN $\lambda$  can directly restrict mCMV replication *in vitro*, this is non-essential during the acute and mucosal persistence phases of *in vivo* infection. IFN $\lambda$  predominantly protects mucosal tissue from viral infections [24–28]. Upon mucosal challenge with mCMV, rapid viral dissemination into the spiral ganglia can occur that is independent of local mCMV replication [35]. Thus, initial control of virus replication within mucosal tissue may be unimportant for host protection. Alternatively, given similarities in signaling pathways induced by IFN $\lambda$ R and type I IFN receptor [46], type I IFN may render IFN $\lambda$ R signaling redundant in immune control of mCMV replication *in vivo*. Furthermore, mCMV may blunt direct antiviral activities of IFN $\lambda$ . Indeed, IFN $\lambda$  induces STAT2 activation [46] and the mCMV-encoded pM27 promotes proteasomal degradation of STAT2 [47]. Further, our data does not preclude the possibility that mCMV may interfere with IFN $\lambda$ R abundance and/or localization. Therefore, although IFN $\lambda$  is capable of limiting mCMV replication in *in vitro* assays, it is possible that viral immune evasion mechanisms may blunt this antiviral cytokine pathway *in vivo*.

Rather than acting as an antiviral cytokine during mCMV infection *in vivo*, our data revealed that IFN $\lambda$  exhibits immune modulatory activity by regulating the accumulation of functional NK cells. IFN $\lambda$  did not impact broadly on NK cell responses, in contrast to the induction of NK cell proliferation by IL-28B during influenza infection [48]. Instead, IFN $\lambda$  preferentially promoted the development of IFN $\gamma$ <sup>+</sup> conventional NK cells in the spleen, and our data implied that this occurred via regulation of NK cell responsiveness to IL-12 and, possibly, IL-18. IFN $\lambda$  specifically induced NK cell responses that express IFN $\gamma$  but not granzyme B and did not impact on systemic IFN $\gamma$  secretion. This selective influence on NK cell responses may explain why IFN $\lambda$ R deficiency had no impact on control of virus replication. Overall, our data demonstrate that IFN $\lambda$  modulates conventional NK cell responses in response to systemic virus infection.

## Supporting information

**S1 Fig. IFN $\lambda$  does not induce STAT1 phosphorylation in NK cells from naïve or mCMV-infected mice.** Splenocytes from naïve mice (A&B) or from mice infected (i.p) for 4 days with mCMV (C&D) were stimulated with/without IFN $\beta$  or IFN $\lambda$  and after 1, 3 and 6 hours, STAT1 phosphorylation was measured. (A&C) STAT1 phosphorylation by NK1.1<sup>+</sup>CD3<sup>-</sup> is expressed as Median fluorescent intensity (MFI) and mean + SEM of 5 mice is shown. (B&D) Representative histogram overlays of STAT1 phosphorylation in viable NK1.1<sup>+</sup>CD3<sup>-</sup> cells after 1 hour of stimulation with/without cytokines. Data are representative of 5 separate mice from infected or naïve groups. (TIFF)

## Acknowledgments

The authors are very grateful to Gordon Dougan for facilitating studies at the WTSI.

## Author Contributions

**Conceptualization:** Simon Clare, Ian R. Humphreys.

**Funding acquisition:** Ian R. Humphreys.

**Investigation:** Silvia Gimeno Brias, Jessica Forbester, Mathew Clement, Cordelia Brandt, Katherine Harcourt, Leanne Kane, Simon Clare.

**Methodology:** Silvia Gimeno Brias, Morgan Marsden, Jessica Forbester, Lucy Chapman.

**Project administration:** Morgan Marsden, Simon Clare, Ian R. Humphreys.

**Supervision:** Simon Clare, Ian R. Humphreys.

**Writing – original draft:** Ian R. Humphreys.

**Writing – review & editing:** Silvia Gimeno Brias, Jessica Forbester, Mathew Clement, Simon Clare, Ian R. Humphreys.

## References

1. Britt WJ, and Alford C.A. Cytomegalovirus. In: Knipe DM BN Fa, editor. *Virology*. Philadelphia, PA: Lipincott-Raven Publishers; 1996. p. 2493–523.
2. Biron CA, Tarrio ML. Immunoregulatory cytokine networks: 60 years of learning from murine cytomegalovirus. *Med Microbiol Immunol*. 2015; 204(3):345–54. <https://doi.org/10.1007/s00430-015-0412-3> PMID: 25850988
3. Ouyang W, Rutz S, Crellin NK, Valdez PA, Hymowitz SG. Regulation and functions of the IL-10 family of cytokines in inflammation and disease. *Annu Rev Immunol*. 2011; 29:71–109. <https://doi.org/10.1146/annurev-immunol-031210-101312> PMID: 21166540
4. Kotenko SV, Sacconi S, Izotova LS, Mirochnitchenko OV, Pestka S. Human cytomegalovirus harbors its own unique IL-10 homolog (cmvIL-10). *Proc Natl Acad Sci U S A*. 2000; 97(4):1695–700. PMID: 10677520
5. Stack G, Stacey MA, Humphreys IR. Herpesvirus exploitation of host immune inhibitory pathways. *Viruses*. 2012; 4(8):1182–201. <https://doi.org/10.3390/v4081182> PMID: 23012619
6. Slobedman B, Barry PA, Spencer JV, Avdic S, Abendroth A. Virus-encoded homologs of cellular interleukin-10 and their control of host immune function. *J Virol*. 2009; 83(19):9618–29. <https://doi.org/10.1128/JVI.01098-09> PMID: 19640997
7. Chang WL, Barry PA. Attenuation of innate immunity by cytomegalovirus IL-10 establishes a long-term deficit of adaptive antiviral immunity. *Proc Natl Acad Sci U S A*. 2010; 107(52):22647–52. <https://doi.org/10.1073/pnas.1013794108> PMID: 21149711
8. Krmpotic A, Bubic I, Polic B, Lucin P, Jonjic S. Pathogenesis of murine cytomegalovirus infection. *Microbes Infect*. 2003; 5(13):1263–77. PMID: 14623023
9. Redpath S, Angulo A, Gascoigne NR, Ghazal P. Murine cytomegalovirus infection down-regulates MHC class II expression on macrophages by induction of IL-10. *J Immunol*. 1999; 162(11):6701–7. PMID: 10352288
10. Mandaric S, Walton SM, Rulicke T, Richter K, Girard-Madoux MJ, Clausen BE, et al. IL-10 suppression of NK/DC crosstalk leads to poor priming of MCMV-specific CD4 T cells and prolonged MCMV persistence. *PLoS Pathog*. 2012; 8(8):e1002846. <https://doi.org/10.1371/journal.ppat.1002846> PMID: 22876184
11. Jost NH, Abel S, Hutzler M, Sparwasser T, Zimmermann A, Roers A, et al. Regulatory T cells and T-cell-derived IL-10 interfere with effective anti-cytomegalovirus immune response. *Immunol Cell Biol*. 2014; 92(10):860–71. <https://doi.org/10.1038/icb.2014.62> PMID: 25070242
12. Clement M, Marsden M, Stacey MA, Abdul-Karim J, Gimeno Brias S, Costa Bento D, et al. Cytomegalovirus-Specific IL-10-Producing CD4+ T Cells Are Governed by Type-I IFN-Induced IL-27 and Promote Virus Persistence. *PLoS Pathog*. 2016; 12(12):e1006050. <https://doi.org/10.1371/journal.ppat.1006050> PMID: 27926930
13. Humphreys IR, de Trez C, Kinkade A, Benedict CA, Croft M, Ware CF. Cytomegalovirus exploits IL-10-mediated immune regulation in the salivary glands. *J Exp Med*. 2007; 204(5):1217–25. <https://doi.org/10.1084/jem.20062424> PMID: 17485516
14. Jones M, Ladell K, Wynn KK, Stacey MA, Quigley MF, Gostick E, et al. IL-10 restricts memory T cell inflation during cytomegalovirus infection. *J Immunol*. 2010; 185(6):3583–92. <https://doi.org/10.4049/jimmunol.1001535> PMID: 20713884

15. Lee SH, Kim KS, Fodil-Cornu N, Vidal SM, Biron CA. Activating receptors promote NK cell expansion for maintenance, IL-10 production, and CD8 T cell regulation during viral infection. *J Exp Med*. 2009; 206(10):2235–51. <https://doi.org/10.1084/jem.20082387> PMID: 19720840
16. Stacey MA, Marsden M, Wang EC, Wilkinson GW, Humphreys IR. IL-10 restricts activation-induced death of NK cells during acute murine cytomegalovirus infection. *J Immunol*. 2011; 187(6):2944–52. <https://doi.org/10.4049/jimmunol.1101021> PMID: 21849677
17. Oakley OR, Garvy BA, Humphreys S, Qureshi MH, Pomeroy C. Increased weight loss with reduced viral replication in interleukin-10 knock-out mice infected with murine cytomegalovirus. *Clin Exp Immunol*. 2008; 151(1):155–64. <https://doi.org/10.1111/j.1365-2249.2007.03533.x> PMID: 18005264
18. Stacey MA, Marsden M, Pham NT, Clare S, Dolton G, Stack G, et al. Neutrophils recruited by IL-22 in peripheral tissues function as TRAIL-dependent antiviral effectors against MCMV. *Cell Host Microbe*. 2014; 15(4):471–83. <https://doi.org/10.1016/j.chom.2014.03.003> PMID: 24721575
19. Gimeno Brias S, Stack G, Stacey MA, Redwood AJ, Humphreys IR. The Role of IL-22 in Viral Infections: Paradigms and Paradoxes. *Front Immunol*. 2016; 7:211. <https://doi.org/10.3389/fimmu.2016.00211> PMID: 27303405
20. Lazear HM, Nice TJ, Diamond MS. Interferon-lambda: Immune Functions at Barrier Surfaces and Beyond. *Immunity*. 2015; 43(1):15–28. <https://doi.org/10.1016/j.immuni.2015.07.001> PMID: 26200010
21. Griffiths SJ, Koegl M, Boutell C, Zenner HL, Crump CM, Pica F, et al. A systematic analysis of host factors reveals a Med23-interferon-lambda regulatory axis against herpes simplex virus type 1 replication. *PLoS Pathog*. 2013; 9(8):e1003514. <https://doi.org/10.1371/journal.ppat.1003514> PMID: 23950709
22. Ank N, West H, Bartholdy C, Eriksson K, Thomsen AR, Paludan SR. Lambda interferon (IFN-lambda), a type III IFN, is induced by viruses and IFNs and displays potent antiviral activity against select virus infections in vivo. *J Virol*. 2006; 80(9):4501–9. <https://doi.org/10.1128/JVI.80.9.4501-4509.2006> PMID: 16611910
23. Sommereyns C, Paul S, Staeheli P, Michiels T. IFN-lambda (IFN-lambda) is expressed in a tissue-dependent fashion and primarily acts on epithelial cells in vivo. *PLoS Pathog*. 2008; 4(3):e1000017. <https://doi.org/10.1371/journal.ppat.1000017> PMID: 18369468
24. Nice TJ, Baldrige MT, McCune BT, Norman JM, Lazear HM, Artyomov M, et al. Interferon-lambda cures persistent murine norovirus infection in the absence of adaptive immunity. *Science*. 2015; 347(6219):269–73. <https://doi.org/10.1126/science.1258100> PMID: 25431489
25. Hernandez PP, Mahlakoiv T, Yang I, Schwierzeck V, Nguyen N, Guendel F, et al. Interferon-lambda and interleukin 22 act synergistically for the induction of interferon-stimulated genes and control of rotavirus infection. *Nat Immunol*. 2015; 16(7):698–707. <https://doi.org/10.1038/ni.3180> PMID: 26006013
26. Pott J, Mahlakoiv T, Mordstein M, Duerr CU, Michiels T, Stockinger S, et al. IFN-lambda determines the intestinal epithelial antiviral host defense. *Proc Natl Acad Sci U S A*. 2011; 108(19):7944–9. <https://doi.org/10.1073/pnas.1100552108> PMID: 21518880
27. Mordstein M, Kochs G, Dumoutier L, Renaud JC, Paludan SR, Klucher K, et al. Interferon-lambda contributes to innate immunity of mice against influenza A virus but not against hepatotropic viruses. *PLoS Pathog*. 2008; 4(9):e1000151. <https://doi.org/10.1371/journal.ppat.1000151> PMID: 18787692
28. Davidson S, McCabe TM, Crotta S, Gad HH, Hessel EM, Beinke S, et al. IFNlambda is a potent anti-influenza therapeutic without the inflammatory side effects of IFNalpha treatment. *EMBO Mol Med*. 2016; 8(9):1099–112. <https://doi.org/10.15252/emmm.201606413> PMID: 27520969
29. Skarnes WC, Rosen B, West AP, Koutsourakis M, Bushell W, Iyer V, et al. A conditional knockout resource for the genome-wide study of mouse gene function. *Nature*. 2011; 474(7351):337–42. <https://doi.org/10.1038/nature10163> PMID: 21677750
30. Ding S, Khoury-Hanold W, Iwasaki A, Robek MD. Epigenetic reprogramming of the type III interferon response potentiates antiviral activity and suppresses tumor growth. *PLoS Biol*. 2014; 12(1):e1001758. <https://doi.org/10.1371/journal.pbio.1001758> PMID: 24409098
31. Souza-Fonseca-Guimaraes F, Young A, Mittal D, Martinet L, Bruedigam C, Takeda K, et al. NK cells require IL-28R for optimal in vivo activity. *Proc Natl Acad Sci U S A*. 2015; 112(18):E2376–84. <https://doi.org/10.1073/pnas.1424241112> PMID: 25901316
32. Hermant P, Demarez C, Mahlakoiv T, Staeheli P, Meuleman P, Michiels T. Human but not mouse hepatocytes respond to interferon-lambda in vivo. *PLoS One*. 2014; 9(1):e87906. <https://doi.org/10.1371/journal.pone.0087906> PMID: 24498220
33. Oduro JD, Redeker A, Lemmermann NA, Ebermann L, Marandu TF, Dekhtiarenko I, et al. Murine cytomegalovirus (CMV) infection via the intranasal route offers a robust model of immunity upon mucosal CMV infection. *J Gen Virol*. 2016; 97(1):185–95. <https://doi.org/10.1099/jgv.0.000339> PMID: 26555192
34. Jordan MC. Interstitial pneumonia and subclinical infection after intranasal inoculation of murine cytomegalovirus. *Infect Immun*. 1978; 21(1):275–80. PMID: 213384

35. Farrell HE, Lawler C, Tan CS, MacDonald K, Bruce K, Mach M, et al. Murine Cytomegalovirus Exploits Olfaction To Enter New Hosts. *MBio*. 2016; 7(2):e00251–16. <https://doi.org/10.1128/mBio.00251-16> PMID: 27118588
36. Misumi I, Whitmire JK. IFN-lambda exerts opposing effects on T cell responses depending on the chronicity of the virus infection. *J Immunol*. 2014; 192(8):3596–606. <https://doi.org/10.4049/jimmunol.1301705> PMID: 24646741
37. Stacey MA, Clare S, Clement M, Marsden M, Abdul-Karim J, Kane L, et al. The antiviral restriction factor IFN-induced transmembrane protein 3 prevents cytokine-driven CMV pathogenesis. *J Clin Invest*. 2017; 127(4):1463–74. <https://doi.org/10.1172/JCI84889> PMID: 28240600
38. Blazek K, Eames HL, Weiss M, Byrne AJ, Perocheau D, Pease JE, et al. IFN-lambda resolves inflammation via suppression of neutrophil infiltration and IL-1beta production. *J Exp Med*. 2015; 212(6):845–53. <https://doi.org/10.1084/jem.20140995> PMID: 25941255
39. Broggi A, Tan Y, Granucci F, Zanoni I. IFN-lambda suppresses intestinal inflammation by non-translational regulation of neutrophil function. *Nat Immunol*. 2017; 18(10):1084–93. <https://doi.org/10.1038/ni.3821> PMID: 28846084
40. Sojka DK, Plougastel-Douglas B, Yang L, Pak-Wittel MA, Artyomov MN, Ivanova Y, et al. Tissue-resident natural killer (NK) cells are cell lineages distinct from thymic and conventional splenic NK cells. *Elife*. 2014; 3:e01659. <https://doi.org/10.7554/eLife.01659> PMID: 24714492
41. Pien GC, Sato AR, Takeda K, Akira S, Biron CA. Cutting edge: selective IL-18 requirements for induction of compartmental IFN-gamma responses during viral infection. *J Immunol*. 2000; 165(9):4787–91. PMID: 11046000
42. Orange JS, Biron CA. An absolute and restricted requirement for IL-12 in natural killer cell IFN-gamma production and antiviral defense. *Studies of natural killer and T cell responses in contrasting viral infections*. *J Immunol*. 1996; 156(3):1138–42. PMID: 8557990
43. Madera S, Sun JC. Cutting edge: stage-specific requirement of IL-18 for antiviral NK cell expansion. *J Immunol*. 2015; 194(4):1408–12. <https://doi.org/10.4049/jimmunol.1402001> PMID: 25589075
44. Dring MM, Morrison MH, McSharry BP, Guinan KJ, Hagan R, Irish HCVR, et al. Innate immune genes synergize to predict increased risk of chronic disease in hepatitis C virus infection. *Proc Natl Acad Sci U S A*. 2011; 108(14):5736–41. <https://doi.org/10.1073/pnas.1016358108> PMID: 21402922
45. Kramer B, Eisenhardt M, Glassner A, Korner C, Sauerbruch T, Spengler U, et al. Do lambda-IFNs IL28A and IL28B act on human natural killer cells? *Proc Natl Acad Sci U S A*. 2011; 108(34):E519–20; author reply E21–2. <https://doi.org/10.1073/pnas.1108850108> PMID: 21825163
46. Zhou Z, Hamming OJ, Ank N, Paludan SR, Nielsen AL, Hartmann R. Type III interferon (IFN) induces a type I IFN-like response in a restricted subset of cells through signaling pathways involving both the Jak-STAT pathway and the mitogen-activated protein kinases. *J Virol*. 2007; 81(14):7749–58. <https://doi.org/10.1128/JVI.02438-06> PMID: 17507495
47. Trilling M, Le VT, Fiedler M, Zimmermann A, Bleifuss E, Hengel H. Identification of DNA-damage DNA-binding protein 1 as a conditional essential factor for cytomegalovirus replication in interferon-gamma-stimulated cells. *PLoS Pathog*. 2011; 7(6):e1002069. <https://doi.org/10.1371/journal.ppat.1002069> PMID: 21698215
48. Wang Y, Li T, Chen Y, Wei H, Sun R, Tian Z. Involvement of NK Cells in IL-28B-Mediated Immunity against Influenza Virus Infection. *J Immunol*. 2017; 199(3):1012–20. <https://doi.org/10.4049/jimmunol.1601430> PMID: 28637903