

CORRECTION

Correction: Cost and time-effective method for multi-scale measures of rugosity, fractal dimension, and vector dispersion from coral reef 3D models

G. C. Young, S. Dey, A. D. Rogers, D. Exton

[Fig 6](#) is incorrect. The authors have provided a corrected version here.



OPEN ACCESS

Citation: Young GC, Dey S, Rogers AD, Exton D (2018) Correction: Cost and time-effective method for multi-scale measures of rugosity, fractal dimension, and vector dispersion from coral reef 3D models. PLoS ONE 13(7): e0201847. <https://doi.org/10.1371/journal.pone.0201847>

Published: July 31, 2018

Copyright: © 2018 Young et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Fig 6. Accuracy of 3D model in terms of point-to-point distances. The root mean square errors (RMSE) of our models were 1.48 cm in X-Y and 1.35 cm in Z, with models underestimating dimensions in both X-Y and Z.

<https://doi.org/10.1371/journal.pone.0201847.g001>

Reference

1. Young GC, Dey S, Rogers AD, Exton D (2017) Cost and time-effective method for multi-scale measures of rugosity, fractal dimension, and vector dispersion from coral reef 3D models. *PLoS ONE* 12(4): e0175341. <https://doi.org/10.1371/journal.pone.0175341> PMID: 28406937