

CORRECTION

Correction: Impact of clinical severity of stroke on the severity and recovery of visuospatial neglect

The *PLOS ONE* Staff

Notice of republication

This article was republished on July 30, 2018, to correct for errors in the Data Availability statement introduced during the production process. The publisher apologizes for the errors. Please download this article again to view the correct version. The originally published, uncorrected article and the republished, corrected articles are provided here for reference.

Supporting information

S1 File. Originally published, uncorrected article.


(PDF)

S2 File. Republished, corrected article.

(PDF)

Reference

1. Nijboer TCW, Winters C, Kollen BJ, Kwakkel G (2018) Impact of clinical severity of stroke on the severity and recovery of visuospatial neglect. *PLoS ONE* 13(7): e0198755. <https://doi.org/10.1371/journal.pone.0198755> PMID: 29966012



OPEN ACCESS

Citation: The *PLOS ONE* Staff (2018) Correction: Impact of clinical severity of stroke on the severity and recovery of visuospatial neglect. *PLoS ONE* 13(8): e0202434. <https://doi.org/10.1371/journal.pone.0202434>

Published: August 10, 2018

Copyright: © 2018 The PLOS ONE Staff. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.