

RESEARCH ARTICLE

Multi-informant path models of the influence of psychosocial and treatment-related variables on adherence and metabolic control in adolescents with type 1 diabetes mellitus

Lene Juel Kristensen^{1*}, Niels Holtum Birkebaek², Anne Hvarregaard Mose², Morten Berg Jensen³, Mikael Thastum¹

1 Department of Psychology and Behavioural Sciences, Aarhus University, Aarhus, Denmark, **2** Department of Pediatrics, Aarhus University Hospital, Aarhus N., Denmark, **3** Department of Economics and Business Economics, Aarhus University, Aarhus, Denmark

* lene_juelk@hotmail.com



OPEN ACCESS

Citation: Kristensen LJ, Birkebaek NH, Mose AH, Berg Jensen M, Thastum M (2018) Multi-informant path models of the influence of psychosocial and treatment-related variables on adherence and metabolic control in adolescents with type 1 diabetes mellitus. PLoS ONE 13(9): e0204176. <https://doi.org/10.1371/journal.pone.0204176>

Editor: David Meyre, McMaster University, CANADA

Received: June 7, 2018

Accepted: September 3, 2018

Published: September 20, 2018

Copyright: © 2018 Kristensen et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data are from the Danish National Psychosocial Diabetes Survey whose authors may be contacted. The data include person sensitive information and are protected by the Danish national Data Protection Agency. For further information regarding how to obtain a de-identified, and select portion of the data, contact either the corresponding author or the Department of Psychology and Behavioural Sciences, Aarhus University (psykologi@psy.au.dk).

Abstract

Background

We assessed the associations between metabolic control and adherence and a broad range of adolescent and family characteristics (e.g., gender, family structure), treatment-related variables (e.g., disease duration, treatment modality), and psychosocial factors (e.g., symptoms of depression and anxiety, parental support, self-efficacy) in a nationwide study of Danish adolescents (age 12–17 years) with type 1 diabetes mellitus (T1DM).

Methods

Sixty-four percent of invited families participated by completing a survey and providing a blood sample. Two path models of associations between generic and diabetes-related family factors, adolescent self-efficacy, emotional difficulties, and metabolic control and adherence were tested, one for adolescents and one for caregivers. Demographic variables were included as covariates.

Results

Both path models demonstrated a satisfying model fit. In both models, metabolic control was associated with adherence, age, and T1DM duration. In the adolescent model, metabolic control was also related to treatment modality, single-parent household, caregiver non-support, and anxiety, whereas in the caregiver model metabolic control was associated with family conflict and caregiver support. In both models, adherence was related to age, duration, treatment modality, family conflict, caregiver support, family functioning, and emotional difficulties of the adolescent. In the adolescent model, adherence was also related to adolescent self-efficacy, whereas in the caregiver model adherence was associated with adolescent gender and caregiver non-support and support. Adolescent self-efficacy, emotional well-being, and difficulties related to adolescent/caregiver interaction appeared to be

Funding: Novo Nordisk A/S (<https://www.novonordisk.com/>) supported the work ('Psychosocial aspects of diabetes' - Grant number: 494887). The financial support was granted to MT. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: Novo Nordisk A/S supported the work financially ('Psychosocial aspects of diabetes' - Grant number: 494887). There are no patents, products in development or marketed products to declare. This does not alter our adherence to all the PLOS ONE policies on sharing data and materials, as detailed online in the guide for authors. We have no other competing interests to declare.

particularly important, as indicated by their stronger association with adherence and/or metabolic control.

Conclusion

The results highlight the value of applying a multi-informant approach to address the psychosocial well-being of adolescents with diabetes in a large national sample. Self-efficacy, emotional, and family-related difficulties are important aspects to address in both clinical care and future research regarding adolescents with T1DM.

Introduction

Achieving adequate metabolic control is crucial in children and adolescents with type 1 diabetes mellitus (T1DM) to prevent both immediate and long-term health complications [1, 2]. The many tasks necessary to achieve successful diabetes management is perceived by many patients and caregivers as a huge burden, with adherence to diabetes treatment a necessary, but not always sufficient, prerequisite for optimal control [3]. Biological, physiological, and psychosocial factors act as intricate parts of the complex processes that influence metabolic control and adherence in adolescents with T1DM.

A range of variables related to individual and family characteristics have been shown to affect T1DM treatment. Older age, longer disease duration, and being treated with multiple daily injections (MDI) vs. insulin pump/continuous subcutaneous insulin infusion (CSII), living in a single-parent household, lower education level of parents, and lower family income have all been associated with decreased adherence and worsening metabolic control [4–7]. However, any association of gender with either metabolic control or adherence is unclear [8–10].

The prevalence of problematic psychological responses by adolescents with T1DM, such as symptoms of depression and anxiety, differs greatly among studies [11–14] but, regardless of the overall prevalence, the presence of these symptoms should warrant attention. Some studies have shown an association between deteriorating metabolic control and mental health problems [15, 16], and adherence has been associated with depression [17, 18] and anxiety [16]. Other studies have not been able to confirm these associations [19].

Previous research has consistently found adolescent self-efficacy (defined as the belief that one can carry out specific behaviors in specified situations [20]) in relation to diabetes management to be an important factor in diabetes care and to be associated not only with adherence, but also metabolic control [21–23], possibly with self-efficacy acting as a mediator between diabetes-specific family issues (e.g., diabetes-related conflict, non-support, or diabetes responsibility) and adherence or metabolic control [24, 25]. A number of studies have focused on conflict and arguments in the family related to diabetes-specific tasks and behaviors and found a direct association with poorer metabolic control [26–28]. Other studies have found an association between conflict and metabolic control to be mediated by adherence [29] or found no association between diabetes-related family conflict and adherence [30].

Both parental support and non-support (as evidenced by critical and negative parenting) in relation to diabetes-specific tasks and behaviors have been shown to be associated with both adherence and metabolic control [31, 32], possibly with adherence mediating the association between critical parenting and glycosylated hemoglobin (HbA_{1c}) as the prevailing indicator of metabolic control [33]. However, further investigations of the link between support, non-support, adherence, and metabolic control is needed due to the so far inconsistent results.

Parental involvement in relation to responsibility in diabetes management tasks has been associated with adherence, and possibly metabolic control through the mediational effect of adherence [34, 35]. A previous study found that adolescents who perceive greater caregiver responsibility engage in better diabetes management, as measured by blood glucose monitoring [36]. Aspects of the family's general way of functioning, such as family cohesion and family dysfunction, have been associated with adherence and metabolic control in some studies [37, 38], whereas other studies have found no relationship with metabolic control [39].

Many of the abovementioned studies relied on relatively small, homogenous groups of participants, and larger studies that are able to test comprehensive models reflecting the multifaceted complexity of factors influencing daily diabetes management, and thereby metabolic control, are needed. Models including a larger number of variables, while still seeking to describe the relative and unique contribution of each, are necessary to guide the multidisciplinary clinical care of adolescents with T1DM and the development of effective screening and interventions. Whittemore, Jaser, Guo and Grey [40] have proposed a theoretical model based on results of previous studies that encompass the possible associations of a multitude of variables relating to individual and family characteristics, psychosocial, individual, and family responses, with adaptation outcomes such as metabolic control. This model is intended as a conceptual framework to guide researchers and health care providers in their understanding of the complexity of diabetes treatment and adaptation to living with this chronic disease. Whittemore et al. stresses the importance of conducting further research to confirm and develop the model and the inconsistent results on which some of the presumed association are based. This model has inspired the development of the path models being tested in the current study.

Thus, the present study set out to investigate possible associations between adherence or metabolic control and patient characteristics, treatment aspects, and psychosocial and psychological variables in Danish children and adolescents (age 12–17 years) with T1DM. As previous research has highlighted the often differing perspectives of adolescents and caregivers in assessing health and the family milieu, we tested separate, independent multivariate path models for children/adolescents and caregivers in order to examine possible direct and indirect pathways between a broad range of variables and metabolic control and adherence. Based on previous research, poorer metabolic control and lower levels of adherence were expected to be independently associated with more diabetes-related family conflict, more parental non-support, and less parental support. These outcomes were also expected to be related to less adolescent self-efficacy in relation to diabetes care and more symptoms of depression and anxiety, and family division of responsibility and general family functioning were expected to be related to adherence, which would act as a mediator between metabolic control and emotional and social difficulties, diabetes-related family conflict, parental support, and non-support. Furthermore, the possible influence of age, gender, diabetes duration, treatment modality, and caregiver socioeconomic factors in relation to both adherence and metabolic control was assessed.

Materials and methods

Ethics statement

The department where the study was conducted did not have an institutional review board, as this is not standard in Denmark. Thus, in accordance with Danish procedure, the regional ethic committees (De Videnskabetiske Komitéer for Region Midtjylland) was consulted. In keeping with the regulations of the committees, questionnaire-based studies do not require permission prior to initiation; however, a study protocol was provided to the committees who

confirmed, that although participating children and adolescents were asked to submit a small blood sample (comparable to their daily blood glucose testings), no biological samples were collected with the intent of establishing a research bio-bank, therefore the project was not encompassed by the term 'Bio-medical research', and as such not eligible for Committee review and approval. The project was registered with the Danish Data Protection Agency (Ref no. 2013-41-1528). Written informed consent was obtained from all participants or their caregivers through the online version of the questionnaires or on paper. All families were given thorough written information informing them that their participation was voluntary and anonymous to everyone other than the first author, and that their consent could be withdrawn at any time, just as refusal to participate did not in any way influence the diabetes treatment that the child/adolescent was receiving.

Participants and procedure

The present study was part of a nationwide web survey initiated to assess the influence of psychosocial variables on adherence, metabolic control, and quality of life in all Danish children and adolescents with T1DM (age 2–17 years). The study was conducted in collaboration with the Danish Society for Diabetes in Childhood and Adolescence, who administers the Danish Registry for Childhood and Adolescent Diabetes (DanDiabKids). Since 1996, DanDiabKids has collected data on all children and adolescents in Denmark with a diagnosis of T1DM, including annual registration of current HbA_{1c} levels, which are analyzed centrally to ensure conformity.

Based on information from DanDiabKids, all families in Denmark with a child or adolescent between 2 and 17 years of age with a diagnosis of T1DM ($n = 1739$) were invited to participate. We excluded 258 families who were registered as being unwilling to participate in scientific research, had a protected address, or were no longer residing at the address registered in the Danish Civil Registration System from which all participant addresses were collected. All families received a written invitation by post, asking them to participate in the national web survey. They were also given the option of completing a paper version of the questionnaire if they preferred.

The caregiver primarily involved with the diabetes-related care of the child/adolescent was requested to complete the survey. All families were asked to provide a blood sample from the child for HbA_{1c} analysis. A total of 1075 of the invited families had a child/adolescent (hereafter referred to as 'adolescents') with T1DM between 12 and 17 years of age. Based on their age, these adolescents were deemed mature enough to complete the full self-report questionnaire battery by themselves if desiring to do so. 519 of these adolescents, and 531 of the caregivers completed the required questionnaires, and consequently, data from these adolescents and one of their caregivers form the basis of the present study and analysis (see [S1 Fig](#)).

A paired samples t-test revealed no significant differences between the HbA_{1c} values provided for the study ($M = 8.23$, $SD = 1.24$) and those obtained from DanDiabKids for the same participants ($M = 8.21$, $SD = 1.28$, $t(579) = 0.65$, $p = 0.52$). The HbA_{1c} values from DanDiabKids were used for the analyses if participants did not provide a blood sample for the study.

Patient characteristics, including family structure and Danish as primary language in the home (a proxy marker for ethnicity), are summarized in [Table 1](#).

Independent samples t-tests revealed no significant difference between participants and non-participants regarding age ($t(1165) = 1.22$, $p = 0.22$), but non-participants had been diagnosed with T1DM for a slightly longer duration ($M = 6.83$, $SD = 1.61$, $t(1160) = 3.71$, $p = 0.00$) and were in worse metabolic control ($M = 8.89$, $SD = 1.53$, $t(1153) = 7.29$, $p = 0.00$). Of the participating adolescents, 26.7% met the recommended HbA_{1c} level ($< 7.5\%/58$ mmol/mol).

Table 1. Participant characteristics.

	Percentage or mean (standard deviation)	Adolescent report	Parent report
Age of participating adolescents, years	14.6 (1.6)		
Gender of participating adolescents, girls	50.0%		
HbA1c - %	8.25 (1.25)		
HbA1c - mmol/mol	66.7 (13.7)		
Diabetes duration, years	6.07 (3.47)		
Participants using insulin pump	37.8%		
Gender of participating caregiver, female	83.6%		
Living with parents who are no longer together	20.8%		
No longer living with either parent (living with foster parents, at a continuation school, a treatment facility, or living by themselves)	5.2%		
Danish primary language at home	98.7%		
Household income:			
• Less than 400.000 kroner*	10.5%		
• Between 400.000–499.999 kroner*	12.8%		
• 500.000 kroner or more*	76.7%		
Adherence (ADQ)		3.95 (0.58)	3.96 (0.62)
Anxiety (BAI)		8.11 (7.78)	
Depression (BDI)		7.58 (8.85)	
Emotional and behavioral problems (SDQ)			7.48 (5.81)
Self-efficacy (SEDM)		7.19 (1.71)	
General family functioning (FAD)		20.70 (5.66)	20.61 (5.65)
Diabetes-related family conflict (DFCS)		20.57 (5.25)	20.46 (4.65)
Supportive caregiver behavior (DFBC sup)		26.35 (6.50)	27.94 (5.54)
Non-supportive caregiver behavior (DFBC non-sup)		14.62 (5.05)	14.38 (4.50)
Responsibility for diabetes-related tasks (DFRQ)		33.52 (4.82)	31.96 (5.03)

* Annual household income before taxes.

<https://doi.org/10.1371/journal.pone.0204176.t001>

Measures

Treatment adherence. Adherence to diabetes treatment was assessed using the Adherence in Diabetes Questionnaire (ADQ), which was developed for this study. The psychometric properties of the questionnaire were described previously [41]. Caregivers and adolescents completed the 17 or 19 items (depending on treatment modality) of the ADQ assessing adherence to different aspects of diabetes treatment. The questionnaires were scored by calculating the mean of all items. Higher scores indicate better adherence. Within this study sample, the ADQ demonstrated acceptable internal consistency, with Cronbach’s alpha ranging from 0.89 (MDI version) to 0.85 (CSII version) on the caregivers’ responses, and 0.85 (MDI version) to 0.82 (CSII version) on the adolescents’ reports.

Supportive and non-supportive caregiver behavior. Parental support and non-support in relation to diabetes care were assessed using the Diabetes Family Behavior Checklist (DFBC) [42, 43]. Both caregivers and adolescents completed the questionnaire, which consists of two separate subscales: nine items comprising the support scale assessing both affective and practical support, and seven items comprising the non-support scale assessing diabetes-related critical parenting behavior. Higher scores on the support scale indicate the child’s or caregiver’s perception of more parental support in relation to diabetes, and higher scores on the non-support scale indicate a perception of more critical parenting/non-support. The reliability and

internal consistency of the subscales of the DFBC were previously found to be adequate [31, 44]. In the present study, the Cronbach's alpha was 0.69 (caregivers)/0.74 (adolescents) for the support subscale and 0.70 (caregivers)/0.64 (adolescents) for the non-support subscale.

Diabetes-related family conflict. The presence of diabetes-related family conflict was assessed using the Diabetes Family Conflict Scale (DFCS). Both adolescents and caregivers completed the revised version of the DFCS, which has demonstrated satisfactory internal consistency, adequate concurrent validity, and predictive validity in relation to metabolic control [26]. The DFCS is scored by calculating the sum score for all items, with higher scores indicating more conflict. The Cronbach's alpha was 0.87.

Responsibility for diabetes-related tasks. Both caregivers and adolescents completed the Diabetes Family Responsibility Questionnaire (DFRQ) to assess the division of responsibility in relation to regimen tasks, general health maintenance, and social presentation of diabetes [45]. This instrument is widely used and has proven to be psychometrically sound [36]. In the present study, a total responsibility score was calculated by summing the item responses, with a higher score indicating more adolescent responsibility. The Cronbach's alpha was 0.84 for the caregiver scale and 0.82 for the adolescent scale.

General family functioning. The General Functioning subscale of the Family Assessment Device (FAD) was chosen as a measure of generic family functioning. The FAD is based on the McMaster Model of Family Functioning [46] and comprises six subscales describing six dimensions of family functioning. The 12-item General Functioning subscale was completed by both caregivers and adolescents, with a higher score indicating unhealthier functioning. The psychometric properties of both the overall questionnaire and the General Functioning subscale were described previously and deemed satisfactory [46, 47], just as the FAD has been used extensively in pediatric samples [48].

The Cronbach's alpha of the subscale in the present study was 0.89 for caregivers and 0.87 for adolescents.

Social and emotional difficulties of the adolescent. Adolescents completed the depression (BDI) and anxiety (BAI) subscales of the Beck's Youth Inventories—second edition (BYI-II) [49]. Each subscale consists of 20 questions and is scored by calculating a total score ranging from 0 to 60, with higher scores indicating more symptoms of depression or anxiety. The reliability and test-retest stability of the Danish version of the BYI was found to be adequate [50]. In the present study, the Cronbach's alpha was 0.94 for the BDI subscale and 0.92 for the BAI subscale.

The caregivers completed the Strength and Difficulties Questionnaire (SDQ). The SDQ is a brief, 25-item behavioral screening instrument consisting of five separate subscales that generate scores for Emotional Symptoms, Conduct Problems, Hyperactivity-Inattention, Peer Problems, and Prosocial Behavior [51]. Only the Total Difficulties Score calculated by summing the scores on the first four subscales was used in the present study. The SDQ has been used worldwide, with satisfactory psychometric properties, including reliability and validity, being established [52, 53]. In this study, the Cronbach's alpha for the 20 items of the Total Difficulties Score was 0.84.

Self-efficacy. The adolescents' self-efficacy in relation to managing diabetes-related tasks was measured using the Self-Efficacy for Diabetes Self-Management (SEDM). The 10-item questionnaire is scored by calculating the mean of all items. Higher scores indicate a more positive perception of self-efficacy. The SEDM was previously shown to be both valid and reliable [54], and in this study the Cronbach's alpha was 0.89.

Medical and sociodemographic information. Information regarding diabetes duration was provided by DanDiabKids, whereas caregivers provided information regarding family structure, caregiver education level, household income, and the adolescent's current diabetes treatment.

Analyzing the blood samples provided by participants, HbA_{1c} was used as an objective measure of metabolic control over the most recent 8–12 weeks. Blood samples were analyzed at a central laboratory using high-pressure liquid chromatography (Tosoh Bioscience, South San Francisco, CA, USA) and standardized according to the American National Glycohemoglobin Standardization Program (NGSP).

The web-based survey took approximately 45 minutes to complete depending on reading proficiency and computer skills.

Statistical analysis

Descriptive analyses were performed in SPSS version 24 (SPSS Inc., 2016).

Bivariate correlations between observed measures, including demographic covariates and metabolic control, were examined. In line with Cohen [55], we considered a correlation of 0.5 as large, 0.3 as moderate, and 0.1 as small. Path modeling was performed using Mplus 7.0 software [56].

Mediation was assessed using bias-corrected bootstrapped confidence intervals for the indirect effects as advocated by Shrout and Bolger [57]. Thus, we followed recent recommendations and did not use significance of the total effect as a prerequisite for assessing mediation [57]. The hypothesized models consisted of 10 and 8 variables for adolescents and caregivers, respectively. The analyses were carried out separately for adolescents and caregivers.

Data were screened for outliers and assessed for normality. Diabetes-related family conflict and the social and emotional difficulties of the adolescent exhibited a pronounced pattern of non-normality (right-skewed); therefore, we carried out the analysis using the robust maximum likelihood estimator, providing standard errors and measures of model fit, which are robust to non-normality. In the path analysis missing data was handled via maximum-likelihood for the endogenous variables. For the exogenous variables list-wise deletion is utilized. The latter leads to a reduction in the sample size of slightly below 25% for both children and adults (see S1 Fig). The overall model fit was assessed using the chi-square value with $p > 0.05$, root mean square error of approximation (RMSEA) < 0.05 , and standardized root mean square residual (SRMR) < 0.08 , indicating an acceptable fit of the model. Furthermore, a comparative fit index (CFI) and Tucker-Lewis Index (TLI) in excess of 0.95 indicates an acceptable fit of the model [58].

Results

Bivariate correlations

Correlations between study variables. Bivariate correlations between all included variables were examined (Table 2), mostly confirming the above-stated hypotheses showing small to large correlations. However, no significant associations were found between metabolic control and adolescent or caregiver Diabetes Family Responsibility Questionnaire scores, or between metabolic control and gender, adherence and gender regardless of respondent, or caregivers' perception of adherence and caregiver education. In addition, caregiver perception of adherence was unrelated to their perception of supportive behavior in relation to diabetes care.

Parent-adolescent correlations. Significant correlations were found between caregiver and adolescent reports for all measures, indicating relatively high agreement between the respondents. Large correlations were found for measures of adherence ($r = 0.61$), parental diabetes-related non-support ($r = 0.53$), and diabetes family responsibility ($r = 0.57$). Medium correlations were found for general family functioning ($r = 0.47$), diabetes-related family conflict ($r = 0.45$), and parental diabetes-related support ($r = 0.45$). In addition, medium correlations were found between parental Strength and Difficulties Questionnaire reports and

Table 2. Intercorrelations between health outcomes, psychosocial variables, and patient characteristics.

	HbA1c	ADQ (A)	ADQ (C)	BDI (A)	BAI (A)	SDQ (C)	SEDQ (A)	FAD (A)	FAD (C)	DFCS (A)	DFCS (C)	DFBC sup. (A)	DFBC sup. (C)	DFBC non-sup. (A)	DFBC non-sup. (C)	DFRQ (A)	DFRQ (C)	Age	Gender	Diabetes duration (years)	Care-giver education level ^a	Family structure	Insulin pump
HbA1c	1.00	-0.37**	-0.41**	0.17**	0.09*	0.25**	-0.27**	0.16**	0.19**	0.28**	0.35**	-0.10**	-0.13**	0.24**	0.24**	0.04	0.03	0.19**	-0.02	0.19**	-0.09*	0.13**	-0.12**
ADQ (A)		1.00	0.61**	-0.24**	-0.14**	0.62**	-0.38**	-0.31**	-0.37**	-0.39**	-0.39**	0.35**	0.23**	-0.29**	-0.32**	0.11**	-0.02	-0.27**	-0.07	-0.17**	0.11*	-0.13**	0.17**
ADQ (C)			1.00	-0.23**	-0.14**	0.52**	-0.27**	-0.39**	-0.40**	-0.55**	0.15**	0.08	0.08	-0.37**	-0.51**	0.15**	0.18**	-0.15**	-0.06	-0.15**	0.04	-0.16**	0.23**
BDI (A)				1.00	0.84**	0.44**	0.37**	0.22**	0.37**	0.18**	0.18**	-0.15**	0.03	0.27**	0.18**	0.00	0.05	0.08*	-0.22**	0.00	-0.11**	0.08	-0.02
BAI (A)					1.00	0.38**	0.32**	0.16**	0.34**	0.12**	0.12**	-0.12**	0.08	0.21**	0.13**	-0.02	0.02	0.04	-0.23**	-0.04	-0.10*	0.05	-0.02
SDQ (C)						1.00	-0.38**	0.42**	-0.29**	0.43**	0.43**	0.05	0.05	0.31**	0.40**	-0.19**	-0.30**	-0.03	0.13**	0.06	-0.14**	0.13**	-0.19**
SEDQ (A)							1.00	0.29**	-0.40**	-0.34**	0.22**	0.04	-0.32**	-0.32**	0.31**	0.21**	0.21**	-0.00	0.03	-0.05	0.11**	-0.12**	0.09*
FAD (A)								1.00	0.47**	0.28**	-0.33**	-0.15**	-0.11**	0.22**	0.22**	-0.03	-0.09*	0.13**	0.03	0.04	-0.10*	0.11**	0.03
FAD (C)									1.00	0.26**	0.29**	-0.11**	-0.11**	0.26**	0.32**	-0.07	-0.13**	0.05	0.07	0.12**	-0.14**	0.05	-0.14**
DFCS (A)										1.00	0.45**	-0.04	0.01	0.49**	0.34**	-0.21**	-0.12**	0.05	-0.07	0.06	-0.09*	0.07	0.18**
DFCS (C)											1.00	-0.04	0.12**	0.65**	-0.21**	-0.27**	-0.27**	0.01	0.12**	0.11**	-0.09*	0.04	-0.07
DFBC sup. (A)												1.00	0.45**	0.09*	0.05	-0.23**	-0.25**	-0.34**	0.08	-0.10*	-0.01	-0.01	0.04
DFBC sup. (C)													1.00	0.07	0.19**	-0.31**	-0.37**	-0.40**	0.01	-0.12**	-0.03	-0.01	0.11**
DFBC non-sup. (A)														1.00	0.53**	-0.08*	-0.20**	0.03	0.06	0.06	-0.07	0.03	-0.07
DFBC non-sup. (C)															1.00	-0.16**	-0.26**	-0.07	0.09*	0.05	-0.07	-0.11**	-0.08*
DFRQ (A)																1.00	0.57**	0.48**	-0.13**	0.05	-0.06	-0.09*	-0.04
DFRQ (C)																	1.00	0.51**	-0.17**	0.10*	0.02	-0.07	-0.05
Age																		1.00	0.02	0.19**	-0.05	0.02	-0.15**
Gender																			-1.00	0.03	0.01	0.03	-0.16**
Diabetes duration (years)																				1.00	0.01	-0.02	0.16**
Care-giver education level ^a																					1.00	-0.04	0.03
Family structure																						1.00	-0.07
Insulin pump																							1.00

HbA1c, hemoglobin A1c; ADQ, Adherence in Diabetes Questionnaire; BDI, Depression subscale of the Beck's Youth Inventories-Second edition; SDQ, Strength and Difficulties Questionnaire; SEDM, Self-efficacy for Diabetes Self-Management; FAD, Family Assessment Device; DFCS, Diabetes Family Conflict Scale; DFBC sup., Diabetes Family Behavior Checklist-caregiver support subscale; DFBC non-sup., Diabetes Family Behavior Checklist-caregiver non-support subscale; DFRQ, Diabetes Family Responsibility Questionnaire. (A) indicates adolescent self-report, (C) indicates caregiver report.

^a Higher scores indicate higher levels of education.

** p < 0.01

* p < 0.05

<https://doi.org/10.1371/journal.pone.0204176.t002>

adolescents' self-reports on the Beck's Depression Inventory ($r = 0.44$) and Beck's Anxiety Inventory ($r = 0.38$).

Path modeling

Two structural equation models of associations between adherence and metabolic control and diabetes-related family conflict, supportive and non-supportive caregiver behavior, responsibility for diabetes-related tasks, general family functioning, self-efficacy (only in the adolescent model), and social and emotional difficulties of the adolescent were tested, including the possible mediational role of adherence. Demographic variables were included as covariates.

Adolescent path model. For adolescents, the model fit was satisfactory [$X^2(2) = 3.19$, $p = 0.20$, CFI = 0.997, TLI = 0.955, SRMR = 0.004, RMSEA = 0.034] (Fig 1).

Higher non-supportive caregiver behavior ($\beta = 0.11$, $p = 0.031$), fewer symptoms of anxiety ($\beta = -0.17$, $p = 0.017$), and poorer adherence ($\beta = -0.21$, $p = 0.001$) were associated with poorer metabolic control. Less diabetes-related conflict ($\beta = -0.11$, $p = 0.007$), more supportive caregiver behavior ($\beta = 0.16$, $p = 0.000$), higher self-efficacy ($\beta = 0.44$, $p = 0.000$), lower family functioning (less unhealthy functioning) ($\beta = -0.10$, $p = 0.011$), fewer depressive symptoms ($\beta = -0.13$, $p = 0.038$), and more symptoms of anxiety ($\beta = 0.23$, $p = 0.000$) were associated with better adherence.

Longer diabetes duration ($\beta = 0.16$, $p = 0.000$), use of multiple daily injections vs. insulin pump ($\beta = -0.10$, $p = 0.006$), and living alone with one parent vs. living with both parents ($\beta = 0.10$, $p = 0.030$) were associated with poorer metabolic control. Lower age ($\beta = -0.18$, $p = 0.000$), shorter diabetes duration ($\beta = -0.10$, $p = 0.003$), and using an insulin pump vs. multiple daily injections ($\beta = 0.11$, $p = 0.000$) were associated with better adherence.

Finally, adherence fully mediated the relationship between metabolic control and diabetes-related family conflict, supportive caregiver behavior, and self-efficacy with bias-corrected bootstrapped confidence intervals (CIs) of [0.000; 0.043], [-0.057; -0.009], and [-0.149; -0.033], respectively. Adherence acted as a partial mediator for the relation between anxiety and metabolic control (CI [-0.084; -0.010]).

Overall, the proposed adolescent path model tested here accounted for 25% of the variance in metabolic control and 51% of the variance in adherence. Metabolic control had the strongest association with adherence ($r = -0.21$), and symptoms of anxiety ($r = -0.17$), whereas the variables most strongly associated with adherence were self-efficacy ($r = 0.44$) and symptoms of anxiety ($r = 0.23$).

Caregiver path model. For caregivers, the model fit was satisfactory [$X^2(2) = 1.78$, $p = 0.41$, CFI = 1.000, TLI = 1.008, SRMR = 0.003, RMSEA = 0.000] (Fig 2).

Higher diabetes-related conflict ($\beta = 0.22$, $p = 0.001$), less supportive caregiver behavior ($\beta = -0.11$, $p = 0.013$), and poorer adherence ($\beta = -0.24$, $p = 0.001$) were associated with poorer metabolic control. Less diabetes-related conflict ($\beta = -0.14$, $p = 0.003$), less non-supportive caregiver behavior ($\beta = -0.30$, $p = 0.000$), higher responsibility for diabetes-related tasks placed on the youth ($\beta = 0.17$, $p = 0.006$), more supportive caregiver behavior ($\beta = 0.12$, $p = 0.001$), and fewer social and emotional difficulties ($\beta = -0.20$, $p = 0.000$) were associated with better adherence. The higher the age of the adolescent ($\beta = 0.13$, $p = 0.003$) and the longer the duration of diabetes ($\beta = 0.08$, $p = 0.027$), the poorer the metabolic control, but lower age ($\beta = -0.12$, $p = 0.002$), being male ($\beta = 0.09$, $p = 0.009$), shorter duration of diabetes ($\beta = -0.07$, $p = 0.022$), and using an insulin pump vs. multiple daily injections ($\beta = 0.16$, $p = 0.000$) were associated with better adherence.

Finally, adherence fully mediated the relationship between metabolic control and lower non-supportive caregiver behavior and social and emotional difficulties (CIs: [0.021; 0.121]).




Fig 1. Standardized path coefficients-adolescent path model.

<https://doi.org/10.1371/journal.pone.0204176.g001>

and [0.011; 0.084], respectively). Adherence partially mediated the relationship between metabolic control and diabetes-related family conflict, as well as supportive caregiver behavior (CIs: [0.004; 0.064] and [0.054; 0.002], respectively). With a slightly better model fit for the caregiver model than the adolescent model, the explanatory value regarding explained variances revealed substantial resemblance between the two models, with the caregiver path model accounting for 26% of the variance in HbA_{1c} and 49% of the variance in adherence. The variables most strongly associated with metabolic control were adherence ($r = -0.24$) and diabetes-related family conflict ($r = 0.22$), whereas adherence was most associated with parental non-support ($r = -0.30$) and the total Strength and Difficulties Questionnaire score ($r = -0.20$), reflecting the parental assessment of the social and emotional difficulties of the adolescent.




Fig 2. Standardized path coefficients—caregiver path model.

<https://doi.org/10.1371/journal.pone.0204176.g002>

Discussion

The present study investigated associations between patient characteristics, treatment aspects, and psychosocial and psychological variables on diabetes management and metabolic control. The associations between variables were assessed in two separate path models, one for adolescents and one for caregivers, and both had satisfactory fit.

The strengths of this study include the large and nationally representative sample size, multiple informants (adolescents and parents), and measures that included multiple methods (self-reports, register based, and biological). The response rate of this study was also relatively high compared to previous questionnaire-based studies.

The hypotheses tested here were derived from previous research based primarily on bivariate associations between a limited number of variables. By replicating this type of bivariate statistical procedure, our data confirmed the majority of these hypotheses. However, when integrating the same variables in adolescent and caregiver path models, the nature, and significance of some of these associations changed or disappeared, rejecting some of the hypotheses.

Self-efficacy, emotional well-being of the adolescent, and difficulties in relation to adolescent/caregiver interactions (parental non-support and family conflict) proved to be particularly important aspects for understanding individual variability in adherence and metabolic control among adolescents with T1DM.

Family-related factors

Both adolescent and caregiver assessment of diabetes-related parental support were related to adherence, whereas only caregiver assessment was related directly to metabolic control. Moreover, caregiver appraisal of adolescent adherence behavior partially mediated the relationship between parental assessment of support and metabolic control. Previous studies have supported an association between parental assessment of family support and metabolic control [31]. Parents' perception of support could be speculated to be a reflection of a positive family milieu and a parenting style that increases the social and diabetes-related competencies of the adolescent, potentially decreasing the stress associated with living with T1DM in some adolescents. Other studies have suggested that worsening metabolic control, could lead parents to withdraw their support [32].

The adolescent model concurs with that of Lewin et al. [29], who found reports of parental support by children to be associated with adherence but not HbA_{1c}. However, the relationship between adolescent assessment of parental support and adherence in the current study was relatively weak. Previous studies suggest that, in adolescents, it may be beneficial to consider the influence of both caregivers' and friends' support in predicting better adjustment to living with a chronic illness [59, 60].

Adolescent assessment of diabetes-related parental non-support was related to metabolic control but not to adherence. The reverse was found for the caregiver assessments; non-support was related to adherence but not to metabolic control. Perhaps adolescents' perception of parental non-supportive behavior, such as critical and negative parenting, contributes to a stressful environment for the youth with diabetes which might affect their HbA_{1c} level, regardless of their adherence efforts. High levels of stress have previously been linked to poor metabolic control [61]. Ott et al. [25] found that parental non-support as rated by adolescents was related to blood glucose monitoring but not to other components of adherence. We did not assess the association between individual aspects of adherence and metabolic control, which may have altered the result regarding the lack of association between non-support and adherence. Moreover, it may be the lack of supportive behavior, such as positive encouragement and affirmation, more than adolescent perception of parental non-support that interacts with adolescents' adherence behavior. Criticizing and nagging behaviors by their caregivers may influence other aspects of the daily lives of adolescents with T1DM. Caregiver reports in the current study indicate that caregivers who perceive their actions towards their youngsters as more non-supportive, such as criticizing and nagging, also perceive their adolescent as less adherent to diabetes care. Perhaps, caregivers who feel prone to criticizing are more focused on any sign of what they perceive as non-adherent behavior in their child or are more reactive when it comes to non-adherence.

In the adolescent model, no direct association between diabetes-related family conflict and metabolic control was found, challenging a number of previous findings based on either adolescent report or merged adolescent-caregiver conflict scores [26, 28]. Instead, mediation analysis revealed that the relationship between HbA_{1c} and conflict was fully mediated by adherence, which is corroborated by previous studies applying similar statistical methods [29, 62]. A family climate characterized by high levels of conflict has been speculated to lead to adolescent stress, and that stress may affect metabolic control [28]. However, the adolescent-

reported level of conflict in the current sample was significantly lower ($M = 20.6$) than the normative sample for the Diabetes Family Conflict Scale ($M = 24.4$) [26], which may indicate an equally low level of stress and no subsequent effect on metabolic control.

In contrast, in the caregiver model, higher levels of family conflict were associated with poorer HbA_{1c} levels. This result is more in concordance with Drotar et al. [28] and Hood et al. [26], who also found an association between diabetes-related conflict in the family (based on either an adolescent-caregiver merged conflict score, or caregiver scores, respectively) and metabolic control. However, in the caregiver path model, the relationship between conflict and HbA_{1c} levels was partially mediated by adherence, indicating that conflict as assessed by caregivers is both directly and indirectly associated with metabolic control, possibly due to the physiological stress caused by the presence of family conflict.

The hypothesized relationship between the division of diabetes-related responsibilities and adherence was not confirmed in the adolescent path model. Perhaps the result would have been different had we used the Diabetes Family Responsibility Questionnaire to assess parent-child *agreement/disagreement* regarding the taking of responsibility. This factor has previously been linked to adherence [36]. Helgeson et al. [63] used the Diabetes Family Responsibility Questionnaire to assess *sharing of diabetes-related responsibility* and found that parent and child sharing of responsibility was associated with self-management, whereas either parent or child taking responsibility was not. Thus, they also highlighted a possible problem in scoring the Diabetes Family Responsibility Questionnaire as a continuous scale ranging from child to caregiver responsibility, as was done in this study.

The caregiver model showed an association between caregiver assessment of responsibility and adherence, indicating that adolescents who the caregiver saw as taking more responsibility for diabetes care tasks were also perceived as exhibiting better adherence toward treatment. This finding is somewhat contradictory with previous studies, which found that higher levels of adolescent responsibility are associated with a decrease in self-management behavior [64]. One possible explanation could be that caregivers who transfer the majority of treatment responsibilities to their adolescents are confident, or at least hopeful, regarding their child's ability to take on these tasks, which is then reflected in their assessment of adherence. Whether this is an accurate appraisal could be questioned, as it is not reflected in the adolescents' responses.

For both adolescents' and caregiver's reports, good general family functioning was related to better adherence. This finding is in line with previous research indicating family functioning to be associated with adherence based on both adolescent and caregiver reports [38, 65] and also our results regarding caregiver support, which could be considered an aspect of a positive family milieu.

Assessment of emotional health and diabetes-related self-efficacy

Better metabolic control was associated with increased anxiety, which contradicts the results of our bivariate analysis and the majority of previous studies [11]. However, some other studies have found that internalizing problems, or specifically anxiety, is related to improved metabolic control based on either adolescent or caregiver assessments [66, 67]. One possible explanation for this association could be that fear of hypoglycemia leads adolescents to strive for a higher blood sugar level, reducing the risk of hypoglycemia, and perhaps also their level of anxiety [68].

We could also speculate that a certain level of anxiety in adolescents with T1DM could be adaptive. We previously found that the overall level of anxiety symptoms in Danish children and adolescents with T1DM are comparable or lower than in a normative Danish sample, indicating that the mental health of this patient group is fairly good [13]; the current sample was

part of that study and we therefore assume that, the level of anxiety is generally not at a critical or impairing level. Symptoms of anxiety or an anxiety-prone personality in some individuals with T1DM may lead to more adherence toward treatment recommendations to prevent or reduce the anxiety-inducing consequences of non-adherence and poor metabolic control, leading to better metabolic control. The adolescent path model also showed that, though adherence moderately correlated with anxiety, it also acted as a partial mediator in the association between anxiety and HbA_{1c}.

Even though the bivariate correlation analysis showed a significant association between symptoms of depression and HbA_{1c}, in the path model the relationship between these two variables became non-significant. This contradicts a number of previous studies in which depression was shown to be associated with metabolic control [12]. A possible explanation for our finding could be that symptoms of depression do not always have a direct effect on metabolic control, but lead instead to a decrease in diabetes self-management tasks, as evidenced by our analysis showing adherence as a partial mediator of the relationship between HbA_{1c} and symptoms of depression, which is also supported by previous findings based on adolescent assessment [29]. Symptoms of depression may also have an even more indirect effect on adapting to living with diabetes, which was not tested in this study, through effects on the general family milieu and functioning, child/adolescent self-efficacy, or an increase in the diabetes-related conflict in the family. In their review of anxiety and depression in juvenile diabetes, Dantzer et al. [19] highlighted how several studies have suggested that adaptation plays a more important role in predicting metabolic control than symptoms of depression, but that depression may influence this process of adaptation. Again, it is important to note that the general psychological well-being of Danish adolescents with T1DM in the presents study appears to be less impacted compared to other samples which might also have affected the results [11, 13]. More studies are needed to understand the complex interaction of diabetes and depression, not disregarding the complex contributions of physiological factors [69].

The association between caregiver assessments of the social and emotional health (the Strength and Difficulties Questionnaire) of adolescents and metabolic control was fully mediated by adherence. This result appears to support the lack of direct correlation between symptoms of depression and metabolic control found in the adolescent path model.

Regarding adolescent assessment of self-efficacy in relation to diabetes care, no direct association with metabolic control was found, contradicting a number of previous studies [21, 54]. Instead, the association was fully mediated by adherence, supporting the work of Herge et al. [70]. Self-efficacy has previously been found to be related to active coping behaviors, and it could be that it is these behaviors, more than self-efficacy, that affect metabolic control [71]. The association between adherence and self-efficacy ($r = 0.44$) was the strongest association found in the adolescent path model and highlights the importance of this aspect of adolescents' adaption to living with T1DM.

Demographic and treatment-related variables

Both the adolescent and caregiver path models showed that better adherence is associated with lower HbA_{1c} levels, just as longer duration of T1DM and older age of the adolescent is related to both decreased adherence level and worse metabolic control. Both the adolescent and caregiver models showed an association between CSII treatment and better adherence. However, only the adolescent model found CSII treatment to be related to improved metabolic control.

In contrast to the adolescent model, the caregivers' report of the living situation of the adolescent was not associated with HbA_{1c}. In addition, the education level of the caregiver was unrelated to adherence and metabolic control in both models. The gender of the adolescent

was also not associated with metabolic control in either model, but weakly associated with adherence in the caregiver model, indicating that caregivers perceive boys with T1DM to be slightly more adherent than girls, replicating the conclusion of others [72].

Overall, the adolescent path model accounted for 25% of the variance in metabolic control and 51% of the variance in adherence, whereas the caregiver path model accounted for 26% of the variance in HbA_{1c} and 49% of the variance in adherence. Thus, our results regarding explained variance are comparable, or even an improvement on previous multivariate models [5].

Comparing the two models based on explained variance, no substantial differences were found. However, the caregiver model had a slightly better model fit. Based on findings regarding associations between included variables, this study highlights the additional information gained from consulting both adolescents and caregivers.

Limitations

Our results should be interpreted in light of several limitations. First, although several risk factors for poor adherence and metabolic control are identified in this study, the cross-sectional design prevents us from offering causal or bidirectional explanations for the relationships between the included variables. In particular, as demonstrated by Maxwell and Cole [73], cross-sectional analyses of mediation may be biased and hence causality cannot be determined. Second, the fact that non-participants had less optimal metabolic control than adolescents from participating families may also have influenced the outcome, though this bias is commonly found in this type of research. We can only speculate that the inclusion of these participants may have strengthened the associations between the investigated variables and outcomes. Third, with only 1.3% of the participants reporting the preferred language spoken in the home not being Danish, an insufficient number of minority families participated in the study to determine whether the results may generalize to this group. Furthermore, 83.6% of the responding caregivers were female, with 98.5% being the biological mother of the adolescent. Previous studies have shown that fathers have a somewhat differing view of the treatment efforts and emotional well-being of the child with diabetes [74]. Fourth, the relatively low internal consistency found for both the supportive and non-supportive subscales of the Diabetes Family Behavior Checklist may have had effect on the results involving these subscales. The reliability of this questionnaire has previously been found to be somewhat low [75], and future research could benefit from a more reliable measure of family support to confirm the results of the current study.

Lastly, dividing participants into smaller age segments may have revealed different patterns of interactions between psychosocial variables and outcomes that did not appear in our models, as others have found age to act as a moderator of the family functioning-adherence-metabolic control associations [76], and that developmental needs and reliance on caregivers vary greatly throughout the course of adolescence [77].

Although one of the major strengths of this study was the large, national, demographically diverse sample, the relatively low level of poverty in the Danish community, the financial support, and free access to a public health care system that covers many of the medically related expenses of children and adolescents with T1DM distinguishes this population, to some degree, from many other national samples.

Clinical implications

This study adds to the growing body of research highlighting the importance of addressing the psychological and psychosocial well-being of adolescents with T1DM.

As the relationship between adherence and self-efficacy was the strongest association in the adolescent model, this study emphasizes the importance of adolescent self-efficacy in relation to diabetes care. Finding a way to build or strengthen the adolescents' beliefs in their ability to overcome obstacles in relation to their daily self-management of diabetes care may increase the attention given to the adherence behavior of this group, improving metabolic control.

This study agrees with previous results stressing the importance of monitoring the emotional well-being of adolescents with T1DM, as the presence of emotional difficulties could affect the adolescents' adherence level. Interventions focusing on the importance of a supportive family milieu would probably prove beneficial, both with respect to adherence and metabolic control.

Diabetes duration may also be a focal point in the clinical care of children and adolescents, not only because of the physiological implications of a longer duration of T1DM, but also because adolescents with a longer duration have been shown to be in need of additional support to re-commit and improve diabetes care activities and goals [4].

Adolescent self-efficacy, finding a way to handle anxiety without damaging diabetes outcomes, and educating caregivers with regard to developmentally appropriate levels of support appear to be valuable focal points.

Supporting information

S1 Fig. Study inclusion.

(PDF)

Acknowledgments

The authors thank the Danish Society for Diabetes in Childhood and Adolescence for their cooperation, their contribution of data from the Danish Registry of Childhood Diabetes, and for encouraging the Society's members to participate. The authors extend their gratitude to all of the health care providers at the pediatric diabetes units throughout Denmark who supported this research and are most grateful to all of the families who took the time to participate in our survey.

Author Contributions

Conceptualization: Lene Juel Kristensen, Niels Holtum Birkebaek, Anne Hvarregaard Mose, Morten Berg Jensen, Mikael Thastum.

Data curation: Lene Juel Kristensen, Niels Holtum Birkebaek, Anne Hvarregaard Mose, Mikael Thastum.

Formal analysis: Lene Juel Kristensen, Morten Berg Jensen, Mikael Thastum.

Funding acquisition: Niels Holtum Birkebaek, Mikael Thastum.

Investigation: Lene Juel Kristensen, Niels Holtum Birkebaek, Anne Hvarregaard Mose, Mikael Thastum.

Methodology: Lene Juel Kristensen, Niels Holtum Birkebaek, Anne Hvarregaard Mose, Morten Berg Jensen, Mikael Thastum.

Project administration: Lene Juel Kristensen, Niels Holtum Birkebaek, Anne Hvarregaard Mose, Mikael Thastum.

Resources: Lene Juel Kristensen, Niels Holtum Birkebaek, Mikael Thastum.

Software: Lene Juel Kristensen, Morten Berg Jensen.

Supervision: Niels Holtum Birkebaek, Anne Hvarregaard Mose, Mikael Thastum.

Validation: Lene Juel Kristensen, Niels Holtum Birkebaek, Anne Hvarregaard Mose, Morten Berg Jensen, Mikael Thastum.

Writing – original draft: Lene Juel Kristensen.

Writing – review & editing: Lene Juel Kristensen, Niels Holtum Birkebaek, Anne Hvarregaard Mose, Morten Berg Jensen, Mikael Thastum.

References

1. Nathan DM, Genuth S, Lachin J, Cleary P, Crofford O, Davis M, et al. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med.* 1993; 329(14):977–86. <https://doi.org/10.1056/NEJM199309303291401> PMID: 8366922
2. Nathan DM. The Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications Study at 30 Years: Overview. *Diabetes Care.* 2014; 37(1):9–16. <https://doi.org/10.2337/dc13-2112> PMID: 24356592
3. Hood KK, Beavers DP, Yi-Frazier J, Bell R, Dabelea D, McKeown RE, et al. Psychosocial Burden and Glycemic Control During the First 6 Years of Diabetes: Results From the SEARCH for Diabetes in Youth Study. *Journal of Adolescent Health.* 2014; 55(4):498–504. <https://doi.org/10.1016/j.jadohealth.2014.03.011> PMID: 24815959
4. Chao A, Whittemore R, Minges KE, Murphy KM, Grey M. Self-Management in Early Adolescence and Differences by Age at Diagnosis and Duration of Type 1 Diabetes. *The Diabetes Educator.* 2014; 40(2):167–77. <https://doi.org/10.1177/0145721713520567> PMID: 24470042
5. Agarwal S, Jawad AF, Miller VA. A multivariate model exploring the predictive value of demographic, adolescent, and family factors on glycemic control in adolescents with type 1 diabetes. *Pediatric diabetes.* 2016; 17(7):500–8. <https://doi.org/10.1111/pedi.12331> PMID: 26486450
6. Rechenberg K, Whittemore R, Grey M, Jaser S, Group tTR. Contribution of income to self-management and health outcomes in pediatric type 1 diabetes. *Pediatric diabetes.* 2016; 17(2):120–6. <https://doi.org/10.1111/pedi.12240> PMID: 25545117
7. Swift EE, Chen R, Hershberger A, Holmes CS. Demographic risk factors, mediators, and moderators in youths' diabetes metabolic control. *Ann Behav Med.* 2006; 32(1):39–49. https://doi.org/10.1207/s15324796abm3201_5 PMID: 16827628
8. Samuelsson U, Anderzen J, Gudbjornsdottir S, Steineck I, Akesson K, Hanberger L. Teenage girls with type 1 diabetes have poorer metabolic control than boys and face more complications in early adulthood. *J Diabetes Complications.* 2016; 30(5):917–22. <https://doi.org/10.1016/j.jdiacomp.2016.02.007> PMID: 27052153
9. Nordly S, Jorgensen T, Andreassen AH, Hermann N, Mortensen HB. Quality of diabetes management in children and adolescents in Denmark. *Diabet Med.* 2003; 20(7):568–74. PMID: 12823239
10. Guo J, Whittemore R, Grey M, Wang J, Zhou ZG, He GP. Diabetes self-management, depressive symptoms, quality of life and metabolic control in youth with type 1 diabetes in China. *J Clin Nurs.* 2013; 22(1–2):69–79. <https://doi.org/10.1111/j.1365-2702.2012.04299.x> PMID: 23106340
11. Buchberger B, Huppertz H, Krabbe L, Lux B, Mattivi JT, Siafarikas A. Symptoms of depression and anxiety in youth with type 1 diabetes: A systematic review and meta-analysis. *Psychoneuroendocrinology.* 2016; 70:70–84. <https://doi.org/10.1016/j.psyneuen.2016.04.019> PMID: 27179232
12. Reynolds KA, Helgeson VS. Children with diabetes compared to peers: depressed? Distressed? A meta-analytic review. *Ann Behav Med.* 2011; 42(1):29–41. <https://doi.org/10.1007/s12160-011-9262-4> PMID: 21445720
13. Kristensen LJ, Birkebaek NH, Mose AH, Hohwu L, Thastum M. Symptoms of emotional, behavioral, and social difficulties in the danish population of children and adolescents with type 1 diabetes—results of a national survey. *PloS one.* 2014; 9(5):e97543. <https://doi.org/10.1371/journal.pone.0097543> PMID: 24842772
14. Sivertsen B, Petrie KJ, Wilhelmsen-Langeland A, Hysing M. Mental health in adolescents with Type 1 diabetes: results from a large population-based study. *BMC endocrine disorders.* 2014; 14:83. Epub 2014/10/12. <https://doi.org/10.1186/1472-6823-14-83> PMID: 25303963

15. Helgeson VS, Siminerio L, Escobar O, Becker D. Predictors of Metabolic Control among Adolescents with Diabetes: A 4-Year Longitudinal Study. *Journal of Pediatric Psychology*. 2009; 34(3):254–70. <https://doi.org/10.1093/jpepsy/jsn079> PMID: 18667479
16. Herzer M, Hood KK. Anxiety Symptoms in Adolescents with Type 1 Diabetes: Association with Blood Glucose Monitoring and Glycemic Control. *Journal of Pediatric Psychology*. 2010; 35(4):415–25. <https://doi.org/10.1093/jpepsy/jsp063> PMID: 19684117
17. Gonzalez JS, Peyrot M, McCarl LA, Collins EM, Serpa L, Mimiaga MJ, et al. Depression and Diabetes Treatment Nonadherence: A Meta-Analysis. *Diabetes Care*. 2008; 31(12):2398–403. <https://doi.org/10.2337/dc08-1341> PMID: 19033420
18. Armstrong B, Mackey ER, Streisand R. Parenting behavior, child functioning, and health behaviors in preadolescents with type 1 diabetes. *J Pediatr Psychol*. 2011; 36(9):1052–61. <https://doi.org/10.1093/jpepsy/jsr039> PMID: 21828111
19. Dantzer C, Swendsen J, Maurice-Tison S, Salamon R. Anxiety and depression in juvenile diabetes: a critical review. *Clin Psychol Rev*. 2003; 23(6):787–800. PMID: 14529698
20. Bandura A. *Self-efficacy: The exercise of control*. New York: W. H. Freeman; 1997 1997.
21. Chih AH, Jan CF, Shu SG, Lue BH. Self-efficacy Affects Blood Sugar Control Among Adolescents With Type I Diabetes Mellitus. *Journal of the Formosan Medical Association*. 2010; 109(7):503–10. [https://doi.org/10.1016/S0929-6646\(10\)60084-8](https://doi.org/10.1016/S0929-6646(10)60084-8) PMID: 20654789
22. Rasbach LE, Volkening LK, Markowitz JT, Butler DA, Katz ML, Laffel LM. Youth and parent measures of self-efficacy for continuous glucose monitoring: survey psychometric properties. *Diabetes Technol Ther*. 2015; 17(5):327–34. <https://doi.org/10.1089/dia.2014.0366> PMID: 25695341
23. Stupiansky NW, Hanna KM, Slaven JE, Weaver MT, Fortenberry JD. Impulse Control, Diabetes-Specific Self-Efficacy, and Diabetes Management Among Emerging Adults With Type 1 Diabetes. *Journal of Pediatric Psychology*. 2013; 38(3):247–54. <https://doi.org/10.1093/jpepsy/jss110> PMID: 23115219
24. Sander EP, Odell S, Hood KK, Ph.D. Diabetes-Specific Family Conflict and Blood Glucose Monitoring in Adolescents With Type 1 Diabetes: Mediation Role of Diabetes Self-Efficacy. *Diabetes Spectrum*. 2010; 23(2):89–94.
25. Ott J, Greening L, Palardy N, Holderby A, Debell WK. Self-efficacy as a mediator variable for adolescents' adherence to treatment for insulin-dependent diabetes mellitus. *Children's Health Care*. 2000; 29(1):47–63.
26. Hood KK, Butler DA, Anderson BJ, Laffel LMB. Updated and Revised Diabetes Family Conflict Scale. *Diabetes Care*. 2007; 30(7):1764–9. <https://doi.org/10.2337/dc06-2358> PMID: 17372149
27. Ingerski LM, Anderson BJ, Dolan LM, Hood KK. Blood Glucose Monitoring and Glycemic Control in Adolescence: Contribution of Diabetes-Specific Responsibility and Family Conflict. *Journal of Adolescent Health*. 2010; 47(2):191–7. <https://doi.org/10.1016/j.jadohealth.2010.01.012> PMID: 20638012
28. Drotar D, Ittenbach R, Rohan JM, Gupta R, Pendley JS, Delamater A. Diabetes management and glycemic control in youth with type 1 diabetes: test of a predictive model. *J Behav Med*. 2013; 36(3):234–45. <https://doi.org/10.1007/s10865-012-9426-0> PMID: 22569775
29. Whittemore R, Liberti L, Jeon S, Chao A, Jaser SS, Grey M. Self-management as a mediator of family functioning and depressive symptoms with health outcomes in youth with type 1 diabetes. *West J Nurs Res*. 2014; 36(9):1254–71. <https://doi.org/10.1177/0193945913516546> PMID: 24357648
30. Williams LB, Laffel LM, Hood KK. Diabetes-specific family conflict and psychological distress in paediatric Type 1 diabetes. *Diabet Med*. 2009; 26(9):908–14. <https://doi.org/10.1111/j.1464-5491.2009.02794.x> PMID: 19719712
31. Lewin AB, Geffken GR, Heidgerken AD, Duke DC, Novoa W, Williams LB, et al. The Diabetes Family Behavior Checklist: A Psychometric Evaluation. *Journal of clinical psychology in medical settings*. 2005; 12(4):315–22.
32. Seiffge-Krenke I, Laursen B, Dickson DJ, Hartl AC. Declining Metabolic Control and Decreasing Parental Support Among Families With Adolescents With Diabetes: The Risk of Restrictiveness. *Journal of Pediatric Psychology*. 2013; 38(5):518–30. <https://doi.org/10.1093/jpepsy/jst006> PMID: 23564837
33. Duke DC, Geffken GR, Lewin AB, Williams LB, Storch EA, Silverstein JH. Glycemic Control in Youth with Type 1 Diabetes: Family Predictors and Mediators. *Journal of Pediatric Psychology*. 2008; 33(7):719–27. <https://doi.org/10.1093/jpepsy/jsn012> PMID: 18296726
34. Holmes CS, Chen R, Streisand R, Marschall DE, Souter S, Swift EE, et al. Predictors of Youth Diabetes Care Behaviors and Metabolic Control: A Structural Equation Modeling Approach. *Journal of Pediatric Psychology*. 2006; 31(8):770–84. <https://doi.org/10.1093/jpepsy/jsj083> PMID: 16221954
35. Marker AM, Noser AE, Clements MA, Patton SR. Shared Responsibility for Type 1 Diabetes Care Is Associated With Glycemic Variability and Risk of Glycemic Excursions in Youth. *J Pediatr Psychol*. 2018; 43(1):61–71. <https://doi.org/10.1093/jpepsy/jsx081> PMID: 28541572

36. Vesco AT, Anderson BJ, Laffel LMB, Dolan LM, Ingerski LM, Hood KK. Responsibility Sharing between Adolescents with Type 1 Diabetes and Their Caregivers: Importance of Adolescent Perceptions on Diabetes Management and Control. *Journal of Pediatric Psychology*. 2010; 35(10):1168–77. <https://doi.org/10.1093/jpepsy/jsq038> PMID: 20444852
37. Leonard BJ, Jang YP, Savik K, Plumbo MA. Adolescents With Type 1 Diabetes: Family Functioning and Metabolic Control. *Journal of Family Nursing*. 2005; 11(2):102–21. <https://doi.org/10.1177/1074840705275152> PMID: 16287821
38. Moore SM, Hackworth NJ, Hamilton VE, Northam EP, Cameron FJ. Adolescents with type 1 diabetes: parental perceptions of child health and family functioning and their relationship to adolescent metabolic control. *Health Qual Life Outcomes*. 2013; 11:50. <https://doi.org/10.1186/1477-7525-11-50> PMID: 23521786
39. Gowers SG, Jones JC, Kiana S, North CD, Price DA. Family Functioning: A Correlate of Diabetic Control? *Journal of Child Psychology and Psychiatry*. 1995; 36(6):993–1001. PMID: 7593406
40. Whittmore R, Jaser S, Guo J, Grey M. A conceptual model of childhood adaptation to type 1 diabetes. *Nurs Outlook*. 2010; 58(5):242–51. <https://doi.org/10.1016/j.outlook.2010.05.001> PMID: 20934079
41. Kristensen LJ, Thastum M, Mose AH, Birkebaek NH, Adolescence TDSfDiCa. Psychometric Evaluation of the Adherence in Diabetes Questionnaire. *Diabetes Care*. 2012; 35(11):2161–6. <https://doi.org/10.2337/dc11-2342> PMID: 22837365
42. Schafer LC, Glasgow RE, McCaul KD, Dreher M. Adherence to IDDM regimens: relationship to psychosocial variables and metabolic control. *Diabetes Care*. 1983; 6(5):493–8. PMID: 6336344
43. Schafer LC, Glasgow RE, McCaul KD. Increasing the adherence of diabetic adolescents. *J Behav Med*. 1982; 5(3):353–62. PMID: 7131549
44. Schafer LC, McCaul KD, Glasgow RE. Supportive and nonsupportive family behaviors: relationships to adherence and metabolic control in persons with type I diabetes. *Diabetes Care*. 1986; 9(2):179–85. PMID: 3698784
45. Anderson BJ, Auslander WF, Jung KC, Miller JP, Santiago JV. Assessing Family Sharing of Diabetes Responsibilities. *Journal of Pediatric Psychology*. 1990; 15(4):477–92. PMID: 2258796
46. Epstein NB, Baldwin LM, Bishop DS. The McMaster Family Assessment Device. *Journal of Marital and Family Therapy*. 1983; 9(2):171–80.
47. Miller IW, Epstein NB, Bishop DS, Keitner GI. The McMaster Family Assessment Device: Reliability and Validity. *Journal of Marital and Family Therapy*. 1985; 11(4):345–56.
48. Alderfer MA, Fiese BH, Gold JI, Cutuli JJ, Holmbeck GN, Goldbeck L, et al. Evidence-based Assessment in Pediatric Psychology: Family Measures. *Journal of Pediatric Psychology*. 2008; 33(9):1046–61. <https://doi.org/10.1093/jpepsy/jsm083> PMID: 17905801
49. Beck JS, Beck AT, Jolly JB, Steer RA. *Beck Youth Inventories™ for children and adolescents—Second Edition, Vejledning—Dansk version*: NCS Pearson, Inc.; 2012 2012.
50. Thastum M, Ravn K, Sommer S, Trillingsgaard A. Reliability, validity and normative data for the Danish Beck Youth Inventories. *Scandinavian Journal of Psychology*. 2009; 50:47–54. <https://doi.org/10.1111/j.1467-9450.2008.00690.x> PMID: 18980602
51. Goodman R. The Strengths and Difficulties Questionnaire: a research note. *J Child Psychol Psychiatry*. 1997; 38(5):581–6. PMID: 9255702
52. Niclasen J, Teasdale TW, Andersen AM, Skovgaard AM, Elberling H, Obel C. Psychometric Properties of the Danish Strength and Difficulties Questionnaire: The SDQ Assessed for More than 70,000 Raters in Four Different Cohorts. *PLoS One*. 2012; 7(2):e32025. <https://doi.org/10.1371/journal.pone.0032025> PMID: 22384129
53. Goodman R. Psychometric properties of the strengths and difficulties questionnaire. *J Am Acad Child Adolesc Psychiatry*. 2001; 40(11):1337–45. <https://doi.org/10.1097/00004583-200111000-00015> PMID: 11699809
54. Iannotti RJ, Schneider S, Nansel TR, Haynie DL. Self-efficacy, outcome expectations, and diabetes self-management in adolescents with type 1 diabetes. *Journal of Developmental and Behavioral Pediatrics*. 2006; 27(2):98–105. PMID: 16682872
55. Cohen J. *Statistical power analysis for the behavioral sciences*. 2nd ed. ed. New Jersey: Lawrence Erlbaum; 1988.
56. Muthén LKM B.O. *Mplus User's Guide*. Seventh Edition. Los Angeles, CA: Muthén & Muthén; 1998–2015.
57. Shrout PE, Bolger N. Mediation in experimental and nonexperimental studies: New procedures and recommendations. *Psychological Methods*. 2002; 7(4):422–45. PMID: 12530702
58. Li Hu, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling: A Multidisciplinary Journal*. 1999; 6(1):1–55.

59. Drew LM, Berg C, Wiebe DJ. The mediating role of extreme peer orientation in the relationships between adolescent-parent relationship and diabetes management. *J Fam Psychol*. 2010; 24(3):299–306. <https://doi.org/10.1037/a0019352> PMID: 20545403
60. Oris L, Seiffge-Krenke I, Moons P, Goubert L, Rassart J, Goossens E, et al. Parental and peer support in adolescents with a chronic condition: a typological approach and developmental implications. *Journal of behavioral medicine*. 2016; 39(1):107–19. Epub 2015/09/16. <https://doi.org/10.1007/s10865-015-9680-z> PMID: 26369633
61. Hilliard ME, Yi-Frazier JP, Hessler D, Butler AM, Anderson BJ, Jaser S. Stress and A1c Among People with Diabetes Across the Lifespan. *Current diabetes reports*. 2016; 16(8):67–. <https://doi.org/10.1007/s11892-016-0761-3> PMID: 27287017
62. Hilliard ME, Guilfoyle SM, Dolan LM, Hood KK. Prediction of adolescents' glycemic control 1 year after diabetes-specific family conflict: the mediating role of blood glucose monitoring adherence. *Arch Pediatr Adolesc Med*. 2011; 165(7):624–9. <https://doi.org/10.1001/archpediatrics.2011.86> PMID: 21727273
63. Helgeson VS, Reynolds KA, Siminerio L, Escobar O, Becker D. Parent and Adolescent Distribution of Responsibility for Diabetes Self-care: Links to Health Outcomes. *Journal of Pediatric Psychology*. 2008; 33(5):497–508. <https://doi.org/10.1093/jpepsy/jsm081> PMID: 17848390
64. Hsin O, La Greca AM, Valenzuela J, Moine CT, Delamater A. Adherence and Glycemic Control among Hispanic Youth with Type 1 Diabetes: Role of Family Involvement and Acculturation. *Journal of Pediatric Psychology*. 2010; 35(2):156–66. <https://doi.org/10.1093/jpepsy/jsp045> PMID: 19491214
65. Hauser ST, Jacobson AM, Lavori P, Wolfsdorf JI, Herskowitz RD, Milley JE, et al. Adherence Among Children and Adolescents With Insulin-Dependent Diabetes Mellitus Over a Four-Year Longitudinal Follow-Up: II. Immediate and Long-Term Linkages With the Family Milieu. *Journal of Pediatric Psychology*. 1990; 15(4):527–42. PMID: 2258799
66. Dusan V, Jovan V, Nada K, Dragan K, Georgios K, Biroo M. Psychological aspects of adolescents with diabetes mellitus. *Procedia—Social and Behavioral Sciences*. 2010; 5:1788–93.
67. Cohen DM, Lumley MA, Naar-King S, Partridge T, Cakan N. Child Behavior Problems and Family Functioning as Predictors of Adherence and Glycemic Control in Economically Disadvantaged Children with Type 1 Diabetes: A Prospective Study. *Journal of Pediatric Psychology*. 2004; 29(3):171–84. PMID: 15131135
68. Driscoll KA, Raymond J, Naranjo D, Patton SR. Fear of Hypoglycemia in Children and Adolescents and Their Parents with Type 1 Diabetes. *Current diabetes reports*. 2016; 16(8):77. <https://doi.org/10.1007/s11892-016-0762-2> PMID: 27370530
69. Lang UE, Borgwardt S. Molecular Mechanisms of Depression: Perspectives on New Treatment Strategies. *Cellular Physiology and Biochemistry*. 2013; 31(6):761–77. <https://doi.org/10.1159/000350094> PMID: 23735822
70. Herge WM, Streisand R, Chen R, Holmes C, Kumar A, Mackey ER. Family and Youth Factors Associated With Health Beliefs and Health Outcomes in Youth With Type 1 Diabetes. *Journal of Pediatric Psychology*. 2012; 37(9):980–9. <https://doi.org/10.1093/jpepsy/jss067> PMID: 22661616
71. Rose M, Fliege H, Hildebrandt M, Schirop T, Klapp BF. The network of psychological variables in patients with diabetes and their importance for quality of life and metabolic control. *Diabetes Care*. 2002; 25(1):35–42. PMID: 11772898
72. Stewart SM, Lee PWH, Waller D, Hughes CW, Low LCK, Kennard BD, et al. A Follow-Up Study of Adherence and Glycemic Control Among Hong Kong Youths With Diabetes. *Journal of Pediatric Psychology*. 2003; 28(1):67–79. PMID: 12490633
73. Maxwell SE, Cole DA. Bias in cross-sectional analyses of longitudinal mediation. *Psychol Methods*. 2007; 12(1):23–44. <https://doi.org/10.1037/1082-989X.12.1.23> PMID: 17402810
74. Berg CA, Butner JE, Turner SL, Lansing AH, King P, Wiebe DJ. Adolescents', mothers', and fathers' reports of adherence across adolescence and their relation to HbA1c and daily blood glucose. *Journal of behavioral medicine*. 2016; 39(6):1009–19. <https://doi.org/10.1007/s10865-016-9771-5> PMID: 27501733
75. Hanna KM. Existing Measures of Diabetes-Specific Support for Use With Adolescents With Diabetes. *The Diabetes Educator*. 2006; 32(5):741–50. <https://doi.org/10.1177/0145721706291759> PMID: 16971707
76. Lewin AB, Heidgerken AD, Geffken GR, Williams LB, Storch EA, Gelfand KM, et al. The Relation Between Family Factors and Metabolic Control: The Role of Diabetes Adherence. *Journal of Pediatric Psychology*. 2006; 31(2):174–83. <https://doi.org/10.1093/jpepsy/jsj004> PMID: 16467317
77. Lee SL, Lo FS, Lee YJ, Chen BH, Wang RH. Predictors of Glycemic Control in Adolescents of Various Age Groups With Type 1 Diabetes. *J Nurs Res*. 2015; 23(4):271–9. <https://doi.org/10.1097/JNR.000000000000096> PMID: 26562458