

RESEARCH ARTICLE

# Intragenus (*Homo*) variation in a chemokine receptor gene (*CCR5*)

Kara C. Hoover <sup>1,2\*</sup>

**1** Department of Anthropology, University of Alaska Fairbanks, Fairbanks, Alaska, United States of America, **2** Biochemistry and Molecular Biology, University of Alaska Fairbanks, Fairbanks, Alaska, United States of America

\* [kchoover@alaska.edu](mailto:kchoover@alaska.edu)

## Abstract

Humans have a comparatively higher rate of more polymorphisms in regulatory regions of the primate *CCR5* gene, an immune system gene with both general and specific functions. This has been interpreted as allowing flexibility and diversity of gene expression in response to varying disease loads. A broad expression repertoire is useful to humans—the only globally distributed primate—due to our unique adaptive pattern that increased pathogen exposure and disease loads (e.g., sedentism, subsistence practices). The main objective of the study was to determine if the previously observed human pattern of increased variation extended to other members of our genus, *Homo*. The data for this study are mined from the published genomes of extinct hominins (four Neandertals and two Denisovans), an ancient human (Ust'-Ishim), and modern humans (1000 Genomes). An average of 15 polymorphisms per individual were found in human populations (with a total of 262 polymorphisms). There were 94 polymorphisms identified across extinct *Homo* (an average of 13 per individual) with 41 previously observed in modern humans and 53 novel polymorphisms (32 in Denisova and 21 in Neandertal). Neither the frequency nor distribution of polymorphisms across gene regions exhibit significant differences within the genus *Homo*. Thus, humans are not unique with regards to the increased frequency of regulatory polymorphisms and the evolution of variation patterns across *CCR5* gene appears to have originated within the genus. A broader evolutionary perspective on regulatory flexibility may be that it provided an advantage during the transition to confrontational foraging (and later hunting) that altered human-environment interaction as well as during migration to Eurasia and encounters with novel pathogens.

## Introduction

Chemokine receptors facilitate communication between cells and the environment [1, 2] and mediate the activity of chemokines, proteins secreted by the immune system genes to chemically recruit immune cells to infection sites via chemotaxis [2, 3]. The cell surface chemokine receptor *CCR5* (a G protein-coupled receptor) is best known for its adaptive immune system role in binding the M-tropic human immunodeficiency virus (HIV) and creating a gateway to host cell infection [3–12]. In several mammals, *CCR5* genes present high levels of gene



## OPEN ACCESS

**Citation:** Hoover KC (2018) Intragenus (*Homo*) variation in a chemokine receptor gene (*CCR5*). PLoS ONE 13(10): e0204989. <https://doi.org/10.1371/journal.pone.0204989>

**Editor:** Francesc Calafell, Universitat Pompeu Fabra, SPAIN

**Received:** September 27, 2017

**Accepted:** September 18, 2018

**Published:** October 2, 2018

**Copyright:** © 2018 Kara C. Hoover. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** The author received no specific funding for this work.

**Competing interests:** The authors have declared that no competing interests exist.

conversion with the chromosomally adjacent *CCR2* [13–18]. Primate *CCR5* gene structure, open reading frame (ORF), and amino acid identity are evolutionary highly conserved [2, 19–21] and interspecific gene sequences are functionally similar [20]. There is, however, common and significant variation across species outside conserved regions. Most of these polymorphisms are not deleterious to health and tolerated due to the redundancy of the chemokine family in ligand binding [2, 22]. New World Monkeys have a high number of functional polymorphisms due to lentivirus resistance [23]. Further, humans have been found to have a significantly high number of *cis*-regulatory region polymorphisms in comparison to 36 non-human primate species of apes, Old World Monkeys, and New World Monkeys [20]. Humans also have a specific a 32bp deletion in Exon 3, *CCR5Δ32* [24, 25], that results in a non-functional protein [24–29]) associated with HIV-resistance and West Nile Virus susceptibility in northern European populations [30–42].

Located on Chromosome 3 (3p21), human *CCR5* is 6,065 bases long with an ORF of 1,056 bases that codes for a protein with 352 residues. Two common transcripts (B with three exons and the more stable A with four) likely resulted from non-coding upstream polymorphisms in two separate gene promoters (the functionally weaker *cis*-acting promoter (P<sub>U</sub>) upstream of Exon 1 and the downstream promoter (P<sub>D</sub>) upstream of Exon 3 [19, 20]. These transcripts cause alternate splicing (differential inclusion or exclusion of exons) in messenger RNA that affects regulation of cell surface receptor expression levels [20–22].

The plasticity in regulation of gene expression via alternate transcripts and increased polymorphisms [20–22] makes human *CCR5* particularly interesting from a broader evolutionary perspective. *Homo* has one of the broadest adaptive ranges of any species [43] and human *CCR5*'s ability to rapidly respond to new pathogens [19, 20] may have served an adaptive function during evolutionary migration and shifts in human-environment relationships with changes to subsistence. Our genomes carry vast evidence of past disease responses [44–46] that are shared across the genus and reflect a unique disease pattern for *Homo*. For example, there is strong evidence for increased disease risk via genetic load in extinct *Homo* and past human populations [47] and archaeological evidence for past disease treatment (ingestion of anti-biotic and anti-fungal non-food plants) in Neandertals [48–52]. *CCR5* has been well studied due to its role in HIV infection (with a focus on natural selection acting on the 32bp deletion) but no work has explored variation within the genus *Homo* more broadly.

The plethora of research on the evolution *CCR5* was conducted prior to the generation of deep coverage, high quality paleogenomes for extinct hominins, such as Neandertal species and the newer Denisova species. While paleogenomic sample sizes are not robust to make statements on selection or add to a discussion of other evolutionary forces acting on variation, they provide an evolutionary dimension to understanding the patterns of variation characterizing our genus and insights into possible adaptations to new environments, subsistence regimes, and pathogens [53]. Plus, the sample of ancient genome is increasing every year. Just a few insights gained from a single paleogenomes include ground-breaking studies on evolution of skin color in humans [54] and Neandertals [55] and the introgression of functionally adaptive polymorphisms into the human immune system genes from Altai Neandertal [56]. Understanding the differences between derived and specific variation also enables potential differentiation of challenges we overcame as a genus such as obligate bipedalism [57] or high-altitude adaptation [58] versus challenges we overcame as a species such as the biocultural evolution of sickle-cell trait and malaria infection [59]. Thus, the overall aim of this research is to place humans within the context *Homo* and examine if the pattern of humans having a significantly higher number of *cis*-regulatory region polymorphisms (compared to 36 non-human primate species of apes, Old World Monkeys, and New World Monkeys) [20] is specific or one that is shared by our genus.

Variation in *CCR5* was examined in humans and extinct hominins to address the questions: are there shared patterns of variation across the genus for polymorphism frequency and is the distribution of polymorphisms across the gene suggestive of a common evolutionary trajectory? Based on previous studies on human-nonhuman primate gene structure and variation and the finding that human polymorphisms allow flexible *CCR5* gene expression [19, 20], the expectation is that there is a shared pattern of variation that aided adaptation for members of the geographically and ecologically dispersed genus *Homo*. Both expectations were met.

## Materials

Modern human data is from the 1000 Genomes Project [60], which contains data for 2,504 individuals from 26 populations (Table 1). While coverage is low per individual, the data are robust enough to identify the majority of polymorphisms at a frequency of at least 1% in the populations studied, which is suitable for the current study. Extinct *Homo* data are: Denisova 3 [61], Denisova 2 [62], Vindija 33.19 Neandertal [63], Altai Neandertal [61], El Sidron Neandertal [64], Mezmaiskaya 1 Neandertal [63], and an ancient human that contributed no genes to modern populations, Ust'-Ishim [65]—see Table 1 for accession numbers. These species are Pleistocene Eurasian hominins with Denisova representing an eastern Eurasian Pleistocene population and Neandertal a western one (with some overlap with Denisova in Siberia). All genomes have high coverage (excepting Mezmaiskaya and Sidron); contamination with modern human DNA is estimated to be less than 1% for the extinct hominins [61–63, 66, 67].

## Methods

The human reference sequences for two common transcript variants for the *CCR5* gene (NM\_000579 and NM\_001100168) were downloaded from the National Center for Biotechnology Information (NCBI). The modern human variation data (*CCR5* and *cis*-acting elements) were downloaded from 1000 Genomes via ftp as variant call format (VCF) files (<ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/>). All data were downloaded to and analyzed using the University of Alaska Research Computing Systems. All files were aligned to the human genome GRCh37/hg19. VCF files for six extinct hominin species and one extinct human (Ust'-Ishim) were downloaded from the Max Planck Institute Leipzig. Ancient DNA often contains C-to-T deaminations at the end of reads [68]. The lack of variation identified from paleogenomic sequence reads is unlikely to be a result of typical problems associated with ancient DNA sequence reads since chemical processes like deamination would increase SNPs (whether false or not). More significantly, the paleogenomes were generated using protocols that largely eliminates this error [69]. Despite high levels of variation at this locus and evidence for balancing selection in humans at this locus, strong levels of introgression from inter-breeding with Neandertals in Eurasia have not been reported at this locus, as they have for other immune system loci in similar scenarios [27, 70–75]; introgression data from the Reich lab (<https://reich.hms.harvard.edu/datasets>) [76] confirm this is the case (S1 Table). Moreover, the African genetic variation is similar to European genetic variation which suggests that diversity was already present in modern humans prior to any admixture with archaic species in Europe.

Distribution of polymorphisms was guided by the structure provided by Mummidi et al. [19] and included promoter regions ( $P_U$  and  $P_D$ ), ORF, and *CCR5*. The target area for  $P_U$  was the most inclusive range (-1976 to +33) which avoided overlap with  $P_D$  and because little difference was noted between putative  $P_U$  regions studied by Mummidi et al [19]. The target area for  $P_D$  was the most productive range (+119 to +828). Significant difference in distribution of polymorphisms across gene structure for all samples was tested using Monte Carlo methods for the exact test.

**Table 1. Samples plus polymorphism totals.**

| Region   | Population Name                                     | Pop  | n   | Var  | AvVar/Pop |
|--|---|------|-----|------|-----------|
| Africa   | African Caribbeans in Barbados                      | ACB  | 96  | 1464 | 15        |
|  | Americans of African Ancestry in SW USA             | ASW  | 61  | 1049 | 17        |
|  | Esan in Nigeria                                     | ESN  | 99  | 1499 | 15        |
|  | Gambian in Western Divisions in the Gambia          | GWD  | 113 | 1744 | 15        |
|  | Luhya in Webuye, Kenya                              | LWK  | 99  | 1469 | 15        |
|  | Mende in Sierra Leone                               | MSL  | 85  | 1420 | 17        |
|  | Yoruba in Ibadan, Nigeria                           | YRI  | 108 | 1786 | 17        |
| America  | Colombians from Medellin, Colombia                  | CLM  | 94  | 1688 | 18        |
|  | Mexican Ancestry from Los Angeles USA               | MXL  | 64  | 1111 | 17        |
|  | Peruvians from Lima, Peru                           | PEL  | 85  | 1184 | 14        |
|  | Puerto Ricans from Puerto Rico                      | PUR  | 104 | 1699 | 16        |
| East Asia  | Chinese Dai in Xishuangbanna, China                 | CDX  | 93  | 2222 | 24        |
|  | Han Chinese in Beijing, China                       | CHB  | 103 | 2278 | 22        |
|  | Southern Han Chinese                                | CHS  | 105 | 2328 | 22        |
|  | Japanese in Tokyo, Japan                            | JPT  | 104 | 1929 | 19        |
|  | Kinh in Ho Chi Minh City, Vietnam                   | KHV  | 99  | 2411 | 24        |
| Europe   | Utah Residents (CEPH, North/West European ancestry) | CEU  | 99  | 1664 | 17        |
|  | Finnish in Finland                                  | FIN  | 99  | 1484 | 15        |
|  | British in England and Scotland                     | GBR  | 91  | 1417 | 16        |
|  | Iberian Population in Spain                         | IBS  | 107 | 1755 | 16        |
|  | Toscans in Italia                                   | TSI  | 107 | 2132 | 20        |
| South Asia   | Bengali from Bangladesh                             | BEB  | 86  | 1729 | 20        |
|  | Gujarati Indian from Houston, Texas                 | GIH  | 103 | 1818 | 18        |
|  | Indian Telugu from the UK                           | ITU  | 102 | 1907 | 19        |
|  | Punjabi from Lahore, Pakistan                       | PJL  | 96  | 1831 | 19        |
|  | Sri Lankan Tamil from the UK                        | STU  | 102 | 1952 | 19        |
| Extinct Homo                                       | Altai Neandertal (ERP002097) <sup>1</sup>           | Alt  | 1   | 7    | 7         |
|  | Denisova 3 (ERP001519) <sup>2</sup>                 | Den1 | 1   | 10   | 10        |
|  | Denisova 2 (PRJEB20653) <sup>3</sup>                | Den2 | 1   | 39   | 39        |
|  | El Sidron (ERP002457) <sup>4</sup>                  | Sid  | 1   | 0    | 0         |
|  | Mezmaiskaya 1 Neandertal (PRJEB21195) <sup>5</sup>  | Mez  | 1   | 28   | 28        |
|  | Ust'-Ishim (PRJEB6622) <sup>6</sup>                 | Ust  | 1   | 15   | 15        |
| Vindija 33.19 Neandertal (PRJEB21157) <sup>7</sup> | Vin   | 1    | 8   | 8    |           |

<sup>1</sup> <http://cdna.eva.mpg.de/neandertal/altai/AltaiNeandertal/VCF/>

<sup>2</sup> <http://cdna.eva.mpg.de/neandertal/altai/Denisovan/>

<sup>3</sup> <http://cdna.eva.mpg.de/neandertal/Vindija/VCF/Denisova/>

<sup>4</sup> <http://cdna.eva.mpg.de/neandertal/exomes/VCF/Sidron/>

<sup>5</sup> <http://cdna.eva.mpg.de/neandertal/Vindija/VCF/Mez1/>

<sup>6</sup> <http://cdna.eva.mpg.de/ust-ishim/VCF/>

<sup>7</sup> <http://cdna.eva.mpg.de/neandertal/exomes/VCF/Vindija/>

<https://doi.org/10.1371/journal.pone.0204989.t001>

## Results and discussion

Previous research has examined gene structure [19], gene variation within primate species [20], and selection acting on the gene primarily in response to viral load [6, 10, 21, 22, 33, 36, 37, 40, 41, 77, 78]. The goal of this research was to establish if the pattern of human variation and distribution of polymorphisms in *CCR5* [20] is specific (i.e., unique in the human species) or genus-wide (i.e., a pattern shared by *Homo*).

Are there shared patterns of variation across the genus for polymorphism frequency? In the modern human sample, 262 known SNPs were observed (Tables 2 and S2 contains all 1000 Genomes variants). SNP frequency per individual (total SNPs in a population/total number of individuals) within the 26 populations ranged from 14 to 24, with East Asians exhibiting the highest variation and Africa and the Americas the least (Table 1). There were 94 polymorphisms identified across all extinct *Homo* samples (Altai, Denisova 3 and 2, Mezmaiskaya, Ust'-Ishim, Vindija, and El Sid), an average of 13 per individual included in analysis (Tables 1 and 2 and S1). No polymorphisms were found in the El Sidron specimen and, as a result, it is not included in the tables. Some polymorphisms in extinct *Homo* ( $n = 41$ ) have been previously observed in modern humans (Table 2). There were 53 novel polymorphisms identified, 32 in Denisova (1 in Denisova 3, 31 in Denisova 2) and 21 in Mezmaiskaya. Table 2 summarizes extinct *Homo* polymorphisms.

Is the distribution of polymorphisms across the gene suggestive of a common evolutionary trajectory? The frequency of polymorphisms across gene structure are used rather than counts because the human sample is much larger and captures an exponentially greater number of polymorphisms as a result (see S3 Table for raw count summary). Both humans and extinct members of our genus exhibit more polymorphisms in gene regulatory regions (Table 3) suggesting a shared pattern of variation across *Homo*. When polymorphisms occur in both  $P_U$  and  $P_D$ , there is a greater frequency in the functionally stronger regulatory area,  $P_D$ , but in four ancient samples (Altai, Denisova 3, Vindija, and Ust'-Ishim), they only occur in  $P_U$  (see S3 Table); only Denisova 2 and Mezmaiskaya had no polymorphisms in the ORF. The comparatively lower frequency across all samples reflects the conservation trend noted in primates [19, 20]. A structural analysis of the distribution of polymorphisms via an Exact Test indicated no significant statistical differences among all samples (results not shown). Given the expected frequency of polymorphism (based on the perception of *CCR5* covered by an area of interest—see Table 3 footnotes), there is a significant pattern in the samples. First, modern humans and Denisova 2 have a greater than expected number of polymorphisms in the ORF (even if these are exceeded by polymorphisms in regulatory regions). All samples (except Denisova 2) have a greater than expected number of polymorphisms in the promoter regions.

Prior research found that humans have a potentially unique plasticity in gene expression due to the effect of alternate splicing [57]. The distribution of polymorphisms across gene regions in *Homo* suggests plasticity in gene regulation and expression in response to viral loads, as noted in previous studies [19, 20]. The pattern of immune gene introgression, particularly regulatory haplotypes in the antiviral *OAS* gene cluster [70], has suggested that selective forces in our close relatives operated on expression, not protein variation—same as seen in non-human primate *CCR5* variation [19, 20]—and those adaptations were also useful to humans. Thus, an increase in polymorphisms that allowed plasticity in regulation and expression in *CCR5* makes sense even if it is not due to introgression. Without functional testing, the exact nature of the polymorphisms is not known other than by inference and comparative analysis, as done here. And, without more paleo-genomes to compare, we cannot know if the variation in these genomes represents true species variation but the data presented here indicate that the pattern is not human specific, rather one shared by recent members of *Homo*.

The expectation that extinct hominins and modern humans would share this pattern of increased variation in the regulatory areas of the gene is met in the current study. Our genus has several unique behavioral and genetic adaptations compared to nonhuman primates and these adaptations might hold some avenues for further research. For instance, a genus-wide shift in subsistence activities occurred during the Plio-Pleistocene (roughly 2 million years ago) from opportunistic non-confrontational scavenging to confrontational scavenging and, later, top predatory behaviors; this alteration to hominin-environment interaction brought

Table 2. *Homo CCR5* polymorphisms marked by\*.

| Position | RefSeq      | Ref | Alt | Structure | IKG | Alt | Den3 | Den2 | Mez | Ust | Vin |
|----------|-------------|-----|-----|-----------|-----|-----|------|------|-----|-----|-----|
| 46409845 | Novel       | C   | A   | PU        |     |     |      |      | *   |     |     |
| 46410036 | rs7637813   | G   | A   | PU        | *   | *   | *    | *    |     | *   | *   |
| 46410114 | Novel       | C   | G   | PU        |     |     |      |      | *   |     |     |
| 46410146 | Novel       | G   | A   | PU        |     |     |      |      | *   |     |     |
| 46410306 | rs41499550  | T   | C   | PU        | *   |     | *    | *    |     |     |     |
| 46410494 | rs2856757   | A   | C   | PU        | *   |     |      |      |     | *   |     |
| 46410499 | rs41395049  | T   | C   | PU        | *   |     | *    | *    |     |     |     |
| 46410507 | rs41412948  | C   | T   | PU        | *   | *   |      |      | *   |     | *   |
| 46410936 | rs2734225   | G   | T   | PU        | *   |     |      |      |     | *   |     |
| 46410992 | Novel       | G   | A   | PU        |     |     |      |      | *   |     |     |
| 46411074 | Novel       | C   | T   | PU        |     |     |      |      | *   |     |     |
| 46411080 | Novel       | C   | T   | PU        |     |     |      |      | *   |     |     |
| 46411083 | rs553230690 | C   | T   | PU        | *   |     |      |      | *   |     |     |
| 46411133 | rs41475349  | G   | A   | PU        | *   |     | *    | *    |     |     |     |
| 46411295 | Novel       | C   | T   | PU        |     |     |      |      | *   |     |     |
| 46411390 | Novel       | C   | G   | PU        |     |     |      |      | *   |     |     |
| 46411542 | rs2227010   | G   | A   | PU        | *   | *   | *    | *    |     | *   | *   |
| 46411782 | Novel       | G   | C   | CCR5      |     |     |      | *    |     |     |     |
| 46411784 | Novel       | G   | A   | CCR5      |     |     |      | *    |     |     |     |
| 46411790 | Novel       | T   | G   | CCR5      |     |     |      | *    |     |     |     |
| 46411840 | rs2734648   | G   | T   | CCR5      | *   |     | *    |      |     | *   |     |
| 46411869 | Novel       | G   | A   | CCR5      |     |     |      |      | *   |     |     |
| 46411870 | Novel       | G   | A   | CCR5      |     |     |      |      | *   |     |     |
| 46411935 | rs1799987   | A   | G   | CCR5      | *   | *   | *    | *    | *   | *   | *   |
| 46412208 | Novel       | G   | A   | CCR5      |     |     |      |      | *   |     |     |
| 46412259 | rs1799988   | C   | T   | CCR5      | *   | *   | *    |      | *   | *   | *   |
| 46412271 | Novel       | G   | A   | CCR5      |     |     |      |      | *   |     |     |
| 46412285 | Novel       | C   | T   | CCR5      |     |     |      |      | *   |     |     |
| 46412308 | rs1800023   | A   | G   | CCR5      | *   |     |      |      |     | *   |     |
| 46412495 | Novel       | G   | A   | CCR5      |     |     |      |      | *   |     |     |
| 46412559 | rs1800024   | C   | T   | CCR5      | *   |     |      |      |     | *   |     |
| 46413157 | Novel       | G   | A   | CCR5      |     |     |      | *    |     |     |     |
| 46413192 | Novel       | G   | A   | CCR5      |     |     |      | *    |     |     |     |
| 46413193 | Novel       | G   | A   | CCR5      |     |     |      | *    |     |     |     |
| 46413334 | rs2856762   | C   | T   | CCR5      | *   | *   |      |      | *   |     | *   |
| 46413418 | rs2254089   | C   | T   | CCR5      | *   |     |      |      |     | *   |     |
| 46413629 | Novel       | T   | G   | CCR5      |     |     |      | *    |     |     |     |
| 46413631 | Novel       | G   | T   | CCR5      |     |     |      |      | *   |     |     |
| 46413632 | rs188423028 | G   | A   | CCR5      | *   |     |      |      | *   |     |     |
| 46413633 | Novel       | C   | G   | CCR5      |     |     |      | *    |     |     |     |
| 46413638 | Novel       | A   | G   | CCR5      |     |     |      | *    |     |     |     |
| 46413647 | Novel       | C   | A   | CCR5      |     |     |      |      | *   |     |     |
| 46413743 | rs2856764   | C   | T   | CCR5      | *   |     |      |      |     | *   |     |
| 46413911 | Novel       | T   | C   | CCR5      |     |     |      | *    |     |     |     |
| 46413914 | Novel       | A   | C   | CCR5      |     |     |      | *    |     |     |     |
| 46413927 | Novel       | G   | T   | CCR5      |     |     |      | *    |     |     |     |
| 46413950 | rs2856765   | G   | A   | CCR5      | *   |     |      |      |     | *   |     |

(Continued)

Table 2. (Continued)

| Position | RefSeq      | Ref | Alt | Structure     | IKG | Alt | Den3 | Den2 | Mez | Ust | Vin |
|----------|-------------|-----|-----|---------------|-----|-----|------|------|-----|-----|-----|
| 46414035 | rs41515644  | A   | G   | CCR5          | *   |     |      |      |     | *   |     |
| 46414612 | Novel       | C   | G   | CCR5, PD, ORF |     |     |      | *    |     |     |     |
| 46414618 | rs1800941   | T   | C   | CCR5, PD, ORF | *   |     |      | *    |     |     |     |
| 46414794 | Novel       | T   | A   | CCR5, PD, ORF |     |     |      | *    |     |     |     |
| 46414925 | rs199824195 | T   | C   | CCR5, PD, ORF | *   |     |      | *    |     |     |     |
| 46414926 | Novel       | G   | C   | CCR5, PD, ORF |     |     |      | *    |     |     |     |
| 46414975 | Novel       | G   | T   | CCR5, PD, ORF |     |     |      | *    |     |     |     |
| 46415095 | Novel       | G   | A   | CCR5, PD, ORF |     |     |      |      | *   |     |     |
| 46415202 | Novel       | G   | C   | CCR5, PD, ORF |     |     |      | *    |     |     |     |
| 46415216 | Novel       | T   | C   | CCR5, PD, ORF |     |     |      | *    |     |     |     |
| 46415264 | Novel       | T   | G   | CCR5, ORF     |     |     |      | *    |     |     |     |
| 46415383 | Novel       | G   | C   | CCR5, ORF     |     |     |      | *    |     |     |     |
| 46415453 | Novel       | C   | T   | CCR5, ORF     |     |     |      |      | *   |     |     |
| 46415501 | Novel       | T   | A   | CCR5, ORF     |     |     |      | *    |     |     |     |
| 46415771 | Novel       | C   | T   | CCR5          |     |     |      | *    |     |     |     |
| 46415857 | Novel       | G   | A   | CCR5          |     |     |      | *    |     |     |     |
| 46415858 | Novel       | G   | A   | CCR5          |     |     |      | *    |     |     |     |
| 46416030 | Novel       | G   | A   | CCR5          |     |     |      | *    |     |     |     |
| 46416038 | Novel       | G   | T   | CCR5          |     |     |      | *    |     |     |     |
| 46416216 | rs17765882  | C   | T   | CCR5          | *   |     |      |      |     |     | *   |
| 46416413 | Novel       | G   | C   | CCR5          |     |     |      | *    |     |     |     |
| 46416470 | rs1800874   | G   | T   | CCR5          | *   |     |      |      |     | *   |     |
| 46416512 | Novel       | T   | G   | CCR5          |     |     |      | *    |     |     |     |
| 46416517 | Novel       | A   | G   | CCR5          |     |     |      | *    |     |     |     |
| 46416525 | Novel       | T   | C   | CCR5          |     |     |      | *    |     |     |     |
| 46416633 | Novel       | C   | T   | CCR5          |     |     |      |      | *   |     |     |
| 46417004 | Novel       | C   | T   | CCR5          |     |     |      |      | *   |     |     |
| 46417069 | rs41442546  | C   | A   | CCR5          | *   |     |      |      | *   |     |     |
| 46417219 | Novel       | A   | T   | CCR5          |     |     |      | *    |     |     |     |
| 46417302 | Novel       | T   | C   | CCR5          |     |     | *    |      |     |     |     |
| 46417312 | rs746492    | G   | T   | CCR5          | *   | *   | *    |      |     | *   | *   |
| 46417614 | Novel       | G   | A   | CCR5          |     |     |      |      | *   |     |     |

\* Polymorphism present

<https://doi.org/10.1371/journal.pone.0204989.t002>

hominins into greater and regular contact with animal carcasses [79–84]. Neandertals in Europe have also been shown to be active hunters and foragers [85] who experienced increased

Table 3. Variation in gene structure, polymorphisms as percentage.

|                             | Expected | Actual |        |         |        |        |        |        |  |
|-----------------------------|----------|--------|--------|---------|--------|--------|--------|--------|--|
|                             | RefSeq   | 1kg    | alt    | den3    | den2   | mez    | ust    | vin    |  |
| ORF <sup>1</sup>            | 17.46%   | 22.33% | 0.00%  | 0.00%   | 32.35% | 11.11% | 0.00%  | 0.00%  |  |
| P <sub>U</sub> <sup>2</sup> | 33.80%   | 32.94% | 75.00% | 100.00% | 28.89% | 55.00% | 36.36% | 60.00% |  |

<sup>1</sup> Percent CCR5 covered by ORF

<sup>2</sup> Percent Promoter+CCR5 covered by promoters

<https://doi.org/10.1371/journal.pone.0204989.t003>

pathogen exposure and disease load as a result [48, 49, 51, 86]. Humans and European Neandertals would have shared similar ecological adaptive pressures in Europe—broad and varied—whereas Altai Neandertal (related to European Neandertals) and Denisova would have shared similar ecological adaptive pressures in Siberia with Ust'-Ishim—less varied. Key pathogens year-round in tropical to temperate zones are more likely to be viral (vector-borne) or bacterial (zoonotic) with transmission via interaction with the environment [87]; high latitude pathogens year-round are more likely parasitic due to the reliance on marine mammals [88] and the short season for viral vector reproductive cycles to transmit infection from insects to hominins [35]. Evidence for gene introgression from extinct hominin species to modern humans is clustered (among other domains) in immune system genes [27, 72, 76, 89]; in particular, the *OAS* anti-viral gene cluster on Chromosome 12 shows signatures of positive selection [72, 76], which suggests that adaptation to Eurasian pathogens may have been partly facilitated by prior adaptive mutations to local viral loads. At a minimum, the environmental challenge faced by non-human members of *Homo* facilitated human adaptation to a new environment—a shared challenge with a similar solution.

While previous studies have examined variation in *CCR5*, particularly *CCR5Δ32* which has a more recent origin [36–42], as a product of more recent human-disease interaction, the widespread pattern of increased variation in the gene across the genus *Homo* identified in this study suggests a potential evolutionary adaptation. A key event distinguishing members of the genus *Homo* from the last common ancestor with *Australopithecus* was the shift to confrontational scavenging and, later, hunting; this alteration to human-environment interaction added a new point of disease contact as evidenced by modern data showing hunting bushmeat (which ancient hominins did too [83]) alters disease exposure via introduction of retroviruses and other pathogens [90–94]. Given *CCR5*'s role in both innate and adaptive immune system functioning, its plasticity may have provided an advantage to members of *Homo* across these varied disease ecologies and its potentially greater than normal interaction with the environment in foraging and hunting activities. As more ancient genomes become sequenced, we can have more robust data with which to work and invest resources into functional testing and experimentation of what function these polymorphisms might have had.

## Supporting information

### S1 Table. Altai neandertal introgression data.

(DOCX)

### S2 Table. Variants in 1000 Genomes.

(DOCX)

### S3 Table. Raw counts of variants per sample.

(DOCX)

## Author Contributions

**Conceptualization:** Kara C. Hoover.

**Data curation:** Kara C. Hoover.

**Formal analysis:** Kara C. Hoover.

**Investigation:** Kara C. Hoover.

**Methodology:** Kara C. Hoover.

**Project administration:** Kara C. Hoover.

**Resources:** Kara C. Hoover.

**Validation:** Kara C. Hoover.

**Writing – original draft:** Kara C. Hoover.

**Writing – review & editing:** Kara C. Hoover.

## References

1. Allen SJ, Crown SE, Handel TM. Chemokine: receptor structure, interactions, and antagonism. *Annu Rev Immunol.* 2007; 25(1):787–820. <https://doi.org/10.1146/annurev.immunol.24.021605.090529> PMID: 17291188
2. Baggiolini M, Dewald B, Moser B. Interleukin-8 and related chemotactic cytokines—CXC and CC chemokines. *Adv Immunol.* 1994; 55:97–179. PMID: 8304236
3. Samson M, Labbe O, Mollereau C, Vassart G, Parmentier M. Molecular cloning and functional expression of a new human CC-chemokine receptor gene. *Biochemistry.* 1996; 35(11):3362–7. <https://doi.org/10.1021/bi952950g> PMID: 8639485
4. Choe H, Farzan M, Sun Y, Sullivan N, Rollins B, Ponath PD, et al. The beta-chemokine receptors CCR3 and CCR5 facilitate infection by primary HIV-1 isolates. *Cell.* 1996; 85(7):1135–48. PMID: 8674119
5. Dragic T, Litwin V, Allaway GP, Martin SR, Huang Y, Nagashima KA, et al. HIV-1 entry into CD4+ cells is mediated by the chemokine receptor CC-CKR-5. *Nature.* 1996; 381:667–73. <https://doi.org/10.1038/381667a0> PMID: 8649512
6. Alkhatib G, Combadiere C, Broder CC, Feng Y, Kennedy PE, Murphy PM, et al. CC CKR5: a RANTES, MIP-1alpha, MIP-1beta receptor as a fusion cofactor for macrophage-tropic HIV-1. *Science.* 1996; 272(5270):1955–8. PMID: 8658171
7. Doranz BJ, Rucker J, Yi Y, Smyth RJ, Samson M, Peiper SC, et al. A dual-tropic primary hiv-1 isolate that uses fusin and the beta-chemokine receptors CKR-5, CKR-3, and CKR-2b as fusion cofactors. *Cell.* 1996; 85(7):1149–58. [https://doi.org/10.1016/S0092-8674\(00\)81314-8](https://doi.org/10.1016/S0092-8674(00)81314-8). PMID: 8674120
8. Deng H, Liu R, Ellmeier W, Choe S, Unutmaz D, Burkhart M, et al. Identification of a major co-receptor for primary isolates of HIV-1. *Nature.* 1996; 381:661–6. <https://doi.org/10.1038/381661a0> PMID: 8649511
9. Farzan M, Choe H, Martin K, Marcon L, Hofmann W, Karlsson G, et al. Two orphan seven-transmembrane segment receptors which are expressed in CD4-positive cells support Simian Immunodeficiency Virus infection. *J Exp Med.* 1997; 186(3):405. PMID: 9236192
10. Liao F, Alkhatib G, Peden KWC, Sharma G, Berger EA, Farber JM. *STRL33*, A novel chemokine receptor-like protein, functions as a fusion cofactor for both macrophage-tropic and T Cell line-tropic HIV-1. *J Exp Med.* 1997; 185(11):2015. PMID: 9166430
11. Alkhatib G, Liao F, Berger EA, Farber JM, Peden KWC. A new SIV co-receptor, *STRL33*. *Nature.* 1997; 388:238. <https://doi.org/10.1038/40789> PMID: 9230431
12. Deng H, Unutmaz D, KewalRamani VN, Littman DR. Expression cloning of new receptors used by simian and human immunodeficiency viruses. *Nature.* 1997; 388:296–300. <https://doi.org/10.1038/40894> PMID: 9230441
13. Esteves PJ, Abrantes J, Van Der Loo W. Extensive gene conversion between CCR2 and CCR5 in domestic cat (*Felis catus*). *Int J Immunogenet.* 2007; 34(5):321–4. <https://doi.org/10.1111/j.1744-313X.2007.00716.x> PMID: 17845301
14. Vázquez-Salat N, Yuhki N, Beck T, O'Brien SJ, Murphy WJ. Gene conversion between mammalian CCR2 and CCR5 chemokine receptor genes: A potential mechanism for receptor dimerization. *Genomics.* 2007; 90(2):213–24. <https://doi.org/10.1016/j.ygeno.2007.04.009>. PMID: 17544254
15. Perelygin AA, Zharkikh AA, Astakhova NM, Lear TL, Brinton MA. Concerted Evolution of Vertebrate CCR2 and CCR5 Genes and the Origin of a Recombinant Equine CCR5/2 Gene. *J Hered.* 2008; 99(5):500–11. <https://doi.org/10.1093/jhered/esn029> PMID: 18502735
16. Carmo CR, Esteves PJ, Ferrand N, van der Loo W. Genetic variation at chemokine receptor CCR5 in leporids: alteration at the 2nd extracellular domain by gene conversion with CCR2 in *Oryctolagus*, but not in *Sylvilagus* and *Lepus* species. *Immunogenetics.* 2006; 58(5):494–501. <https://doi.org/10.1007/s00251-006-0095-4> PMID: 16596402
17. Abrantes J, Carmo CR, Matthee CA, Yamada F, van der Loo W, Esteves PJ. A shared unusual genetic change at the chemokine receptor type 5 between *Oryctolagus*, *Bunolagus* and *Pentalagus*. *Conservation Genetics.* 2011; 12(1):325–30. <https://doi.org/10.1007/s10592-009-9990-1>

18. Schroder KEE, Carey MP, Venable PA. Methodological Challenges in Research on Sexual Risk Behavior: II. Accuracy of Self-Reports. *Ann Behav Med*. 2003; 26(2):104–23. [https://doi.org/10.1207/S15324796ABM2602\\_03](https://doi.org/10.1207/S15324796ABM2602_03) PMID: 14534028
19. Mummidi S, Ahuja SS, McDaniel BL, Ahuja SK. The human CC chemokine receptor 5 (CCR5) gene. Multiple transcripts with 5'-end heterogeneity, dual promoter usage, and evidence for polymorphisms within the regulatory regions and noncoding exons. *J Biol Chem*. 1997; 272(49):30662–71. PMID: 9388201
20. Mummidi S, Bamshad M, Ahuja SS, Gonzalez E, Feuillet PM, Begum K, et al. Evolution of human and non-human primate CC chemokine receptor 5 gene and mRNA. Potential roles for haplotype and mRNA diversity, differential haplotype-specific transcriptional activity, and altered transcription factor binding to polymorphic nucleotides in the pathogenesis of HIV-1 and simian immunodeficiency virus. *J Biol Chem*. 2000; 275(25):18946–61. <https://doi.org/10.1074/jbc.M000169200> PMID: 10747879
21. Zhang YW, Ryder OA, Zhang YP. Sequence evolution of the CCR5 chemokine receptor gene in primates. *Mol Biol Evol*. 1999; 16(9):1145–54. <https://doi.org/10.1093/oxfordjournals.molbev.a026205> PMID: 10486970
22. Zhang Y-W, Ryder OA, Zhang Y-P. Intra- and interspecific variation of the *CCR5* gene in higher primates. *Mol Biol Evol*. 2003; 20(10):1722–9. <https://doi.org/10.1093/molbev/msg198> PMID: 12949140
23. Ribeiro IP, Schrago CG, Soares EA, Pissinatti A, Seuanez HN, Russo CAM, et al. CCR5 chemokine receptor gene evolution in New World monkeys (Platyrrhini, Primates): implication on resistance to lentiviruses. *Infections, Genetics and Evolution*. 2005; 5:271–80.
24. Dean M, Carrington M, Winkler C, Huttley GA, Smith MW, Allikmets R, et al. Genetic restriction of HIV-1 infection and progression to AIDS by a deletion allele of the *CKR5* structural gene. Hemophilia Growth and Development Study, Multicenter AIDS Cohort Study, Multicenter Hemophilia Cohort Study, San Francisco City Cohort, ALIVE Study. *Science*. 1996; 273(5283):1856–62. <https://doi.org/10.1126/science.273.5283.1856> PMID: 8791590
25. Samson M, Libert F, Doranz BJ, Rucker J, Liesnard C, Farber CM, et al. Resistance to HIV-1 infection in caucasian individuals bearing mutant alleles of the CCR-5 chemokine receptor gene. *Nature*. 1996; 382(6593):722–5. <https://doi.org/10.1038/382722a0> PMID: 8751444
26. Temme S, Zacharias M, Neumann J, Wohlfromm S, König A, Temme N, et al. A novel family of human leukocyte antigen class II receptors may have its origin in archaic human species. *J Biol Chem*. 2014; 289(2):639–53. <https://doi.org/10.1074/jbc.M113.515767> PMID: 24214983
27. Abi-Rached L, Jobin MJ, Kulkarni S, McWhinnie A, Dalva K, Gragert L, et al. The shaping of modern human immune systems by multiregional admixture with archaic humans. *Science*. 2011; 334(6052):89–94. <https://doi.org/10.1126/science.1209202> PMID: 21868630
28. Huang Y, Paxton WA, Wolinsky SM, Neumann AU, Zhang L, He T, et al. The role of a mutant CCR5 allele in HIV-1 transmission and disease progression. *Nat Med*. 1996; 2(11):1240–3. PMID: 8898752
29. Zimmerman PA, Buckler-White A, Alkhatib G, Spalding T, Kubofcik J, Combadiere C, et al. Inherited resistance to HIV-1 conferred by an inactivating mutation in CC chemokine receptor 5: studies in populations with contrasting clinical phenotypes, defined racial background, and quantified risk. *Mol Med*. 1997; 3(1):23–36. PMID: 9132277
30. Glass WG, McDermott DH, Lim JK, Lekhong S, Yu SF, Frank WA, et al. CCR5 deficiency increases risk of symptomatic West Nile Virus infection. *J Exp Med*. 2006; 203(1):35–40. <https://doi.org/10.1084/jem.20051970> PMID: 16418398
31. Lim JK, Louie CY, Glaser C, Jean C, Johnson B, Johnson H, et al. Genetic deficiency of chemokine receptor CCR5 is a strong risk factor for symptomatic West Nile virus infection: a meta-analysis of 4 cohorts in the US epidemic. *J Infect Dis*. 2008; 197(2):262–5. <https://doi.org/10.1086/524691> PMID: 18179388
32. Diamond MS, Klein RS. A genetic basis for human susceptibility of West Nile Virus. *Trends Microbiol*. 2006; 14:287–89. <https://doi.org/10.1016/j.tim.2006.05.004> PMID: 16750369
33. Novembre J, Galvani AP, Slatkin M. The geographic spread of the CCR5 Delta32 HIV-resistance allele. *PLoS Biol*. 2005; 3(11):e339. <https://doi.org/10.1371/journal.pbio.0030339> PMID: 16216086
34. Novembre J, Johnson T, Bryc K, Kutalik Z, Boyko AR, Auton A, et al. Genes mirror geography within Europe. *Nature*. 2008; 456(7218):98–101. <https://doi.org/10.1038/nature07331> PMID: 18758442
35. Hoover KC, Barker CM. West Nile virus, climate change, and circumpolar vulnerability. *Wiley Interdiscip Rev Clim Change*. 2016; 7(2):283–300. <https://doi.org/10.1002/wcc.382>
36. Carrington M, Kissner T, Gerrard B, Ivanov S, O'Brien SJ, Dean M. Novel alleles of the chemokine-receptor gene *CCR5*. *Am J Hum Genet*. 1997; 61(6):1261–7. <https://doi.org/10.1086/301645> PMID: 9399903
37. Stephens JC, Reich DE, Goldstein DB, Shin HD, Smith MW, Carrington M, et al. Dating the origin of the CCR5-Delta32 AIDS-resistance allele by the coalescence of haplotypes. *Am J Hum Genet*. 1998; 62(6):1507–15. <https://doi.org/10.1086/301867> PMID: 9585595

38. Galvani AP, Slatkin M. Evaluating plague and smallpox as historical selective pressures for the CCR5-Delta 32 HIV-resistance allele. *Proc Natl Acad Sci U S A*. 2003; 100(25):15276–9. <https://doi.org/10.1073/pnas.2435085100> PMID: 14645720
39. Limborska SA, Balanovsky OP, Balanovskaya EV, Slominsky PA, Schadrina MI, Livshits LA, et al. Analysis of CCR5Delta32 geographic distribution and its correlation with some climatic and geographic factors. *Hum Hered*. 2002; 53(1):49–54. doi: 48605. <https://doi.org/10.1159/000048605> PMID: 11901272
40. Sabeti PC, Walsh E, Schaffner SF, Varilly P, Fry B, Hutcheson HB, et al. The case for selection at CCR5-Delta32. *PLoS Biol*. 2005; 3(11):e378. <https://doi.org/10.1371/journal.pbio.0030378> PMID: 16248677
41. Hummel S, Schmidt D, Kremeyer B, Herrmann B, Oppermann M. Detection of the CCR5-Delta32 HIV resistance gene in Bronze Age skeletons. *Genes Immun*. 2005; 6(4):371–4. <https://doi.org/10.1038/sj.gene.6364172> PMID: 15815693
42. Meccas J, Franklin G, Kuziel WA, Brubaker RR, Falkow S, Mosier DE. Evolutionary genetics: CCR5 mutation and plague protection. *Nature*. 2004; 427:606. <https://doi.org/10.1038/427606a> PMID: 14961112
43. Winder IC, Devès MH, King GCP, Bailey GN, Inglis RH, Meredith-Williams M. Evolution and dispersal of the genus *Homo*: A landscape approach. *J Hum Evol*. 2015; 87:48–65. <https://doi.org/10.1016/j.jhevol.2015.07.002> PMID: 26235482
44. Brinkworth JF. Infectious Disease and the Diversification of the Human Genome. *Hum Biol*. 2017; 89(1). <https://digitalcommons.wayne.edu/humbiol/vol89/iss1/4>.
45. Quintana-Murci L. Genetic and epigenetic variation of human populations: An adaptive tale. *Comptes Rendus Biologies*. 2016; 339(7):278–83. <https://doi.org/10.1016/j.crv.2016.04.005>.
46. Karlsson EK, Kwiatkowski DP, Sabeti PC. Natural selection and infectious disease in human populations. *Nature reviews Genetics*. 2014; 15(6):379–93. <https://doi.org/10.1038/nrg3734> PMID: 24776769
47. Berens AJ, Cooper TL, Lachance J. The genomic health of ancient hominins. *Hum Biol Oceania*. 2017; 89(1). <https://digitalcommons.wayne.edu/humbiol/vol89/iss1/2>.
48. Hardy K, Buckley S, Collins MJ, Estalrich A, Brothwell D, Copeland L, et al. Neanderthal medics? Evidence for food, cooking, and medicinal plants entrapped in dental calculus. *Die Naturwissenschaften*. 2012; 99(8):617–26. <https://doi.org/10.1007/s00114-012-0942-0> PMID: 22806252
49. Hardy K, Radini A, Buckley S, Blasco R, Copeland L, Burjachs F, et al. Diet and environment 1.2 million years ago revealed through analysis of dental calculus from Europe's oldest hominin at Sima del Elefante, Spain. *The Science of Nature*. 2016; 104(1):2. <https://doi.org/10.1007/s00114-016-1420-x> PMID: 27981368
50. Weyrich LS, Duchene S, Soubrier J, Arriola L, Llamas B, Breen J, et al. Neanderthal behaviour, diet, and disease inferred from ancient DNA in dental calculus. *Nature*. 2017; 544(7650):357–61. <https://doi.org/10.1038/nature21674> PMID: 28273061
51. Hardy K, Buckley S, Huffman M. Doctors, chefs or hominin animals? Non-edible plants and Neanderthals. *Antiquity*. 2016; 90(353):1373–9. <https://doi.org/10.15184/aqy.2016.134>
52. Essig A, Hofmann D, Munch D, Gayathri S, Kunzler M, Kallio PT, et al. Copsin, a novel peptide-based fungal antibiotic interfering with the peptidoglycan synthesis. *J Biol Chem*. 2014; 289(50):34953–64. <https://doi.org/10.1074/jbc.M114.599878> PMID: 25342741
53. Marciniak S, Perry GH. Harnessing ancient genomes to study the history of human adaptation. *Nature Reviews Genetics*. 2017; advance online publication. <https://doi.org/10.1038/nrg.2017.65> <http://www.nature.com/nrg/journal/vaop/ncurrent/abs/nrg.2017.65.html> PMID: 28890534
54. Olalde I, Allentoft ME, Sanchez-Quinto F, Santpere G, Chiang CWK, DeGiorgio M, et al. Derived immune and ancestral pigmentation alleles in a 7,000-year-old Mesolithic European. *Nature*. 2014; advance online publication. <https://doi.org/10.1038/nature12960> <http://www.nature.com/nature/journal/vaop/ncurrent/abs/nature12960.html> PMID: 24463515
55. Lalueza-Fox C, Römpler H, Caramelli D, Stäubert C, Catalano G, Hughes D, et al. A melanocortin 1 receptor allele suggests varying pigmentation among Neanderthals. *Science*. 2007; 318(5855):1453–5. <https://doi.org/10.1126/science.1147417> PMID: 17962522
56. Racimo F, Sankararaman S, Nielsen R, Huerta-Sanchez E. Evidence for archaic adaptive introgression in humans. *Nature Review Genetics*. 2015; 16(6):359–71. <https://doi.org/10.1038/nrg3936> <http://www.nature.com/nrg/journal/v16/n6/abs/nrg3936.html> PMID: 25963373
57. Niemitz C. The evolution of the upright posture and gait—a review and a new synthesis. *Die Naturwissenschaften*. 2010; 97(3):241–63. <https://doi.org/10.1007/s00114-009-0637-3> PMID: 20127307
58. Huerta-Sanchez E, Jin X, Asan Bianba Z, Peter BM, Vinckenbosch N, et al. Altitude adaptation in Tibetans caused by introgression of Denisovan-like DNA. *Nature*. 2014; 512:194–7. <https://doi.org/10.1038/>

- nature13408]<http://www.nature.com/nature/journal/v512/n7513/abs/nature13408.html>. PMID: 25043035
59. Livingstone FB. Anthropological implications of sickle cell gene distribution in West Africa. *American Anthropologist*. 1958; 60(3):533–62.
  60. Abecasis GR, Auton A, Brooks LD, DePristo MA, Durbin RM, Handsaker RE, et al. An integrated map of genetic variation from 1,092 human genomes. *Nature*. 2012; 491(7422):56–65. <https://doi.org/10.1038/nature11632> PMID: 23128226
  61. Prüfer K, Racimo F, Patterson N, Jay F, Sankararaman S, Sawyer S, et al. The complete genome sequence of a Neanderthal from the Altai Mountains. *Nature*. 2014; 505(7481):43–9. <https://doi.org/10.1038/nature12886> PMID: 24352235
  62. Slon V, Viola B, Renaud G, Gansauge M-T, Benazzi S, Sawyer S, et al. A fourth Denisovan individual. *Science Advances*. 2017; 3(7). <https://doi.org/10.1126/sciadv.1700186> PMID: 28695206
  63. Prüfer K, de Filippo C, Grote S, Mafessoni F, Korlević P, Hajdinjak M, et al. A high-coverage Neandertal genome from Vindija Cave in Croatia. *Science*. 2017; <https://doi.org/10.1126/science.aao1887> PMID: 28982794
  64. Castellano S, Parra G, Sánchez-Quinto FA, Racimo F, Kuhlwilm M, Kircher M, et al. Patterns of coding variation in the complete exomes of three Neandertals. *Proceedings of the National Academy of Sciences*. 2014; 111(18):6666–71. <https://doi.org/10.1073/pnas.1405138111> PMID: 24753607
  65. Fu Q, Li H, Moorjani P, Jay F, Slepchenko SM, Bondarev AA, et al. Genome sequence of a 45,000-year-old modern human from western Siberia. *Nature*. 2014; 514(7523):445–9. <https://doi.org/10.1038/nature13810> PMID: 25341783
  66. Reich D, Green RE, Kircher M, Krause J, Patterson N, Durand EY, et al. Genetic history of an archaic hominin group from Denisova Cave in Siberia. *Nature*. 2010; 468(7327):1053–60. <https://doi.org/10.1038/nature09710> PMID: 21179161
  67. Meyer M, Kircher M, Gansauge M-T, Li H, Racimo F, Mallick S, et al. A High-Coverage Genome Sequence from an Archaic Denisovan Individual. *Science*. 2012; 338(6104):222–6. <https://doi.org/10.1126/science.1224344> PMID: 22936568
  68. Dabney J, Meyer M, Pääbo S. Ancient DNA damage. *Cold Spring Harb Perspect Biol*. 2013; 5(7): a012567. <https://doi.org/10.1101/cshperspect.a012567> PMID: 23729639
  69. Briggs AW, Stenzel U, Meyer M, Krause J, Kircher M, Pääbo S. Removal of deaminated cytosines and detection of in vivo methylation in ancient DNA. *Nucleic Acids Res*. 2010; 38(6):e87–e. <https://doi.org/10.1093/nar/gkp1163> PMID: 20028723
  70. Gittelman Rachel M, Schraiber Joshua G, Vernot B, Mikacenic C, Wurfel Mark M, Akey Joshua M. Archaic hominin admixture facilitated adaptation to Out-of-Africa environments. *Curr Biol*. 2016; 26(24):3375–82. <https://doi.org/10.1016/j.cub.2016.10.041> PMID: 27839976
  71. Hu Y, Ding Q, Wang Y, Xu S, He Y, Wang M, et al. Investigating the evolutionary importance of Denisovan introgressions in Papua New Guineans and Australians. *bioRxiv*2015.
  72. Mendez FL, Watkins JC, Hammer MF. Neandertal origin of genetic variation at the cluster of OAS immunity genes. *Mol Biol Evol*. 2013; 30(4):798–801. <https://doi.org/10.1093/molbev/mst004> PMID: 23315957
  73. Pimenoff VN, Mendes de Oliveira C, Bravo IG. Transmission between archaic and modern human ancestors during the evolution of the oncogenic human papillomavirus 16. *Mol Biol Evol*. 2017; 34(1):4–19. <https://doi.org/10.1093/molbev/msw214> PMID: 28025273
  74. Racimo F, Marnetto D, Huerta-Sánchez E. Signatures of archaic adaptive introgression in present-day human populations. *Mol Biol Evol*. 2017; 34(2):296–317. <https://doi.org/10.1093/molbev/msw216> PMID: 27756828
  75. Vernot B, Tucci S, Kelso J, Schraiber JG, Wolf AB, Gittelman RM, et al. Excavating Neandertal and Denisovan DNA from the genomes of Melanesian individuals. *Science*. 2016; 352(6282):235–9. <https://doi.org/10.1126/science.aad9416> PMID: 26989198
  76. Sankararaman S, Mallick S, Dannemann M, Prüfer K, Kelso J, Paabo S, et al. The genomic landscape of Neandertal ancestry in present-day humans. *Nature*. 2014; 507:354–7. <https://doi.org/10.1038/nature12961> PMID: 24476815
  77. Holmes EC. On the origin and evolution of the human immunodeficiency virus (HIV). *Biol Rev Camb Philos Soc*. 2001; 76(2):239–54. PMID: 11396848
  78. Galvani AP, Slatkin M. Evaluating plague and smallpox as historical selective pressures for the CCR5-Δ32 HIV-resistance allele. *Proc Natl Acad Sci U S A*. 2003; 100(25):15276–9. <https://doi.org/10.1073/pnas.2435085100> PMID: 14645720
  79. Brantingham PJ. Hominid–carnivore coevolution and invasion of the predatory guild. *J Anthropol Archaeol*. 1998; 17(4):327–53. <https://doi.org/10.1006/jaar.1998.0326>.

80. de Heinzelin J, Clark JD, White T, Hart W, Renne P, WoldeGabriel G, et al. Environment and behavior of 2.5-million-year-old Bouri hominids. *Science*. 1999; 284(5414):625–9. PMID: [10213682](https://pubmed.ncbi.nlm.nih.gov/10213682/)
81. Domínguez-Rodrigo M. Hunting and scavenging by early humans: The state of the debate. *J World Prehist*. 2002; 16(1):1–54. <https://doi.org/10.1023/A:1014507129795>
82. Domínguez-Rodrigo M, Pickering TR. Early hominid hunting and scavenging: a zooarcheological review. *Evol Anthr*. 2003; 12(6):275–82. <https://doi.org/10.1002/evan.10119>
83. Shipman P, Bosler W, Davis KL, Behrensmeier AK, Dunbar RIM, Groves CP, et al. Butchering of giant geladas at an Acheulian site *Current Anthropology*. 1981; 22(3):257–68. <https://doi.org/10.1086/202663>
84. Moleón M, Sánchez-Zapata JA, Margalida A, Carrete M, Owen-Smith N, Donázar JA. Humans and Scavengers: The Evolution of Interactions and Ecosystem Services. *Bioscience*. 2014; 64(5):394–403. <https://doi.org/10.1093/biosci/biu034>
85. Richards MP, Pettitt PB, Trinkaus E, Smith FH, Paunović M, Karavanić I. Neanderthal diet at Vindija and Neanderthal predation: The evidence from stable isotopes. *Proc Natl Acad Sci U S A*. 2000; 97(13):7663–6. <https://doi.org/10.1073/pnas.120178997> PMID: [10852955](https://pubmed.ncbi.nlm.nih.gov/10852955/)
86. Hardy BL. Climatic variability and plant food distribution in Pleistocene Europe: Implications for Neanderthal diet and subsistence. *Quaternary Science Reviews*. 2010; 29(5–6):662–79. <https://doi.org/10.1016/j.quascirev.2009.11.016>
87. Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, Gittleman JL, et al. Global trends in emerging infectious diseases. *Nature*. 2008; 451(7181):990–3. <https://doi.org/10.1038/nature06536> PMID: [18288193](https://pubmed.ncbi.nlm.nih.gov/18288193/)
88. Dudley JP, Hoberg EP, Jenkins EJ, Parkinson AJ. Climate change in the North American Arctic: A one health perspective. *EcoHealth*. 2015; 12(4):713–25. <https://doi.org/10.1007/s10393-015-1036-1> PMID: [26070525](https://pubmed.ncbi.nlm.nih.gov/26070525/)
89. Mendez Fernando L, Watkins Joseph C, Hammer Michael F. A haplotype at STAT2 introgressed from Neanderthals and serves as a candidate of positive selection in Papua New Guinea. *Am J Hum Genet*. 2012; 91(2):265–74. <https://doi.org/10.1016/j.ajhg.2012.06.015>. PMID: [22883142](https://pubmed.ncbi.nlm.nih.gov/22883142/)
90. Steve A-M, Ahidjo A, Placide M-K, Caroline F, Mukulumanya M, Simon-Pierre N-K, et al. High prevalences and a wide genetic diversity of simian retroviruses in non-human primate bushmeat in rural areas of the Democratic Republic of Congo. *EcoHealth*. 2017; 14(1):100–14. <https://doi.org/10.1007/s10393-016-1202-0> PMID: [28050688](https://pubmed.ncbi.nlm.nih.gov/28050688/)
91. van Vliet N, Moreno J, Gomez J, Zhou W, Fa JE, Golden C, et al. Bushmeat and human health: assessing the evidence in tropical and sub-tropical forests. *Ethnobiology and Conservation*. 2017; 6. <https://doi.org/10.15451/ec2017-04-6.3-1-45>
92. Kurpiers LA, Schulte-Herbrüggen B, Ejotre I, Reeder DM. Bushmeat and emerging infectious diseases: Lessons from Africa. In: Angelici FM, editor. *Problematic Wildlife: A Cross-Disciplinary Approach*; [https://doi.org/10.1007/978-3-319-22246-2\\_24](https://doi.org/10.1007/978-3-319-22246-2_24) Cham: Springer International Publishing; 2016. p. 507–51.
93. Mossoun A, Calvignac-Spencer S, Anoh AE, Pauly MS, Driscoll DA, Michel AO, et al. Bushmeat hunting and zoonotic transmission of simian T-Lymphotropic Virus 1 in tropical West and Central Africa. *J Virol*. 2017; 91(10):e02479–16. <https://doi.org/10.1128/JVI.02479-16> PMID: [28298599](https://pubmed.ncbi.nlm.nih.gov/28298599/)
94. Chen M, Reed RR, Lane AP. Acute inflammation regulates neuroregeneration through the NF- $\kappa$ B pathway in olfactory epithelium. *Proc Natl Acad Sci U S A*. 2017; 114(30):8089–94. <https://doi.org/10.1073/pnas.1620664114> PMID: [28696292](https://pubmed.ncbi.nlm.nih.gov/28696292/)