

RESEARCH ARTICLE

Prevalence and prognosis of acutely ill patients with organ failure at arrival to hospital: A systematic review

Peter Bank Pedersen^{1,2*}, Asbjørn Hrobjartsson³, Daniel Lykke Nielsen¹, Daniel Pilsgaard Henriksen^{1,4}, Mikkel Brabrand^{1,5}, Annmarie Touborg Lassen^{1,2}

1 Department of Emergency Medicine, Odense University Hospital, Odense, Denmark, **2** Institute of Clinical Research, University of Southern Denmark, Odense, Denmark, **3** Centre for Evidence-Based Medicine, University of Southern Denmark & Odense University Hospital, Odense, Denmark, **4** Department of Respiratory Medicine, Odense University Hospital, Odense, Denmark, **5** Department of Emergency Medicine, Hospital of South West Jutland, Esbjerg, Denmark

* Peter.Bank.Pedersen@rsyd.dk



Abstract

OPEN ACCESS

Citation: Pedersen PB, Hrobjartsson A, Nielsen DL, Henriksen DP, Brabrand M, Lassen AT (2018) Prevalence and prognosis of acutely ill patients with organ failure at arrival to hospital: A systematic review. PLoS ONE 13(11): e0206610. <https://doi.org/10.1371/journal.pone.0206610>

Editor: Wisit Cheungpasitporn, University of Mississippi Medical Center, UNITED STATES

Received: April 4, 2018

Accepted: October 16, 2018

Published: November 1, 2018

Copyright: © 2018 Pedersen et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: No external funding has been received for this review. PBP was supported by the University of Southern Denmark and the Region of Southern Denmark and DLN was supported by the Foundation of December 17, 1981, but they were not involved in any aspect of this systematic review, and had no influence on the study results or publication, and no role in study design, data

Introduction

Patients in an emergency department are diverse. Some are more seriously ill than others and some even arrive in multi-organ failure. Knowledge of the prevalence of organ failure and its prognosis in unselected patients is important from a diagnostic, hospital planning, and from a quality evaluation point of view, but is not reported systematically.

Objectives

To analyse the prevalence and prognosis of new onset organ failure in unselected acute patients at arrival to hospital.

Methods

A systematic review of studies of prevalence and prognosis of acutely ill patients with organ failure at arrival to hospital. We searched PubMed, Cochrane Library, Embase and Cinahl, and read references in included studies. Two authors decided independently on study eligibility and extracted data. Results were summarised qualitatively.

Results

Four studies were included with a total of 678,960 patients. The number of different organ failures reported in the studies ranged from one to six, and the settings were emergency departments and wards. The definitions of organ failure varied between studies. The prevalence of organ failure was 7%, 14%, 14%, and 23%, and in-hospital mortality was 5%, 11% and 15% respectively. The relative risk of in-hospital mortality for patients with organ failure compared to patients without organ failure varied from 2.58 to 8.65. Numbers of organ failures per 1,000 visits varied from 71 to 256.

collection or analysis. ATL received an unrestricted grant from the private philanthropic fund TrygFonden given to the University of Southern Denmark.

Competing interests: The authors have declared that no competing interests exist.

Conclusion

The results of this review indicate that clinicians have good reasons to be alert when a patient arrives to the emergency department; as a state of organ failure seems both frequent and highly severe.

However, most studies identified were performed in patients after a diagnosis was established, and only very few studies were performed in unselected patients.

Systematic review registration number

PROSPERO: [CRD42017060871](https://doi.org/10.1186/1745-6215-42017060871).

Introduction

Rationale

The unselected patients who attend an Emergency Department(ED) are a heterogeneous group, and identifying the critically ill as soon as possible is crucial. Some patients have one or more organ failures with diverse severity, aetiologies, and affecting different organs. Knowledge of the prevalence and prognosis for these patients are relevant to all healthcare workers in contact with critically ill patients, and are important from a diagnostic, hospital planning, and from a quality point of view, but is not reported systematically. Furthermore definitions of organ failure appear diverse based on different scores or systems [1–3]. Most studies with focus on organ failure are in selected groups of patients, but before patients arrive to the wards or Intensive Care Unit (ICU), most of them have passed through the ED, where the primary evaluation and treatment is performed.

Prevalence of organ failure has been demonstrated in 51–72% of patients in the ICU, mortality was 40–60% in one to five years, and early improvement in organ function improved prognosis [4–7]. Similar results are found in sepsis patients, where organ failure was a risk factor of deterioration [8, 9], and organ failure was persistent and long-term [10].

An efficient method of identifying patients with organ failure has yet to be discovered, but is very much needed, because patients with overt clinical evidence of organ dysfunction receive better care than patients without a clear presentation of their organ failure [11]. It is essential for these acutely ill patients, with organ failure, to be identified and treated as early as possible, preferably in the emergency department, or even in the prehospital setting to prevent deterioration, ICU-transfer, death or long-term morbidity. This has the potential to improve both short-term and long-term prognosis, in these patients [6, 9].

With this review we wanted to highlight knowledge in the early identification of organ failure patients at the hospital doorstep, with the aim to guide future optimisation of the organisation there meets these omnipresent patients.

Objectives

Our main research objectives were to assess the prevalence and prognosis of new onset organ failures in acutely ill patients at arrival.

Methods

Methods were described and published in our protocol developed ahead of this systematic review, and covered below [12].

Protocol and registration

This systematic review is reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement [13, 14]. To describe the conduct, a protocol was developed in advance according to the Preferred Reporting Items for Systematic Reviews and Meta-Analysis Protocols (PRISMA-P) 2015 statement, and with inspiration from the Cochrane Handbook for Systematic Reviews of Interventions [15–17]. The protocol was registered in PROSPERO, as proposed in the reporting guideline, registration number: CRD42017060871.

Eligibility criteria

Study designs. Eligible studies were randomized and non-randomized controlled trials and observational studies, assessing the prevalence of new organ failure, or prognosis of acutely ill patients, at arrival. Case reports and studies with less than 100 patients were excluded.

Participants/Population. We included studies on acutely ill adult patients (adult age limit set by included studies), with one or more of following organ failures: circulatory, respiratory, renal, cerebral, hepatic or coagulation failure. Studies on specific condition or disease were excluded, and studies on both children and adults were eligible, as long as adult data were presented.

Outcomes. We included studies on acutely ill adult patients (adult age limit set by included studies), with one or more of following organ failures: circulatory, respiratory, renal, cerebral, hepatic or coagulation failure. Studies on specific condition or disease were excluded, and studies on both children and adults were eligible, as long as adult data were presented.

Setting. We included studies of patients, who arrived at an emergency department, an acute medical unit, a trauma centre, a general ward, or other entrances for acutely ill patients. Studies were excluded if patients arrived directly at an intensive care unit.

Language. The search were performed without any language filtering, but the included words were all English, and publications in other languages were included in our search, but none were eligible.

Information sources

Electronic databases, references in included studies and authors' personal files, were searched. The databases were: PubMed, Cochrane Library, Embase and Cinahl, and the protocol database, PROSPERO, were searched for ongoing or recently completed systematic reviews on similar topics. The last search was run on 15 November 2017.

Search

Search strategy was developed by input from authors, and with help from an information specialist from The Medical Research Library at University of Southern Denmark.

The reference lists of included studies were scanned for eligible studies. A similar search was performed at all databases, only adapted the exact subjects and syntax, which fit in the particular database. A research record table is included in [S1 Table](#).

Study selection

One author performed title and abstract screening, and excluded obviously ineligible studies. Remaining studies were read in full length, independent and in duplicate, by two review

authors. Disagreements were discussed, at a face to face meeting. Reasons for excluding full text studies were documented and presented in [S2 Table](#).

For evaluation of an included study, conducted in our own research unit, we contacted a researcher outside our unit to double-check the inclusion, to minimize the potential bias.

Data collection process

We contacted the authors of the study conducted in our research unit and were provided with additional information and data. Data from other included studies were collected directly from the publication and some processing was performed to fill in missing data according to our table, and by transforming percentages into numbers and numbers into percentages.

Data items

From included studies bibliographical and study description data, patient characteristics, and data related to prevalence and prognosis of organ failure, were extracted, using a data extraction form.

Risk of bias in individual studies

To assess the risk of bias within included studies, the Quality in Prognosis Studies (QUIPS) tool for prognostic studies and the Newcastle-Ottawa Scale (NOS) for observational studies were used. The QUIPS tool rates 6 bias domains; study participation, study attrition, prognostic factor measurement, outcome measurement, study confounding and statistical analysis and reporting, as having high, moderate, or low risk of bias.[18] The NOS evaluate selection, comparability and outcome in case-control and cohort studies by assigning stars.[19] Two authors assessed every included study for bias independently, neither of the assessors was blinded to the studies, and disagreements were resolved by discussion. For studies describing both prevalence and prognosis, both risks of bias assessment tools were used.

Planned methods of analysis

Reasons for heterogeneity according to clinical (participants and outcomes) and methodological (design and risk of bias) characteristics were explored. On the basis of scoping searches prior to protocol development, we did not expect to perform meta-analysis due to anticipated heterogeneity. However, we performed statistical test for heterogeneity (Cochran's Q) and degree of heterogeneity were described with the I-square statistic, and because of substantial and statistically significant heterogeneity (I-square >60%, $P < 0.10$), a meta-analysis with the aim of assessing a weighted estimate were not performed [20]. Statistical analyses were performed in Stata version 15.

A qualitative synthesis of the included studies based on tables and graphs, to sum up results and findings were presented [21].

In a subgroup of patients with circulatory failure, sub-group descriptions were performed based at available data.

Confidence in cumulative evidence

Certainty of evidence was provided, inspired by the Grading of Recommendations Assessment (GRADE approach) depending on the basic design of the included studies and any downgrading decisions as: high, moderate, low or very low. Following assessment criteria were used, when considering downgrading the certainty of the evidence: risk of bias, inconsistency, indirectness, imprecision, and publication bias to decide how to grade the certainty of evidence

[22, 23]. The assessments were performed by two authors, independently, and disagreements were resolved by discussion.

Results

Study selection

The search conducted 15 November 2017 provided a total of 14,878 citations. Another four citations were identified through reference lists of included studies. Forty-one articles were read in full length and Appendix 2 provides an overview of full-text screening and reasons for excluding articles from the review. The proportion of agreement was 90.2% between the two Authors performing full-text screening. Thirty-seven studies did not meet the inclusion criteria, or fell under the exclusion criteria, leaving four studies for inclusion in the systematic review Fig 1 [24–27]. No unpublished relevant studies were obtained, and the PROSPERO search identified no reviews on similar topics.

Study characteristics

The four included studies were cohort studies published in English between 2013 and 2015. Two studies were conducted in the United States and two in Europe, Denmark and United Kingdom, and the inclusion periods were between 14 days and four years.

The studies included covered a total population of 678,960 patients, and between 45.8% and 60% were males.

The numbers of organ failures included in the studies ranged from one to six. The prognostic outcome also varied; three studies presented in-hospital mortality and one study presented ICU-transfer; while the last study presented no prognostic outcome.

The settings were emergency departments and wards, and the different organ failure definitions for each study, are provided in S1 Text (Table 1) [24–27].

Results of individual studies

The four studies included presented organ failure prevalence ranging from 6.5% to 23.1% (Table 2). Only three studies presented mortality data on organ failure patients, and none presented longer follow up than in-hospital. The in-hospital mortality ranged from 5.3% to 14.7% in the group of patients with organ failure and from 1.1% to 4.4% in the patients without organ failure (Table 2) [24–27].

Syntheses of results

Significant heterogeneity between studies in both analyses was detected, prevalence; $I^2 = 100\%$, and mortality; $I^2 = 99.3\%$, and according to predefined criteria, no meta-analysis was performed. Relative risk of in-hospital mortality for patients with organ failure compared to patients without is presented in Table 2 and Fig 2.

Numbers of organ failures were calculated per 1,000 visits and varied from 71 to 256. Prevalence and mortality for patients with one to at least four organ failures are presented in Table 3, calculated based on data from included studies, and from prevalence-data collected from the authors of Lindvig et al.

Three of the included studies described circulatory organ failure, two published data on in-hospital mortality at 12.3% and 41.5% respectively. Further data on patients with organ failure following trauma, bleeding, cardiac or sepsis were not published in the included studies.




Fig 1. [28]: PRISMA-flow diagram of selection process, from search to inclusion.

<https://doi.org/10.1371/journal.pone.0206610.g001>

Risk of bias within studies

All four studies reported prevalence of organ failure and three of the studies reported in-hospital mortality. They were judged as having low to moderate risk of bias [S3 Table](#).

Table 1. Characteristics of the four included studies, for organ failure definitions, see S1 Text.

Study	Year	Country	Design	Period	Setting	Population, N	Age	Gender	Type of organ failure included
Benns et al	2013	United States of America	Cohort study	2006	Trauma and non-trauma centres in 38 US states	396,276 injured patients	64 (IQR 42–81), median	Male 48%	Respiratory, Circulatory, Hepatic, Renal
Challiner et al	2014	United Kingdom	Cohort study	2 x 7 days, September and February	Emergency admissions to Manchester Royal Infirmary	745 emergency admissions	NS	Male 54.6%	Renal
Churpek et al	2015	United States of America	Cohort study	November 2008—January 2013	Wards at five University Hospitals	269,951 ward patients	60/61, mean	Male 60%	Cerebral, Coagulatory, Renal, Respiratory, Circulatory, Hepatic
Lindvig et al	2014	Denmark	Cohort study	August 2009—August 2011	Acute Medical Ward at Odense University Hospital	11,988 first time admissions	66 (range 15–103), median	Male 45.8%	Renal, Respiratory, Circulatory

<https://doi.org/10.1371/journal.pone.0206610.t001>

Confidence in cumulative evidence

The outcomes prevalence, based on all four studies, and mortality, based on three studies, was graded as having low certainty of evidence, mainly because of the inconsistency between studies. The outcome ICU-transfer was also graded as low certainty of evidence due to serious risk of bias and few events, based on one study S4 Table.

Discussion

Four eligible studies were included in our systematic review, which only provided an indication of an answer to our research question. The studies covered in total 678,960 patients from EDs and wards, with a prevalence of organ failure between 6.5% and 23.1%, and in-hospital mortality between 5.3% and 14.7%. As expected and without regard for the differences between included studies, prognosis worsens by increasing number of organ failures per patient.

Specific conditions or diseases, or arrival at ICU were the most frequent reasons for exclusion at full-text screening, particular studies on sepsis patients, trauma patients or patients with suspected infection.

Due to significant heterogeneity between studies, meta-analysis was not performed, but results were presented in text and tables. Overall the included studies were judged as having moderate risk of bias and the certainty of evidence were graded as low.

Organ failure appeared frequent and was a very serious condition, and the anticipation is that all health care providers in an acute setting meet organ failure patients frequently. The

Table 2. Results of individual studies, total population included, prevalence of organ failure, in-hospital mortality and relative risk of death for patients with compared to patient without organ failure. NS = Not specified.

Study	Year	Population, N	Population, patients with organ failure, total N (%)	In-hospital mortality, patients with organ failure, N (%)	In-hospital mortality, patients without organ failure, N (%)	Relative risk, organ failure/no organ failure, in-hospital mortality, (CI 95%)
Benns et al	2013	396,276	25,758 (6.5)	3,788 (14.7)	6,299 (1.7)	8.65 (8.32–8.99)
Challiner et al	2014	745	106 (14.2)	12 (11.3)	28 (4.4)	2.58 (1.36–4.92)
Churpek et al	2015	269,951	36,767 (13.6)	1,934 (5.3)	2,565 (1.1)	4.78 (4.51–5.07)
Lindvig et al	2014	11,988	2,769 (23.1)	NS	NS	NS

<https://doi.org/10.1371/journal.pone.0206610.t002>




Fig 2. Forrest plot, relative risk of in-hospital mortality, patients with organ failure compared to patients without organ failure.

<https://doi.org/10.1371/journal.pone.0206610.g002>

health care personnel in emergency departments have to pass attention to baseline health, comorbidities, and clinical presentation to recognize patients with organ failure. But as revealed in this review, lack of research prevented us from the possibility of presenting an exact estimate for prevalence and prognosis; the included studies were too heterogeneous.

Most important this review highlighted large knowledge gaps surrounding the prevalence and prognosis of these omnipresent patients with organ failure. This had the potential to bring forward where to put the efforts at the hospital doorstep to recognize and treat organ failure patients as early as possible. The mortality associated with organ failure, the risk of ICU-transfer and even an accepted and consistent definition of organ failures in the Emergency Department remains topics for future research.

To explain heterogeneity we were looking at study characteristics, and organ failure definitions turned out very different. Two studies used definitions adapted from the 2001 Sepsis consensus definitions, one study used three different classifications of acute kidney injury, where organ failure at arrival were not possible to extract from presented data, and one study defined

Table 3. Prevalence of organ failures per 1000 visits, patients with 1, 2, 3, and 4 organ failures, and in-hospital mortality for patients with 1, 2, 3, and 4 organ failures. NS = Not specified.

Author	Number of organ failures, N/1000 visits	1 organ failure, % (N)	2 organ failures, % (N)	3 organ failures, % (N)	4 organ failures, % (N)	1 organ failure, in-hospital mortality, % (N)	2 organ failures, in-hospital mortality, % (N)	3 organ failures, in-hospital mortality, % (N)	4 organ failures, in-hospital mortality, % (N)
Bennis et al	71	92.3% (23771)	6.6% (1702)	1 % (259)	0.1% (26)	12.4% (2948)	39.4% (671)	59.6% (169)	NS
Challiner et al	142	100% (106)	NS	NS	NS	11.3% (12)	NS	NS	NS
Churpek et al	159	85.5% (31448)	12.6% (4627)	1.7% (617)	0.2% (75)	4.1% (1289)	10.4% (481)	21.3% (131)	44.6% (33)
Lindvig et al	256	89.8% (1486)	9.4% (261)	0.8% (22)	NS	NS	NS	NS	NS

<https://doi.org/10.1371/journal.pone.0206610.t003>

organ failures based on ICD-9 diagnostic codes, where patients without obvious organ failure might be missed. Previous studies have revealed that sepsis diagnosis based on symptoms and clinical findings at arrival, had higher prevalence than discharge diagnosis [29].

Other characteristics which differed across studies were numbers of organ failures included, and the setting. Two studies included patients from the emergency department, and two studies included ward patients, furthermore the organ failures studied ranged from one to six. One study presented a very broad definition of acute kidney injury and included no further organ failure, while another included ward patients with six different organ failures. One study presented a prevalence of organ failure at 23.1%, based on unselected patients at the acute medical ward. Studies including patients with suspected infection in the ED, presented a prevalence of organ failure at 48.7% [30], where mortality was 3–6% [31, 32], the latter in accordance with one included study, but only half of another. In patients with sepsis, the prevalence of organ failure has been described between 17 and 47%, and mortality between 5% and 11% [33, 34], a finding in accordance with the results of this review. If narrowing the inclusion of patients to severe sepsis the in-hospital mortality increases, as expected [35–37].

Finally there could be some differences between pre-hospital care organisations in the studies, according to criteria for admission to ICU, number of available ICU-beds, and more seriously ill patients might be admitted directly to the biggest centres. This information is not described in the included studies.

Early improvement better the prognosis in ICU patients, and as presented in this review, with increasing number of organ failures, prognosis deteriorates [4–7]. Maybe focus at organ failure patients at arrival could reduce number of flowing organ failures, lower the severity, diminish need of ICU-transfer, and thereby improve prognosis. Until further data are available, healthcare professionals in acute settings should bring careful attention and assessment to the presence of organ failure in the meeting with an acutely ill patient.

Strengths

This systematic review was conducted on the basis of an accepted reporting guideline, and a protocol was created, registered and published in advance.

The electronic database search was very broad and was developed with help from an information specialist.

The full-text screening was performed independent and in duplicate, by two review authors, with a proportion of agreement at 90.2%. Likewise the data extraction, risk of bias assessment, and certainty of evidence grading were performed independent and in duplicate by two review authors.

Limitations

Eligible studies indexed in searched electronic databases and references of included studies were discovered, but studies might be missed if indexed in other electronic databases.

Only one author conducted title and abstract screening, and when only four studies were included in final analysis, this revealed a risk of bias if just one or two eligible studies were missed. However, we did not identify any of such studies in the reference lists of the included papers.

Furthermore; we were not able to conduct a meta-analysis due to heterogeneity between studies, and we did not contact authors of three included studies to provide complete data, and data to complete planned sub-group analyses.

One study provided data on acute kidney injury instead of failure, and it was not possible to limit data extraction to failure.

Conclusion

The results of this review indicate that clinicians have good reasons to be alert when a patient arrives to the emergency department; as a state of organ failure seems both frequent and highly severe.

However, most studies identified were performed in patients after a diagnosis was established, and only very few studies were performed in unselected patients.

Supporting information

S1 Checklist.

(DOC)

S1 Text. Organ failure definitions for each study included in the systematic review.

(DOCX)

S1 Table. Full search for Prevalence and Prognosis of Acutely Ill Patients with Organ Failure at Arrival: A Systematic Review.

(DOCX)

S2 Table. Full-text screening with reasons for exclusion.

(DOCX)

S3 Table. Risk of bias assessment.

(DOCX)

S4 Table. Quality of evidence, GRADE.

(DOCX)

Acknowledgments

The authors thank; Jesper Hallas, Professor, from the Department of Clinical Biochemistry and Pharmacology, Odense University Hospital, Odense, Denmark, for reading and double-check the inclusion of Lindvig et al, and Mette Brandt Eriksen, Information Specialist from The Medical Research Library at University of Southern Denmark, for search strategy help.

Author Contributions

Conceptualization: Peter Bank Pedersen, Asbjørn Hrobjartsson, Annmarie Touborg Lassen.

Data curation: Peter Bank Pedersen, Daniel Lykke Nielsen.

Formal analysis: Peter Bank Pedersen.

Investigation: Daniel Lykke Nielsen.

Methodology: Asbjørn Hrobjartsson, Annmarie Touborg Lassen.

Project administration: Peter Bank Pedersen.

Supervision: Asbjørn Hrobjartsson, Daniel Pilsgaard Henriksen, Mikkel Brabrand, Annmarie Touborg Lassen.

Validation: Peter Bank Pedersen, Daniel Lykke Nielsen.

Writing – original draft: Peter Bank Pedersen, Asbjørn Hrobjartsson, Daniel Lykke Nielsen, Daniel Pilsgaard Henriksen, Mikkel Brabrand, Annmarie Touborg Lassen.

Writing – review & editing: Peter Bank Pedersen, Asbjørn Hrobjartsson, Daniel Lykke Nielsen, Daniel Pilsgaard Henriksen, Mikkel Brabrand, Annmarie Touborg Lassen.

References

1. Vincent JL, Moreno R, Takala J, Willatts S, De Mendonca A, Bruining H, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On behalf of the Working Group on Sepsis-Related Problems of the European Society of Intensive Care Medicine. *Intensive care medicine*. 1996; 22(7):707–10. PMID: [8844239](#)
2. Vincent JL, de Mendonca A, Cantraine F, Moreno R, Takala J, Suter PM, et al. Use of the SOFA score to assess the incidence of organ dysfunction/failure in intensive care units: results of a multicenter, prospective study. Working group on "sepsis-related problems" of the European Society of Intensive Care Medicine. *Critical care medicine*. 1998; 26(11):1793–800. PMID: [9824069](#)
3. Dellinger RP, Levy MM, Rhodes A, Annane D, Gerlach H, Opal SM, et al. Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock, 2012. *Intensive care medicine*. 2013; 39(2):165–228. <https://doi.org/10.1007/s00134-012-2769-8> PMID: [23361625](#)
4. Bingold TM, Lefering R, Zacharowski K, Meybohm P, Waydhas C, Rosenberger P, et al. Individual Organ Failure and Concomitant Risk of Mortality Differs According to the Type of Admission to ICU—A Retrospective Study of SOFA Score of 23,795 Patients. *PloS one*. 2015; 10(8):e0134329. <https://doi.org/10.1371/journal.pone.0134329> PMID: [26241475](#)
5. Ranzani OT, Zampieri FG, Besen BA, Azevedo LC, Park M. One-year survival and resource use after critical illness: impact of organ failure and residual organ dysfunction in a cohort study in Brazil. *Critical care (London, England)*. 2015; 19:269.
6. Sakr Y, Lobo SM, Moreno RP, Gerlach H, Ranieri VM, Michalopoulos A, et al. Patterns and early evolution of organ failure in the intensive care unit and their relation to outcome. *Critical care (London, England)*. 2012; 16(6):R222.
7. Lone NI, Walsh TS. Impact of intensive care unit organ failures on mortality during the five years after a critical illness. *American journal of respiratory and critical care medicine*. 2012; 186(7):640–7. <https://doi.org/10.1164/rccm.201201-0059OC> PMID: [22837381](#)
8. Angus DC, van der Poll T. Severe sepsis and septic shock. *The New England journal of medicine*. 2013; 369(21):2063.
9. Chen YX, Li CS. Risk stratification and prognostic performance of the predisposition, infection, response, and organ dysfunction (PIRO) scoring system in septic patients in the emergency department: a cohort study. *Critical care (London, England)*. 2014; 18(2):R74.
10. Guirgis FW, Brakenridge S, Sutchu S, Khadpe JD, Robinson T, Westenbarger R, et al. The long-term burden of severe sepsis and septic shock: Sepsis recidivism and organ dysfunction. *The journal of trauma and acute care surgery*. 2016; 81(3):525–32. <https://doi.org/10.1097/TA.0000000000001135> PMID: [27398984](#)
11. Kakebeeke D, Vis A, de Deckere ER, Sandel MH, de Groot B. Lack of clinically evident signs of organ failure affects ED treatment of patients with severe sepsis. *International journal of emergency medicine*. 2013; 6(1):4. <https://doi.org/10.1186/1865-1380-6-4> PMID: [23445761](#)
12. Pedersen PB, Hrobjartsson A, Nielsen DL, Henriksen DP, Brabrand M, Lassen AT. Prevalence and prognosis of acutely ill patients with organ failure at arrival to hospital: protocol for a systematic review. *Systematic reviews*. 2017; 6(1):227. <https://doi.org/10.1186/s13643-017-0622-4> PMID: [29141664](#)
13. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS medicine*. 2009; 6(7):e1000097. <https://doi.org/10.1371/journal.pmed.1000097> PMID: [19621072](#)
14. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JP, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS medicine*. 2009; 6(7):e1000100. <https://doi.org/10.1371/journal.pmed.1000100> PMID: [19621070](#)
15. Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic reviews*. 2015; 4:1. <https://doi.org/10.1186/2046-4053-4-1> PMID: [25554246](#)
16. Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ (Clinical research ed)*. 2015; 349:g7647.

17. Higgins JPT, Green, S. (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5. 1. 0 [Updated March 2011]. Available from www.handbook.cochrane.org; The Cochrane Collaboration; 2011.
18. Hayden JA, van der Windt DA, Cartwright JL, Cote P, Bombardier C. Assessing bias in studies of prognostic factors. *Annals of internal medicine*. 2013; 158(4):280–6. <https://doi.org/10.7326/0003-4819-158-4-201302190-00009> PMID: 23420236
19. Wells GA, Shea B, O'Connell D, Peterson J, Welch V, Losos M, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. Available: http://www.ohrica/programs/clinical_epidemiology/oxford.asp. 2014.
20. Ryan R. Cochrane Consumers and Communication Review Group. Heterogeneity and subgroup analysis in Cochrane Consumers and Communication Group reviews: planning the analysis at protocol stage. <http://ccrg.cochrane.org>. December 2016.
21. Ryan R. Cochrane Consumers and Communication Review Group. Cochrane Consumers and Communication Review Group: data synthesis and analysis. <http://ccrg.cochrane.org> June 2016.
22. Ryan R, Hill S. How to GRADE the quality of the evidence. Cochrane Consumers and Communication Group, available at <http://ccrg.cochrane.org/author-resources>. Version 3.0. December 2016.
23. Guyatt G, Oxman AD, Akl EA, Kunz R, Vist G, Brozek J, et al. GRADE guidelines: 1. Introduction- GRADE evidence profiles and summary of findings tables. *Journal of clinical epidemiology*. 2011; 64(4):383–94. <https://doi.org/10.1016/j.jclinepi.2010.04.026> PMID: 21195583
24. Bennis M, Carr B, Kallan MJ, Sims CA. Benchmarking the incidence of organ failure after injury at trauma centers and nontrauma centers in the United States. *The journal of trauma and acute care surgery*. 2013; 75(3):426–31. <https://doi.org/10.1097/TA.0b013e31829cfa19> PMID: 24089112
25. Challiner R, Ritchie JP, Fullwood C, Loughnan P, Hutchison AJ. Incidence and consequence of acute kidney injury in unselected emergency admissions to a large acute UK hospital trust. *BMC nephrology*. 2014; 15(1) (no pagination)(84).
26. Churpek MM, Zdravetz FJ, Winslow C, Howell MD, Edelson DP. Incidence and Prognostic Value of the Systemic Inflammatory Response Syndrome and Organ Dysfunctions in Ward Patients. *American journal of respiratory and critical care medicine*. 2015; 192(8):958–64. <https://doi.org/10.1164/rccm.201502-0275OC> PMID: 26158402
27. Lindvig KP, Henriksen DP, Nielsen SL, Jensen TG, Kolmos HJ, Pedersen C, et al. How do bacteraemic patients present to the emergency department and what is the diagnostic validity of the clinical parameters; temperature, C-reactive protein and systemic inflammatory response syndrome? *Scandinavian journal of trauma, resuscitation and emergency medicine*. 2014; 22:39. <https://doi.org/10.1186/1757-7241-22-39> PMID: 25027551
28. Moher D, Liberati A, Tetzlaff J, Altman DG. (2009) TPG. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med* 6(7): e1000097. <https://doi.org/10.1371/journal.pmed.1000097> PMID: 19621072
29. Henriksen DP, Laursen CB, Jensen TG, Hallas J, Pedersen C, Lassen AT. Incidence rate of community-acquired sepsis among hospitalized acute medical patients—a population-based survey. *Critical care medicine*. 2015; 43(1):13–21. <https://doi.org/10.1097/CCM.0000000000000611> PMID: 25251760
30. Linder A, Arnold R, Boyd JH, Zindovic M, Zindovic I, Lange A, et al. Heparin-binding protein measurement improves the prediction of severe infection with organ dysfunction in the emergency department. *Critical care medicine*. 2015; 43(11):2378–86. <https://doi.org/10.1097/CCM.0000000000001265> PMID: 26468696
31. Shapiro N, Howell MD, Bates DW, Angus DC, Ngo L, Talmor D. The Association of Sepsis Syndrome and Organ Dysfunction With Mortality in Emergency Department Patients With Suspected Infection. *Annals of emergency medicine*. 2006; 48(5):583–90.e1. <https://doi.org/10.1016/j.annemergmed.2006.07.007> PMID: 17052559
32. Jessen MK, Skibsted S, Shapiro NI. Number of organ dysfunctions predicts mortality in emergency department patients with suspected infection: a multicenter validation study. *European journal of emergency medicine: official journal of the European Society for Emergency Medicine*. 2015.
33. Williams JM, Greenslade JH, McKenzie JV, Chu K, Brown AFT, Lipman J. Systemic Inflammatory Response Syndrome, Quick Sequential Organ Function Assessment, and Organ Dysfunction: Insights From a Prospective Database of ED Patients With Infection. *Chest*. 2017; 151(3):586–96. <https://doi.org/10.1016/j.chest.2016.10.057> PMID: 27876592
34. Quinten VM, van Meurs M, Wolffensperger AE, Ter Maaten JC, Ligtenberg JJ. Sepsis patients in the emergency department: stratification using the Clinical Impression Score, Predisposition, Infection, Response and Organ dysfunction score or quick Sequential Organ Failure Assessment score? *European journal of emergency medicine: official journal of the European Society for Emergency Medicine*. 2017.

35. Levy MM, Artigas A, Phillips GS, Rhodes A, Beale R, Osborn T, et al. Outcomes of the Surviving Sepsis Campaign in intensive care units in the USA and Europe: a prospective cohort study. *The Lancet Infectious diseases*. 2012; 12(12):919–24. [https://doi.org/10.1016/S1473-3099\(12\)70239-6](https://doi.org/10.1016/S1473-3099(12)70239-6) PMID: [23103175](https://pubmed.ncbi.nlm.nih.gov/23103175/)
36. Rohde JM, Odden AJ, Bonham C, Kuhn L, Malani PN, Chen LM, et al. The epidemiology of acute organ system dysfunction from severe sepsis outside of the intensive care unit. *Journal of hospital medicine*. 2013; 8(5):243–7. <https://doi.org/10.1002/jhm.2012> PMID: [23401431](https://pubmed.ncbi.nlm.nih.gov/23401431/)
37. Macdonald SP, Arendts G, Fatovich DM, Brown SG. Comparison of PIRO, SOFA, and MEDS scores for predicting mortality in emergency department patients with severe sepsis and septic shock. *Academic emergency medicine: official journal of the Society for Academic Emergency Medicine*. 2014; 21(11):1257–63.