

## RESEARCH ARTICLE

# Screening for prevalence of current TB disease and latent TB infection in type 2 diabetes mellitus patients attending a diabetic clinic in an Indian tertiary care hospital

Pradipkumar Arvindbhai Dabhi<sup>1</sup>, Balamugesh Thangakunam<sup>1</sup>, Richa Gupta<sup>1</sup>, Prince James<sup>2</sup>, Nihal Thomas<sup>3</sup>, Dukhabandhu Naik<sup>4</sup>, Devasahayam Jesudas Christopher<sup>1\*</sup>

**1** Department of Pulmonary Medicine, Christian Medical College, Vellore, Tamil Nadu, India, **2** Department of Respiratory Medicine, Christian Medical College, Vellore, Tamil Nadu, India, **3** Department of Endocrinology, Diabetes and Metabolism, Christian Medical College, Vellore, India, **4** Department of Endocrinology, Jawaharlal Institute of postgraduate medical education and Research (JIPMER), Pondicherry, India

\* [djchris@cmcvellore.ac.in](mailto:djchris@cmcvellore.ac.in)



## OPEN ACCESS

**Citation:** Dabhi PA, Thangakunam B, Gupta R, James P, Thomas N, Naik D, et al. (2020) Screening for prevalence of current TB disease and latent TB infection in type 2 diabetes mellitus patients attending a diabetic clinic in an Indian tertiary care hospital. PLoS ONE 15(6): e0233385. <https://doi.org/10.1371/journal.pone.0233385>

**Editor:** Frederick Quinn, The University of Georgia, UNITED STATES

**Received:** February 21, 2020

**Accepted:** May 3, 2020

**Published:** June 5, 2020

**Peer Review History:** PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0233385>

**Copyright:** © 2020 Dabhi et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are available from the Harvard Dataverse (DOI: [10.7910/DVN/HOTY7Y](https://doi.org/10.7910/DVN/HOTY7Y)).

## Abstract

### Background

Diabetes triples the risk of developing tuberculosis (TB). This study was designed to determine the prevalence of past and current TB disease and Latent TB infection (LTBI) in type 2 Diabetes Mellitus (NIDDM) patients.

### Design

This was a prospective descriptive study on all NIDDM patients attending a Diabetic clinic. Detailed history, included details of previous history of TB (Past TB) and symptoms of active TB and a thorough physical exam was also done. When clinical suspicion of TB was present, appropriate investigations were carried out to diagnose 'Current TB'. Subsequently, 200 consecutive patients who were negative for Past and Current TB were screened for Latent TB infection (LTBI) by tuberculin skin test.

### Results

Of 1000 NIDDM patients enrolled, 43(4.3%) had Past TB. Of remaining 957 patients, 50 were evaluated for New TB on the basis of suggestive symptoms and 10(1%) patients were confirmed to have Current TB. Risk factors for Past or Current TB 'DM-TB' in comparison with 'DM Only' group were; male sex (72% VS 57%; P = 0.033), manual laborer (28% VS 15%; P = 0.012), smoking (26% VS 14%; P = 0.015), alcohol consumption (23% VS 9%; P<0.001) & being on treatment with Insulin (40% VS 20%; P<0.001). There was a protective effect with being a home maker (17% VS 37%; P = 0.034) & overweight status (53% VS 71%; P = 0.004). Of the 200 patient without Past or Current TB, who were screened for LTBI, 96

**Funding:** This study was partly supported by the Department of Biotechnology (India) and National Institute for Allergy and Infectious Diseases & National Institutes of Health (USA). The support was administered by CRDF Global under the aegis of the RePORT India consortium. We also acknowledge part funding from Fluid Research grant CMC, Vellore.

**Competing interests:** The authors have declared that no competing interests exist.

(48%) patients were found to have LTBI. Male sex was the only significant risk factor for LTBI (72% VS 59%;  $P = 0.05$ ).

## Conclusion

Past and Current TB was substantial in patients attending a Diabetic Clinic. Active symptom screening for TB in these clinics could lead to increase in case detection and earlier diagnosis.

## Background

While significant gains in global TB control have been achieved, an estimated 10 million people developed TB disease and 1.2 million HIV negative patients died from the disease in 2018 [1]. India is the highest TB burden country accounting for an incidence of 2.69 million cases. As per India TB report 2017 India accounts for a quarter of world's TB cases and Government of India plans to achieve a rapid decline in burden of TB, morbidity and mortality while working towards elimination of TB in India by 2025 [2]. Type 2 diabetes mellitus (DM) triples the risk of developing tuberculosis (TB), and rates of TB are higher in people with DM than in the general population [3]. The link of DM and TB is important in developing countries where TB is endemic and the prevalence of DM is rising. The prevalence of diabetes in adults aged 20 years or older in India increased from 5.5% in 1990 to 7.7% in 2016 [4]. Although the definite pathophysiological mechanism of the effect of DM as a risk factor for TB is unknown, some hypotheses are: depressed cellular immunity, dysfunction of alveolar macrophages, low levels of interferon gamma, pulmonary microangiopathy, and micronutrient deficiency [5,6]. The coexistence of these TB and DM in low and middle income country is an example of a bidirectional association between a communicable and non-communicable disease that increases the dual burden of both the diseases [7].

Some believe that screening for active TB among diabetics could improve case detection, earlier treatment and prevent transmission of disease [8]. The World Health Organization (WHO) and the international union against lung and tuberculosis disease (IUALTD) launched the 'Collaborative framework for the Care and Control of Diabetes and Tuberculosis', with one important recommendation being the routine implementation of bi-directional screening of the two diseases [9]. While studies in India have show that bidirectional screening for DM and TB is feasible [10]. High yield of DM has been reported among TB patients but the yield of TB among DM patients was low and so further research using new, improved TB diagnostic tools was recommended [11]. Other low income countries have reported lower prevalence of DM among patients with TB [12].

In the present study, we aimed to determine the prevalence of Past and Current TB disease and Latent TB infection (LTBI) in patients with NIDDM attending a Diabetic clinic and to describe the prevalence, socio-demographic characteristics, clinical features, microbiological and biochemical variables of these patients.

## Study design

A prospective screening of the DM patients attending the Diabetic clinic of the department of Endocrinology and Diabetes for Past and Current TB disease and LTBI, between September 2014 to April 2015.

## Setting

The study was conducted at the Christian Medical College and Hospital, Vellore, a tertiary care referral hospital in India.

## Inclusion criteria

Patients with NIDDM 18 years or older.

## Exclusion criteria

1. Acute medical illness
2. Known HIV or other immunosuppressive disease
3. Unable or unwilling to comply with study protocol
4. Pregnancy

## Methods

Consecutive NIDDM patients, attending the Diabetic clinic of the Christian Medical College, Vellore, a referral teaching hospital in Southern India, who consented to participate in the trial and provide informed consent were recruited. A detailed history was obtained and in particular past history TB, details of organ involvement and treatment details.

In addition, information about socio-demographic characteristics, history of smoking and alcohol consumption and duration of DM and current medication for DM were obtained. Weight, height, BMI and the most recent blood glucose measurements (fasting & post prandial) and glycosylated hemoglobin (HBA1c) results were recorded. BMI of greater than 23 was considered as high risk for DM in our population [13]. For patients with no chest radiographs within past 1 year, a fresh chest X-Ray (PA view) was obtained. Both these were reviewed by experienced pulmonary physicians and the radiological features were recorded

They fell in one of these 4 categories:

1. Patients with past history of TB (Past TB)
2. Patients with features of pulmonary or extra-pulmonary TB (TB suspects)
3. TB suspects who were proven to have TB (Current TB)
4. Patient with no past history or current clinical features of pulmonary or extra-pulmonary TB (Non TB)

The patients were evaluated as follows:

1. Past TB: Details of organ involvement, history, mode of diagnosis and treatment details were recorded. For those with history of pulmonary TB (PTB), old chest radiographs, if available were reviewed
2. TB suspects: The patients with current symptoms of active TB with the following symptoms:
  - Productive cough for >2 weeks, fever, night sweats, loss of appetite and weight loss.
  - Symptoms and signs of extra-pulmonary TB like lymphadenopathy, pleural effusion, typical skin lesion and joint swelling/pain etc.

- These patients were further investigated to confirm active TB disease. Chest x-ray and 3 sputum examinations (2 for AFB smear and 1 for X-pert MTB/Rif and MGIT culture) were done for suspected active PTB patients. Sputum for AFB smear was done by Ziehl-Neelsen staining method, X-pert MTB/Rif & MGIT cultures were performed as per the manufacturer's instruction. For extra-pulmonary TB (EPTB), appropriate aspirate or tissue was obtained for histopathological evaluation and MGIT culture.
3. Those subsequently diagnosed with active TB (Current TB) were initiated on standard anti TB treatment.
  4. Among these patients with no Past or Current TB (Non TB), 200 consecutive patients were tested for LTBI using one step tuberculin skin test (TST). TST was performed by using tuberculin 2TU solution (PPD RT 23). TST was considered as positive if the indurations were  $\geq 10$  mm and these were categorized as having 'LTBI'. Those who underwent TST test and were negative ( $\leq 10$  mm) and those without LTBI were categorized as 'No LTBI'.
  5. Those with Past and Current were grouped as 'DM-TB' the rest of the patients were grouped as 'DM Only'

The study subjects flowchart is shown in [Fig 1](#). All patient data was recorded on case report form and captured in an electronic data base.

### Statistical analysis

Analysis and statistics Data were extracted from the electronic database and analyzed using SPSS (Statistical Package and Service Solutions, version 16.0, SPSS Inc, Chicago, IL). The socio-demographic characteristics were tabulated. Mean and standard deviations (SD) were calculated. Continuous variables such as age, BMI and duration of DM were converted to categorical variables and compared using the  $\chi^2$  test where appropriate. Levels of significance were set at 5%. The clinical features, diabetic control variables and diabetic treatment details of patients with 'DM-TB' were compared with the 'DM Only' group and the 'LTBI' group was compared with 'No LTBI' group.

### Ethics approval

Ethics approval for the study was obtained from the Institutional review board of the Christian Medical College Vellore.

### Results

Of 1000 DM patients screened for TB, the mean age was 51 [SD 11.3] years, 586 [59%] were male and 414 [41%] were female). In all, 43 (4.3%) had Past TB, and none of these patients had features of active TB (relapse) at screening. Among the other 957 patients, there were 50 patients with clinical suspicion of TB, who were investigated further to confirm/rule out TB. Ten (20%) of those investigated turned out to have Current TB. In all, there were 53 cases of Past and Current TB 'DM-TB'.

[Table 1](#) shows the details of the 10 patients diagnosed to have Current TB-9 of them had pulmonary TB (PTB) and 1 extra pulmonary TB. Out of the 9 pulmonary TB patients, 8 were positive by smear as well as X-pert MTB/Rif, while 1 patient was negative by both these test but had MTB growth on MGIT culture. Out of 9 PTB patients, 8 were drug sensitive and 1 drug resistant (MDR) TB. The only patient with extra-pulmonary TB had cervical lymphadenopathy, confirmed by biopsy showing caseating granuloma on histopathology. MTB did not grow on culture, the patient responded to anti-TB treatment.




Fig 1. Study subjects flowchart.

<https://doi.org/10.1371/journal.pone.0233385.g001>

**Table 1. Types and categories of active TB in DM patients screened for TB in a tertiary care hospital between September 2014 to April 2015.**

Newly diagnosed with TB-‘Current TB’ (n-10)	
Category and type of TB	
Pulmonary TB	9
AFB Smear positive	8
X-pert MTB Rif positive (Rif sensitive-07,Rif resistance-01)	8
Smear and X-pert MTB Rif negative, Culture positive(drug sensitive)	1
Extra pulmonary TB	1

<https://doi.org/10.1371/journal.pone.0233385.t001>

The socio-demographic ‘DM-TB’ and ‘DM Only’ patients are shown in Table 2. Risk factors for Past or Current TB ‘DM-TB’ in comparison with ‘DM Only’ group were; male sex (72% VS 57%;  $P = 0.033$ ), manual laborer (28% VS 15%;  $P = 0.012$ ), smoking (26% VS 14%;  $P = 0.015$ ), alcohol consumption (23% VS 9%;  $P < 0.001$ ), I & being on treatment with Insulin (40% VS 20%;  $P < 0.001$ ). There was a protective effect with being a home maker (17% VS 37%;  $P = 0.034$ ) & overweight status (53% VS 71%;  $P = 0.004$ ).

The clinical variables and DM treatment are compared between ‘DM-TB’ and ‘DM Only’ patients (Table 2). Those with normal BMI was significantly lower in the ‘DM-TB’ group (52.8% VS 71.2%), conversely those with high BMI were higher in the ‘DM Only’ group. With regards to treatment, significantly less patients in the ‘DM-TB’ group were on oral hypoglycemic agents (OHAs) only (26% VS 51%) and significantly more were on insulin (40% VS 20%). There was no difference between the groups with regards to HbA1c, fasting and post prandial plasma glucose.

LTBI was positive in 96 of the 200 patients tested (48%). The ‘LTBI’ group was compared with the ‘No LTBI’ group (Table 3). A higher proportion of ‘LTBI’ patients were male (72% VS 59%;  $P = 0.05$ ). There was no other difference between the groups.

## Discussion

While there are studies that have addressed the prevalence of DM in TB clinics, prevalence of TB among Diabetics are fewer and to the best of our knowledge, there are no studies from our country on LTBI prevalence among NIDDM patients. There are few studies on LTBI in the general population and these reported a higher prevalence of LTBI among patients with DM, with the rates ranging from 28.2 to 42.4%. [14, 15]

While bi-directional screening has been shown to be feasible, A Majumdar et al. recently reported that the implementation of bidirectional screening was poor in the public health system. Adequate staffing, regular training, continuous laboratory supplies for DM diagnosis and widespread publicity were recommended [11].

In all, there were a total of 53 (5.3%) who either had Past or Current TB among patients attending our Diabetic clinic. This is substantial. Among the 10(1%) patients with current TB, 8(80%) had smear-positive disease and were therefore infectious. The prevalence of active TB in this population works out to 1000 per 100000 patients, which is higher than the total TB case notification rate in India. [16]

S. Kumpatlaetal [17], screened 7083 patient of DM for TB and found that only 50(0.7%) patients had current or Past TB, which was substantially lower than what we found (5.3%). Another study by Mave et al. which screened 630 patients in a public sector tertiary care hospital failed to yield any active TB case, using a WHO-recommended questionnaire among people with DM [18]. A thorough screening by a Pulmonary physician, experienced in TB management could at least partly explain the higher yield in our study. There has been few attempts to

Table 2. The socio-demographic and relevant DM treatment and control variables.

Characteristics	DM patients with TB disease	DM patients with no TB disease	p-value
<b>Total</b>	<b>53</b>	<b>947</b>	
Sex			
Male	38 (71.7)	538 (56.8)	0.033
Female	15 (28.3)	409 (43.2)	
Age			
≤45 years	15 (28.3)	230 (24.3)	0.51
46–60 years	30 (56.6)	490 (51.7)	0.487
≥61 years	8 (15.1)	227 (24.0)	0.137
Occupation			
Business	5 (9.4)	111 (11.7)	0.611
Home makers	9 (17.0)	348 (36.8)	0.034
Office job	17 (32.1)	206 (21.7)	0.077
Manual labourer	15 (28.3)	145 (15.3)	0.012
Retired	7 (13.2)	137 (14.5)	0.793
Smoking	14 (26.4)	134 (14.2)	0.015
Alcohol	12 (22.6)	82 (8.7)	<0.001
BMI, Kg/m <sup>2</sup>			
Mean±SD	26	25	0.363
<18	5 (5.7)	28 (3.0)	0.273
18–23	22 (41.5)	245 (25.9)	0.013
>23	28 (52.8)	674 (71.2)	0.004
Duration of DM years			
Mean ± SD	10	7	0.161
<1	4 (7.6)	90 (9.5)	0.645
5-Jan	13 (24.5)	338 (35.7)	0.096
>5	36 (67.9)	272 (54.8)	0.062
Treatment details			
OHA	14 (26.4)	487 (51.4)	<0.001
Insulin	21 (39.6)	193 (20.4)	<0.001
Both	18 (34.0)	267 (28.2)	0.363
Fasting Glucose, Median(IQR)	156 (124, 222)	157 (121, 214)	0.746
Post Prandial Glucose, Median (IQR)	257 (205, 344)	239 (182, 316)	0.084 <sup>†</sup>
HbA1c			
<7	10 (18.9)	243 (25.7)	0.268
7–8.9	23 (43.4)	369 (38.9)	0.514
> = 9	20 (37.7)	335 (35.4)	0.734

<sup>a</sup> Chi-square test was used to assess the association between given factors and two groups

<https://doi.org/10.1371/journal.pone.0233385.t002>

use ultrasound to screen for extra pulmonary tuberculosis in high risk individuals in resource limited settings [19,20].

Rieder [21] has reported that age and sex are strong determinants of TB, with the highest risks being found in elderly people, and this was evident in the study by S. Kumpatla et al. [17] also. Systematic reviews have shown that under nutrition, smoking, diabetes and alcohol consumption are individual risk factors that can double or triple the risk of developing active TB. [22] A large part of the TB burden in India has been attributed to smoking (40%) and DM (15%). [23,24] We also observed that male gender, history of smoking and alcohol

**Table 3. Socio-demographic characteristics of 'LTBI' and 'DM Only' group.**

Characteristics			p-value
	LTBI	No TB	
<b>Total</b>	<b>96</b>	<b>104</b>	
Sex			
Male	69 (71.9)	61 (58.7)	0.05
Female	27 (28.1)	43 (41.3)	
Age			
≤45 years	29 (30.2)	24 (23.1)	0.256
46–60 years	45 (46.9)	56 (53.8)	0.329
≥61 years	22 (22.9)	24 (23.1)	0.973
Occupation			
Business	18 (18.8)	11 (10.6)	0.1
Home maker	20 (20.8)	34 (32.7)	0.058
Job	32 (33.3)	24 (23.1)	0.109
Labourer	16 (16.7)	14 (13.5)	0.527
Retired	10 (10.4)	21 (20.2)	0.056
Smoking	24 (25.0)	17 (16.4)	0.13
Alcohol	15 (15.6)	8 (7.7)	0.079
BMI			
<18	2 (2.1)	7 (6.7)	0.117
18–23	26 (27.1)	35 (33.7)	0.311
>23	68 (70.8)	62 (59.6)	0.097
Duration			
<1 year	6 (6.2)	11 (10.6)	0.265
1–5 years	31 (32.3)	31 (29.8)	0.703
>5 years	59 (61.5)	62 (59.6)	0.784
Treatment			
OHA	51 (53.1)	58 (55.8)	0.702
Insulin	19 (19.8)	19 (19.3)	0.929
Both	26 (27.1)	27 (25.9)	0.848

<sup>a</sup> Chi-square test was used to assess the association between given factors and two groups

<https://doi.org/10.1371/journal.pone.0233385.t003>

consumption were strongly associated with TB diagnosis in Diabetic patients. This makes a stronger case for screening DM patients who have the above additional risk factors.

Compared to no 'DM-TB' patients, 'DM Only' patients had higher body weight [25]. In our study this was true and significantly more 'DM Only' had high BMI.

S. Kumpatlaetal [17], observed that 'DM-TB' group were more likely to have metabolic disease, (>10 years), to be on combination therapy with oral and insulin medication and have higher Hb1AC levels (>7). This indicates that DM patients with TB had more severe disease requiring more treatment for control. [26] Our 'DM-TB' patients were more likely to be on insulin rather than oral hypoglycemic agents only. This could imply that 'DM-TB' patients had more severe DM. It is very important that control of diabetes is given due importance in patients with tuberculosis [27].

Tuberculin skin test surveys in India show a very high annual risk of TB infection. Given the high TB burden of active TB in India, it is not surprising that nearly 40% of Indians are estimated to be latently infected. Despite the likelihood of a large number of LTBI reactivating to active TB, the National program (RNTCP) does not give priority to LTBI detection and



treatment in the public sector. This is true for most of the other high TB burden countries around the world. [28]

We have demonstrated a high (48%) prevalence of LTBI in our diabetic patients, which is higher than the prevalence of LTBI reported for the Indian population. We did not observe any socio-demographic risk factor for latent TB infection in diabetic patients, however the population tested was small. With regard to the duration of Diabetes, treatment and glycemic control, there was no difference between 'DM-LTBI' and 'No LTBI' suggesting that other factors may also be responsible for the susceptibility to LTBI. However, Gerardo Martinez-Aguilar et al [29] found a higher proportion of subjects with HbA1c values of more than 7% in the LTBI (TST-positive) than in the TST negative group.

### Limitations

One of the drawbacks of the study is not having done Interferon Gamma Release Assays due to local issues with availability, when the study commenced.

### Conclusion

The study highlights that screening of DM patients for active TB is feasible and productive in DM clinics in referral hospitals. This could lead to increase on case detection as well as earlier diagnosis. Screening should include assessment for extra-pulmonary TB as well.

### Author Contributions

**Conceptualization:** Pradipkumar Arvindbhai Dabhi, Balamugesh Thangakunam, Devasahayam Jesudas Christopher.

**Data curation:** Devasahayam Jesudas Christopher.

**Formal analysis:** Pradipkumar Arvindbhai Dabhi.

**Funding acquisition:** Devasahayam Jesudas Christopher.

**Investigation:** Pradipkumar Arvindbhai Dabhi, Dukhabandhu Naik.

**Methodology:** Pradipkumar Arvindbhai Dabhi, Balamugesh Thangakunam, Richa Gupta, Prince James, Nihal Thomas, Dukhabandhu Naik, Devasahayam Jesudas Christopher.

**Project administration:** Devasahayam Jesudas Christopher.

**Supervision:** Balamugesh Thangakunam, Richa Gupta, Prince James, Nihal Thomas, Devasahayam Jesudas Christopher.

**Validation:** Balamugesh Thangakunam.

**Writing – original draft:** Pradipkumar Arvindbhai Dabhi, Devasahayam Jesudas Christopher.

**Writing – review & editing:** Balamugesh Thangakunam, Richa Gupta, Prince James, Nihal Thomas, Dukhabandhu Naik, Devasahayam Jesudas Christopher.

### References

1. World Health Organization. Global tuberculosis report, 2012. WHO/HTM/TB/2012.6. Geneva, Switzerland: WHO; 2012: <https://www.who.int/tb/publications/globalreport/en/> Accessed on 22Oct2019
2. (TB India Report 2018.: Ministry of Health and Family Welfare [Internet]. [cited 2020 Apr 9]. Available from: <https://tbcindia.gov.in/showfile.php?lid=3314> Accessed on 22Apr2020

3. Jeon C Y, Murray M B. Diabetes mellitus increases the risk of active tuberculosis: a systematic review of 13 observational studies. *PLOS Med* 2008; 5:e152 <https://doi.org/10.1371/journal.pmed.0050152> PMID: 18630984
4. Tandon N, Anjana RM, Mohan V, Kaur T, Afshin A, Ong K, et al. The increasing burden of diabetes and variations among the states of India: the Global Burden of Disease Study 1990–2016. *The Lancet Global Health*. 2018 Dec 1; 6(12):e1352–62. [https://doi.org/10.1016/S2214-109X\(18\)30387-5](https://doi.org/10.1016/S2214-109X(18)30387-5) PMID: 30219315
5. Webb EA, Hesseling AC, Schaaf HS, Gie RP, Lombard CJ, Spitaels A, Delport S, et al. High prevalence of mycobacterium tuberculosis infection and disease in children and adolescents with type 1 diabetes mellitus. *Int J Tuberc Lung Dis* 2009, 13(7):868–874. PMID: 19555537
6. Ottmani SE, Murray MB, Jeon CY, Baker MA, Kapur A, Lonroth K, et al. Consultation meeting on tuberculosis and diabetes mellitus: meeting summary and recommendations. *Int J Tuberc Lung Dis* 2010, 14(12):1513–1517 PMID: 21180207
7. (Pizzol D, Gennaro FD, Chhaganlal KD, Fabrizio C, Monno L, Putoto G, et al. Tuberculosis and diabetes: current state and future perspectives. *Tropical Medicine & International Health*. 2016; 21(6):694–702.)
8. Jeon CY, Harries AD, Baker MA, Hart JE, Kapur A, Lonroth K, et al. Bi-directional screening for tuberculosis and diabetes: a systematic review. *Trop Med Int Health* 2010, 15(11):1300–1314. <https://doi.org/10.1111/j.1365-3156.2010.02632.x> PMID: 20958887
9. World Health Organization/International Union Against Tuberculosis and Lung Disease. Provisional collaborative framework for care and control of tuberculosis and diabetes. WHO/HTM/TB/2011.15. Geneva, Switzerland: WHO, 2011.
10. Screening of patients with diabetes mellitus for tuberculosis in India. India Diabetes Mellitus–Tuberculosis Study Group\*
11. Majumdar A., Wilkinson E., Rinu P. K., Maung T. M., Bachani D., Punia J. S., et al. Tuberculosis–diabetes screening: how well are we doing? A mixed-methods study from North India *Public Health Action*. 2019 Mar 21; 9(1): 3–10. Published online 2019 Mar 21. PMID: PMC6436489. <https://doi.org/10.5588/pha.18.0048> PMID: 30963036
12. (Pizzol D, Di Gennaro F, Chhaganlal KD, Fabrizio C, Monno L, Putoto G, et al. Prevalence of diabetes mellitus in newly diagnosed pulmonary tuberculosis in Beira, Mozambique. *Afr Health Sci*. 2017 Sep; 17(3):773–779) <https://doi.org/10.4314/ahs.v17i3.20> PMID: 29085405
13. Snehalatha C, Viswanathan V, Ramachandran A. Cutoff Values for Normal Anthropometric Variables in Asian Indian Adults. *Diabetes Care*. 2003 May 1; 26(5):1380–4. <https://doi.org/10.2337/diacare.26.5.1380> PMID: 12716792
14. Leow MK, Dalan R, Chee CB, et al. Latent tuberculosis in patients with diabetes mellitus: prevalence, progression and public health implications. *Exp Clin Endocrinol Diabetes* 2014; 122:528e532.
15. Mansilla Bermejo MJ, Sanz Gil MJ, Moraleda Velasco P, et al. Tuberculin test in diabetic patients in a health center. *Aten Primaria* 1995; 16:154e157.
16. Sachdeva K S, Kumar A, Dewan P, Kumar A M V, Satyanarayana S. New vision for Revised National Tuberculosis Control Programme (RNTCP): universal access—‘Reaching the un-reached’. *Ind J Med Res* 2012; 135: 690–694.
17. Kumpatla S., Sekar A., Achanta S., Sharath B. N., Kumar A. M. V., Harries A. D., Viswanathan V Characteristics of patients with diabetes screened for tuberculosis in a tertiary care hospital in South India *Public Health Action*. 2013 November 4; 3(Suppl 1): S23–S28. PMID: PMC4463146 <https://doi.org/10.5588/pha.13.0035> PMID: 26393064
18. Mave Vidya, Nimkar Smita, Prasad Haridas, Kadam Dileep, Meshram Sushant, Lokhande Rahul, et al. Tuberculosis screening among persons with diabetes mellitus in Pune, India. *BMC Infect Dis*. 2017; 17: 388. Published online 2017 Jun 2. PMID: PMC5457599 <https://doi.org/10.1186/s12879-017-2483-9> PMID: 28577535
19. (Bobbio F, Di Gennaro F, Marotta C, Kok J, Akec G, Norbis L, et al. Focused ultrasound to diagnose HIV-associated tuberculosis (FASH) in the extremely resource-limited setting of South Sudan: a cross-sectional study. *BMJ Open*. 2019 02; 9(4):e027179. <https://doi.org/10.1136/bmjopen-2018-027179> PMID: 30944140
20. Di Gennaro F, Pisani L, Veronese N, Pizzol D, Lippolis V, Saracino A, et al. Potential Diagnostic Properties of Chest Ultrasound in Thoracic Tuberculosis—A Systematic Review. *Int J Environ Res Public Health* [Internet]. 2018 Oct; 15(10). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6210728/>
21. Rieder H. Epidemiologic basis of tuberculosis control. Paris, France: International Union against Tuberculosis and Lung Disease, 1999.

22. Lönnroth K, Castro K G, Chakaya J M, et al. Tuberculosis control and elimination 2010–50: cure, care and social development. *Lancet* 2010; 375: 1814–1829. [https://doi.org/10.1016/S0140-6736\(10\)60483-7](https://doi.org/10.1016/S0140-6736(10)60483-7) PMID: 20488524
23. Hassmiller K. The impact of smoking on population level tuberculosis outcomes. TSRU progress report, 2007. The Hague, The Netherlands: KNCV, 2007.
24. Stevenson C R, Forouhi N G, Roglic G, et al. Diabetes and tuberculosis: the impact of the diabetes epidemic on tuberculosis incidence. *BMC Public Health* 2007; 7: 234 <https://doi.org/10.1186/1471-2458-7-234> PMID: 17822539
25. Ruslami R, Aarnoutse RE, Alisjahbana B, van der Ven AJ, Van Crevel R: Implications of the global increase of diabetes for tuberculosis control and patient care. *Trop Med Int Health* 2010, 15(11):1289–1299. <https://doi.org/10.1111/j.1365-3156.2010.02625.x> PMID: 20955495
26. Restrepo B I, Fisher-Hoch S P, Pino P A, et al. Tuberculosis in poorly controlled type 2 diabetes: altered cytokine expression in peripheral white blood cells. *Clin Infect Dis* 2008; 47: 634–641 <https://doi.org/10.1086/590565> PMID: 18652554
27. (Di Gennaro F, Marotta C, Antunes M, Pizzol D. Diabetes in active tuberculosis in low-income countries: to test or to take care? *Lancet Glob Health*. 2019Jun; 7(6):e707.) [https://doi.org/10.1016/S2214-109X\(19\)30173-1](https://doi.org/10.1016/S2214-109X(19)30173-1) PMID: 31097272
28. sharma S K and Amohan, Kadhiavan T. HIV-TB co-infection: epidemiology, diagnosis and management. *Indian J Med Res* 2005: 121: 550–67 PMID: 15817963
29. Martinez-Aguilar Gerardo, Serrano Carmen J., Castaneda-Delgado Julio Enrique, Associated Risk Factors for Latent Tuberculosis Infection in Subjects with Diabetes *Archives of Medical Research* 46 (2015) 221e227 <https://doi.org/10.1016/j.arcmed.2015.03.009> PMID: 25864989