

CORRECTION

Correction: Geographical variation in the diatom communities associated with loggerhead sea turtles (*Caretta caretta*)

The *PLOS ONE* Staff

In [Table 1](#), the data on seven samples are missing. The publisher apologizes for the error. Please see the correct [Table 1](#) here.



OPEN ACCESS

Citation: The *PLOS ONE* Staff (2021) Correction: Geographical variation in the diatom communities associated with loggerhead sea turtles (*Caretta caretta*). *PLoS ONE* 16(2): e0247350. <https://doi.org/10.1371/journal.pone.0247350>

Published: February 17, 2021

Copyright: © 2021 The PLOS ONE Staff. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Table 1. List of samples and information on loggerhead sea turtles.

Sample code	Body part	ID Tag / Turtle name	Sampling Date	Sex	Weight (kg)	SCL (cm)	CCL (cm)	SCW (cm)	CCW (cm)
Sampling location: Amvrakikos Gulf (Greece) 39° 1' 29" N—39° 1' 44" N; 21° 3' 36" E—21° 4' 19" E									
GRE-01	carapace	Y6343- Y6344	8/1/2018			75.4	78.6	55.0	68.5
GRE-02	skin	Y6343- Y6344							
GRE-03	carapace	Y6366- Y6367	8/1/2018			47.4	51.0	36.3	46.8
GRE-04	skin	Y6366- Y6367							
GRE-05	carapace	Y6368- Y6369	8/1/2018			66.1	69.6	51.5	63.8
GRE-06	skin	Y6368- Y6369							
GRE-07	carapace	Y6370- Y6371	8/1/2018			54.5	58.5	43.7	55.2
GRE-08	skin	Y6370- Y6371							
GRE-09	carapace	M9123- M9124	8/1/2018			50.4	53.2	37.5	47.5
GRE-10	skin	M9123- M9124							
Sampling location: Florida Bay (USA) 24° 55'18" N; 80° 48' 28" W									
FLO-U8	carapace	HA5053—HA5054	6/24/2015	F	58.5	74.1			
FLO-U9	carapace	HB5559—HB5560	6/24/2015		48.0	66.9			
FLO-U10	carapace	HB5668—X7596	6/24/2015	M	89.0	87.3			
FLO-U11	carapace	W1924—W2176	6/24/2015	F	75.7	79.5			
FLO-U12	carapace	HB5581—HB5582	6/25/2015		71.2	78.5			
Sampling location: Kosi Bay (South Africa) 26° 59' 38.9" S; 32° 51' 59.8" E									
SA-33	carapace	ZA0447D - ZA0427D	1/16/2018	F			86.4		83.6
SA-37	carapace	ZA0829D - ZA0828D	1/11/2018	F		80.2	85.7	62.0	78.8
SA-45	carapace	ZA0924D - ZA0186D	1/11/2018	F		73.9	80.2	58.2	74.8
SA-47	carapace	ZA0833D - ZA1923D	1/13/2018	F			91.4		81.8
SA-55	carapace	ZA1406D - ZA1407D	1/18/2018	F		84.0	92.7	64.0	88.5
Sampling location: Adriatic Sea (Croatia) 44° 50' 07" N; 13° 49' 58" E									
CRO-13	carapace	Zanja-Mara	5/30/2016		11.0	44.0	43.0		41.0
CRO-19	carapace	Lunga	8/30/2016		28.8	69.8	68.0		63.2
CRO-21	carapace	Palma	9/30/2016	F	45.0	70.2	68.5		71.5
CRO-24	carapace	Shiggy-Lola	12/5/2016		0.9	24.1	24.5		23.0
CRO-42	carapace	Tylago	2/24/2017		28.2	64.0	63.0		58.0

CCL = curved carapace length, CCW = curved carapace width, SCL = straight carapace length, SCW = straight carapace width

<https://doi.org/10.1371/journal.pone.0247350.t001>

Reference

1. Van de Vijver B, Robert K, Majewska R, Frankovich TA, Panagopoulou A, Bosak S (2020) Geographical variation in the diatom communities associated with loggerhead sea turtles (*Caretta caretta*). PLoS ONE 15(7): e0236513. <https://doi.org/10.1371/journal.pone.0236513> PMID: 32726804