

RESEARCH ARTICLE

# Diversity of fungal pathogens associated with loquat and development of novel virulence scales

Muhammad Fahim Abbas<sup>1\*</sup>, Sana Batool<sup>2</sup>, Sobia Khaliq<sup>3</sup>, Sidra Mubeen<sup>4</sup>, Azziz-ud-Din<sup>5</sup>, Naseeb Ullah<sup>5</sup>, Khalida Zafar<sup>6</sup>, Muhammad Rafiq<sup>7\*</sup>, Abdullah M. Al-Sadi<sup>8</sup>, Saqer S. Alotaibi<sup>9</sup>, Ahmed M. El-Shehawi<sup>9</sup>, Yunzhou Li<sup>10\*</sup>, Ali Tan Kee Zuan<sup>11\*</sup>, Mohammad Javed Ansari<sup>12</sup>

**1** Department of Plant Pathology, Faculty of Agriculture, Lasbela University of Water Agriculture and Marine Sciences (LUAWMS) Uthal, Balochistan, Pakistan, **2** Department of Plant Pathology, Pir Mehr Ali Shah Arid Agriculture University, Rawalpindi, Pakistan, **3** Department of Entomology, Pir Mehr Ali Shah Arid Agriculture University, Rawalpindi, Pakistan, **4** Department of Chemistry, Kutchery Campus, The Women University Multan, Multan, Pakistan, **5** Department of Biotechnology and Genetic Engineering, Hazara University, Mansehra, Pakistan, **6** Department of Botany, Islamia College, Peshawar, Pakistan, **7** Department of Plant Pathology, Faculty of Agriculture, University of the Punjab, Lahore, Pakistan, **8** Department of Plant Sciences, College of Agricultural and Marine Sciences, Sultan Qaboos University, Alkhoud, Oman, **9** Department of Biotechnology, College of Science, Taif University, Taif, Saudi Arabia, **10** Department of Plant Pathology, College of Agriculture, Guizhou University, Guiyang, Guizhou, China, **11** Department of Land Management, Faculty of Agriculture, Universiti Putra Malaysia, Selangor, Malaysia, **12** Department of Botany, Hindu College Moradabad, Mahatma Jyotiba Phule Rohilkhand University Bareilly, Bareilly, India

\* [fahim.abbas@luawms.edu.pk](mailto:fahim.abbas@luawms.edu.pk) (MFA); [fiqakha@gmail.com](mailto:fiqakha@gmail.com) (MR); [tkz@upm.edu.my](mailto:tkz@upm.edu.my) (ATKZ); [liyunzhou2007@126.com](mailto:liyunzhou2007@126.com) (YL)



**OPEN ACCESS**

**Citation:** Fahim Abbas M, Batool S, Khaliq S, Mubeen S, Azziz-ud-Din, Ullah N, et al. (2021) Diversity of fungal pathogens associated with loquat and development of novel virulence scales. PLoS ONE 16(10): e0257951. <https://doi.org/10.1371/journal.pone.0257951>

**Editor:** Shahid Farooq, Harran Üniversitesi, Harran Üniversitesi, TURKEY

**Received:** July 12, 2021

**Accepted:** September 15, 2021

**Published:** October 14, 2021

**Copyright:** © 2021 Fahim Abbas et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the manuscript.

**Funding:** This work was supported by projects of the national Nature Science Foundation (No. 32060679) and projects of Guizhou University (No. GuidapeiYU[2019]52 and No. [2017]50). This study was financed by Taif University Researchers Supporting Project number (TURSP-2020/38), Taif University, Taif, Saudi Arabia. There were no additional external funding involved in the study. The funders had no role in study design, data

## Abstract

Loquat [*Eriobotrya japonica* (Thunb.) Lindl.] is an important fruit crop in Pakistan; however, a constant decline in its production is noted due biotic and abiotic stresses, particularly disease infestation. Fungal pathogens are the major disease-causing agents; therefore, their identification is necessary for devising management options. This study explored Taxila, Wah-Cantt, Tret, Chatar, Murree, Kalar-Kahar, Choa-Saidan-Shah and Khan-Pur districts in the Punjab and Khyber Paktoon Khawa (KPK) provinces of Pakistan to explore the diversity of fungal pathogens associated with loquat. The samples were collected from these districts and their microscopic characterizations were accomplished for reliable identification. *Alternaria alternata*, *Curvularia lunata*, *Lasiodiplodia theobromae*, *Aspergillus flavis*, *Botrytis cinerea*, *Chaetomium globosum*, *Pestalotiopsis mangiferae* and *Phomopsis* sp. were the fungal pathogens infesting loquat in the study area. The isolates of *A. alternata* and *C. lunata* were isolated from leaf spots and fruit rot, while the isolates of *L. theobromae* were associated with twig dieback. The remaining pathogens were allied with fruit rot. The nucleotide evidence of internal transcribed spacer (ITS) regions (ITS1, 5.8S, and ITS2) were computed from all the pathogens and submitted in the database of National Center for Biotechnology Information (NCBI). For multigene analysis, beta-tubulin (BT) gene and glyceraldehyde 3-phosphate dehydrogenase (GAPDH) regions were explored for *A. alternata* and *C. lunata* isolates, respectively. The virulence scales of leaf spots, fruit rot, and twig dieback diseases of loquat were developed for the first time through this study. It is the first comprehensive study with morpho-molecular identification, and newly developed virulence scales of the

collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

fungal pathogens associated with loquat, which improves the understanding of these destructive diseases.

## Introduction

Loquat [*Eriobotrya japonica* (Thunb.) Lindl] belongs to the Rosaceae family. It is a subtropical evergreen tree, and its fruit is used in fresh and processed forms. It is cultivated in Australia, Brazil, China, India, Japan, Madagascar, Mauritius, Mediterranean countries, United States, and Pakistan [1]. China is the largest producer of loquat in the world, while it was introduced in Pakistan from China [2]. In Pakistan, Taxila, Wah-Cantt, Khan-Pur, Tret, Chatar, Muree, Kalar Kahar and Choa Saiden Shah are the loquat growing areas. However, the losses caused by fungi infestation in loquat are still unknown [3]. Loquat is produced on an area of 10000 hectares in Pakistan with annual production 8,823 tons of fruits. However, this production is very low as compared to other loquat growing countries of the world. This low production is attributed to various biotic and abiotic stresses, including disease infestation [4]. Loquat fruit is a good source of minerals and carotenoids, and higher phenolic contents are present in the flesh responsible for rapid browning of the fruits [5].

Loquat fruit has a very short shelf life and sensitive to loss of nutrients, moisture, mechanical and physical damage, and microbial decay [6]. Several fungal pathogens, including *Alternaria alternata*, *Aspergillus niger*, *Botrytis cinerea*, *Colletotrichum gloeosporioides*, *Curlaria lunata*, *Diplocarpon mespili*, *Diplodia natalensis*, *Diplodia seriata*, *Geotrichum candidum*, *Fusarium solani*, *Mucor fragilis*, *Pestalotia* sp., *Phytophthora palmivora* and *Spilocaea pyranthae* infest loquat [7–19].

Several fungi genera are difficult to identify at the species level based on morphological characteristics and require considerable expertise for distinguishing closely-related species since their morphological features overlap [20]. The recent introduction of molecular techniques has advanced the detection, reliable identification and screening of many pathogens [21]. Sequence analysis of internal transcribed spacer (ITS) regions (ITS1, 5.8S and ITS2) has been developed for quicker and specific identification of fungi [22, 23]. Although loquat is an important fruit crop in Pakistan, little is known about fungal pathogens causing leaf spots, fruit rot, and twig dieback in the major loquat growing areas of the country. The successful disease management in fruit crops depends on the species' diagnosis and infestation level of casual organisms. There is no database in Pakistan providing such information. Nevertheless, no literature is currently available on the fungal pathogens associated with loquat. Likewise, the detection of casual organisms through molecular techniques and disease rating scales have not been explored yet.

Therefore, current study was conducted to determine the fungal pathogens associated with loquat in major loquat growing areas of Pakistan. The objectives of this study were; i) to determine fungal pathogens associated with fruit rot, leaf spots and twig dieback of loquat, ii), identification of the pathogens through morpho-molecular methods and iii) and development of the virulence scales for the pathogens through pathogenicity tests. The results will help in future disease management strategies for loquat production areas of the country.

## Materials and methods

### Survey and sample collection

The surveys were conducted during the summer season, 2015 in loquat growing areas of Taxila (33.74° N, 72.78° E), Wah-Cantt (33.77° N, 72.75° E), Tret (33.83° N, 73.30° E), Chatar (33.77°

N, 73.24° E), Murree (33.90° N, 73.39° E), Kalar-Kahar (32.78° N, 72.70° E), Choa-Saidan-Shah (32.71° N, 72.98° E) and Khan-Pur (33.80° N, 72.93° E) districts situated in Punjab and Khyber Pakhtunkhwa (KPK) provinces of Pakistan. A total 20 loquat trees were selected randomly from each district and suspected leaves, fruits, and twigs were collected. The samples were placed in sterile polythene bags mentioned with time, date, and locations [16]. There were no permits required for sample collection as survey studies aimed at collecting the samples and analyzing them within the country is exempt from the permit requirements.

### Isolation and frequency of pathogens

The tissue segments from leaves, fruits, and twigs containing symptomatic and asymptomatic area (5 mm<sup>3</sup>) were surface sterilized with 1% sodium hypochlorite for 1 minute, dipped thrice into sterile distilled water and dried on double layer of sterile filter papers. The segments were transferred to potato dextrose agar (PDA) and incubated at 24 ± 2°C for 7 days in the dark. Cultural characteristics such as colony color, growth rate and texture were checked with three days interval and microscopic characteristics were used for the identification of the pathogens. A single spore method was used for the purification of the pathogens and preserved in silica gel. The frequency of pathogen was computed with the following formula [16].

$$\text{Frequency of a pathogen (\%)} = \frac{\text{Isolates of a specific fungi}}{\text{Total number of fungal isolates}} \times 100$$

### Virulence scale on detached leaf

One to two-years old loquat plants with healthy leaves were purchased from the local nursery. The alternate leaves were disinfected with sodium hypochlorite (1%) and its traces were eliminated with double sterile distilled water. The spore suspension of *C. lunata* (1×10<sup>4</sup> spore/mL) and *A. alternata* (10<sup>6</sup> spores/mL) were sprayed on healthy leaves and the mean infected leaf area (mm) was recorded after 7 days [23].

### Virulence scale on fruit

Healthy fruits were initially surface sterilized with 1% sodium hypochlorite for 30 sec and its traces were eliminated by dipping them into sterile distilled water for 1 min. The fruits were dried on three layers of sterile filter paper. Single spore from each isolate was grown on PDA and 100µL spore suspension from an individual isolate of *C. lunata* (1×10<sup>4</sup> spore/mL), *A. alternata* (10<sup>6</sup> conidia/mL), *A. flavis* (1×10<sup>4</sup> spore/mL), *B. cinerea* (1×10<sup>4</sup> spore/mL), *C. globosum* (1×10<sup>4</sup> spore/mL) and *Phomopsis* sp. (1×10<sup>4</sup> spore/mL) was poured to wounded and non-wounded fruits at top, middle and bottom. Artificially inoculated fruits were incubated at 24 ± 2°C with 75% relative humidity in sterile chambers and observed daily.

### Virulence scale on twigs

Three loquat plants (one to two-years old) with healthy twigs were obtained from the local nursery and superficial wounds (5 mm long, 2 mm deep and 30 cm below the tip) were made on twigs. The spore suspension (100 µL) of *L. theobromae* (1×10<sup>6</sup> spore /mL) was dispensed on wounded twigs and covered with plastic bags. The death of the twig was recorded after 15-day intervals. All treatments had three replications and sterile distilled water (without spore suspension) was used as a negative control. The pathogen was re-isolated from the artificially inoculated fruits, leaves, and twigs and compared with the inoculated culture. The virulence categories such as slightly virulent (+), moderately virulent (++) and highly virulent (+++) of fruit rot, leaf spots, and twig dieback were noted with a newly developed scale (Table 1).

**Table 1. Virulence scales of fungal pathogens causing leaf spots, fruit rot, and twig dieback of loquat.**

Virulence category of leaf spots		
Mean infected leaf area (mm)	Virulence category	
50–70	Highly virulent (+++)	
20–49	Moderately virulent (++)	
01–19	Slightly virulent (+)	
Virulence category of fruit rot		
Wounded fruits (days)	Non-wounded (days)	Virulence category
3	4	Highly virulent (+++)
5	6	Moderately virulent (++)
≥ 7	8	Slightly virulent (+)
Virulence category of twig dieback		
Death of twigs (Days)	Virulence category	
45–60	Highly virulent (+++)	
61–75	Moderately virulent (++)	
>75	Slightly virulent (+)	

<https://doi.org/10.1371/journal.pone.0257951.t001>

## Molecular identification

The genomic DNA from purified fungal isolates was extracted with the standard protocol specified in the PrepMan® Ultra sample preparation reagent. A total volume of 50 µL comprising of 1× PCR reaction buffer, 0.5 µM each forward and reverse primer, 0.2 mM of each deoxynucleotide triphosphate (dNTPs), 2.5 mM magnesium chloride (MgCl<sub>2</sub>), 1 U Taq DNA polymerase, and 20 ng DNA template was used in PCR assay. A negative control (without DNA template) was carried out in the PCR. The PCR conditions of each gene loci were optimized according to the details given in Table 2 and 1Kb DNA ladder was used as a marker. The amplified products were electrophoresed on 1% (w/v) agarose gel stained with ethidium bromide. The amplified products were visualized under ultraviolet light and target DNA fragments were purified with Thermo Scientific Gene<sup>JET</sup> PCR Purification Kit (#K0701) [16].

## Sequencing and sequence analysis

The purified products were sequenced in forward and reverse directions from a DNA facility of Iowa State University, USA, and Macrogen Korea. Both sequences from individual isolate were aligned and manipulated with BioEdit version 7 and final sequences were submitted in the GenBank of National Center for Biotechnological Information (NCBI) to obtain the GenBank accessions numbers (Table 3). The Basic Local Alignment Search Tool (BLAST) network services (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) was used to determine the genetic similarity

**Table 2. PCR reaction and conditions for the amplification of gene loci.**

Locus	Designation	Sequence (5' → 3')	PCR reaction conditions	Fungal Pathogens Identified	Reference
Inter transcribed spacer regions (ITS)	ITS1	TCCGTAGGTGAACCTGCGG	94°C 120 s; 30 cycles of 94°C, 55°C and 72°C 1 min each; 72°C 7 min	<i>C. lunata</i> , <i>A. alternata</i> , <i>B. cinerea</i> , <i>Phomopsi</i> sp., <i>A. flavis</i> , <i>L. theobromae</i> , <i>P. mangiferae</i> and <i>C. globosum</i>	White et al. [22]
	ITS1	TCCTCCGCTTTATTGATATG			
Glyceraldehyde-3-phosphate dehydrogenase (GAPDH)	GAPDH1	ACCACAGTCCATGCCATCAC	95°C 3 min; 30 cycles of 94°C 1 min, 56°C 2 min, 72°C 2 min; 72°C 7 min	<i>C. lunata</i>	Jeon et al. [39]
	GAPDH2	TCCACCACCCCTG TTGCTGTA			
Beta Tubulin (BT)	BT1	AACATGCCGTGAGATTGTAAG	95°C 3 min; 30 cycles of 94°C 1 min, 55°C 1.5 min, 72°C 2 min; 72°C 7 min	<i>A. alternata</i>	
	BT2	TCTGGATGTTGTTGGGAATC			

<https://doi.org/10.1371/journal.pone.0257951.t002>

Table 3. Total number of fungal isolates isolated from major loquat growing areas of Pakistan.

Sr. No.	Species identification	Isolate	Isolation source	Symptoms	Pathogenicity tests				Location	ITS	GPDH	BT
					Wounded fruit	Non-wounded Fruit	Leaf spot	Twig dieback				
1	<i>A. flavis</i>	AF1	Fruit	Fruit rot	+++	+++	-	-	Taxila	MN893384	-	-
2	<i>A. flavis</i>	AF2	Fruit	Fruit rot	++	++	-	-	Taxila	MN893385	-	-
3	<i>A. flavis</i>	AF3	Fruit	Fruit rot	+++	+++	-	-	Wah-Cant	MN893386	-	-
4	<i>A. flavis</i>	AF4	Fruit	Fruit rot	+++	+++	-	-	Khan-Pur	MN893387	-	-
5	<i>B. cinerea</i>	BC	Fruit	Fruit rot	+++	+++	-	-	Taxila	MN891765	-	-
6	<i>C. globosum</i>	CG1	Fruit	Fruit rot	++	++	-	-	Taxila	MN891844	-	-
7	<i>C. globosum</i>	CG2	Fruit	Fruit rot	+++	+++	-	-	WahCant	MN891845	-	-
8	<i>C. lunata</i>	CUR7	Leaf	Leaf spot	-	-	+++	-	Khan-Pur	MN897731	MN894806	-
9	<i>C. lunata</i>	CUR11	Leaf	Leaf spot	-	-	+++	-	Wah-Cant	MN897732	MN894807	-
10	<i>C. lunata</i>	CUR15	Leaf	Leaf spot	-	-	+	-	Taxila	MN897733	MN894808	-
11	<i>C. lunata</i>	CUR19	Fruit	Fruit rot	+++	+++	-	-	Taxila	MN897734	MN894809	-
12	<i>C. lunata</i>	CUR23	Fruit	Fruit rot	+++	+++	-	-	Taxila	MN897735	MN894810	-
13	<i>C. lunata</i>	CUR29	Fruit	Fruit rot	+	+	-	-	Tret	MN897736	MN894811	-
14	<i>C. lunata</i>	CUR33	Fruit	Fruit rot	+++	++	-	-	Taxila	MN897737	MN894812	-
15	<i>C. lunata</i>	CUR45	Fruit	Fruit rot	+++	+++	-	-	Khan-Pur	MN897738	MN894813	-
26	<i>C. lunata</i>	CUR49	Leaf	Leaf spot	-	-	++	-	Wah-Cant	MN897739	MN894814	-
17	<i>C. lunata</i>	CUR53	Fruit	Fruit rot	++	++	-	-	Murree	MN897740	MN894815	-
18	<i>C. lunata</i>	CUR58	Fruit	Fruit rot	+++	+++	-	-	Taxila	MN897741	MN894816	-
19	<i>C. lunata</i>	CUR63	Leaf	Leaf spot	-	-	+++	-	Khan-Pur	MN897742	MN894817	-
20	<i>C. lunata</i>	CUR65	Fruit	Fruit rot	++	++	-	-	Khan-Pur	MN897743	MN894818	-
21	<i>C. lunata</i>	CUR69	Fruit	Fruit rot	+++	+++	-	-	Taxila	MN897744	MN894819	-
22	<i>C. lunata</i>	CUR71	Fruit	Fruit rot	+++	+++	-	-	Kalar-Kahar	MN897745	MN894820	-
23	<i>C. lunata</i>	CUR75	Leaf	Leaf spot	-	-	+++	-	Wah-Cant	MN897746	MN894821	-
24	<i>C. lunata</i>	CUR80	Fruit	Fruit rot	+++	+++	-	-	Khan-Pur	MN897747	MN894822	-
25	<i>C. lunata</i>	CUR99	Leaf	Leaf spot	-	-	++	-	Choa-Saidan-Shah	MN897748	MN894823	-
26	<i>L. theobromae</i>	PAK9	Twig	TDB	-	-	-	+++	Murree	KR259518	-	-
27	<i>L. theobromae</i>	PAK8	Twig	TDB	-	-	-	++	Choa-Saidan-Shah	KR259517	-	-
28	<i>L. theobromae</i>	PK7	Twig	TDB	-	-	-	+++	Wah-Cant	KR259516	-	-
29	<i>L. theobromae</i>	PAK40	Twig	TDB	-	-	-	+++	Taxila	KT312819	-	-
30	<i>L. theobromae</i>	TR-73	Twig	TDB	-	-	-	++	Tret	KR092218	-	-
31	<i>L. theobromae</i>	TR-80	Twig	TDB	-	-	-	++	Tret	KR092217	-	-
32	<i>P. mangiferae</i>	PM	Fruit	Fruit rot	+++	+++	-	-	Taxila	MN888956	-	-
33	<i>Phomopsis</i> sp.	PS1	Fruit	Fruit rot	+++	+++	-	-	Khan-Pur	MN892356	-	-
34	<i>Phomopsis</i> sp.	PS2	Fruit	Fruit rot	+	+	-	-	Taxila	MN892357	-	-
35	<i>Phomopsis</i> sp.	PS3	Fruit	Fruit rot	+	+	-	-	Wah-Cant	MN892358	-	-
36	<i>Phomopsis</i> sp.	PS4	Fruit	Fruit rot	++	++	-	-	WahCant	MN892359	-	-
37	<i>Phomopsis</i> sp.	PS5	Fruit	Fruit rot	+++	+++	-	-	Taxila	MN892360	-	-
38	<i>A. alternate</i>	ALT1	Fruit	Fruit rot	+++	+++	-	-	Taxila	MN944580	-	MT003125
39	<i>A. alternate</i>	ALT2	Fruit	Fruit rot	+++	+++	++	-	Khan-Pur	MN944581	-	MT003126
40	<i>A. alternate</i>	ALT3	Leaf	Leaf spot	++	++	+++	-	Wah-Cant	MN944582	-	MT003127
41	<i>A. alternate</i>	ALT4	Leaf	Leaf spot	++	++	+++	-	Wah-Cant	MN944583	-	MT003128
42	<i>A. alternate</i>	ALT5	Fruit	Fruit rot	+++	+++	++	-	Taxila	MN944584	-	MT003129

(Continued)



Table 3. (Continued)

Sr. No.	Species identification	Isolate	Isolation source	Symptoms	Pathogenicity tests				Location	ITS	GPDH	BT
					Wounded fruit	Non-wounded Fruit	Leaf spot	Twig dieback				
43	<i>A. alternate</i>	ALT6	Leaf	Leaf spot	++	++	++	-	Taxila	MN944585	-	MT003130
44	<i>A. alternate</i>	ALT7	Fruit	Fruit rot	++	++	++	-	Choa-Saidan-Shah	MN944586	-	MT003131
45	<i>A. alternate</i>	ALT8	Leaf	Leaf spot	++	++	+	-	Kalar-Kahar	MN944587	-	MT003132

ITS = Internal Transcribed spacer region, GAPDH = glyceraldehyde 3-phosphate dehydrogenase (GAPDH) BT = beta tubulin, - = negative, + = slightly virulent, ++ = moderately virulent and +++ highly Virulent.

<https://doi.org/10.1371/journal.pone.0257951.t003>

between sequenced and earlier reported isolates. The ITS and GAPDH for *Curvularia* isolates and ITS and BT for *Alternaria* isolate to conduct a multi-locus phylogenetic analysis.

The Molecular Evolutionary (ME) tree of single ITS and combined sequences were inferred for 1000 replicates to assess the stability and robustness of each branch by using the Maximum Likelihood method based on the Tamura 3-parameter model. The tree was computed by Neighbor-Join and BioNJ algorithms to a matrix of pairwise distances estimated using the Maximum Composite Likelihood (MCL) approach. The analysis was conducted in Molecular Evolutionary for Genetic Analysis (MEGA) version 7.0 [24].

## Results

### Symptoms

Soft, dark brown to black lesions were observed in immature and mature fruits and entire fruit rot was recorded at later stage of infection. The necrotic leaf spots were visualized on infected leaves, which varied from brown to yellow. Leaf spots and fruit rot were recorded in all locations, while samples collected from Chatar and Kalar-Kahar were free from twig dieback.

### Morpho-molecular identification of *Curvularia* leaf spots and fruit rot

A total eighteen isolates were morphologically identified as the causal agents of *Curvularia* leaf spots and fruit rot. The fungal colonies were fast-growing with fluffy or velvety and suede-like to downy texture, as well as brown to blackish brown with a black reverse.

Pale brown conidia were slightly curved or cylindrical with dark and large central and three or more transverse septa were formed apically through a pore. The range of average conidial length and width was  $28.02 \pm 2.280 \mu\text{m}$  to  $9.97 \pm 1.99 \mu\text{m}$ , respectively.

The isolates were obtained from Khan-Pur (CUR7, CUR45, CUR63, CUR65, and CUR80), Wah-Cantt (CUR11, CUR49, and CUR75), Tret (CUR29), Murree (CUR53), Kalar-Kahar (CUR71), Choa-Saidan-Shah (CUR99), and Taxila (CUR15, CUR19, CUR23, CUR33, CUR58, and CUR69) with different virulence categories (Table 2). These isolates were confirmed with the amplification of ITS regions (650bp) through PCR assay and sequences were submitted at NCBI with GenBank accession numbers MN897731 to MN897748. The *C. lunata* exhibited 40% pathogen frequency. For multigene identification, GAPDH genes were also amplified through PCR assay and sequences were submitted at NCBI with GenBank accession numbers MN894806 to MN894823.

### Morph-molecular identification of *Alternaria* leaf spots and fruit rot

A total eight isolates of *A. alternata* were isolated from Taxila (ALT1, ALT5, and ALT6), Wah Cantt (ALT3 and ALT4), Khan-Pur (ALT2), Kalar-Kahar (ALT8), and Choa-Saidan-Shah (ALT7). At the initial growth stage, white colonies with slight variations and regular to irregular margins were observed on media. The colonies turned black to golden brown after seven days due to sporulation. The conidia were light to dark brown to black, obclavate, and obpyriform with a short conical or cylindrical beak and the range of conidial length was 9.40 to 39.84  $\mu\text{m}$ . The number of longitudinal septa ranged from 0 to 4, whereas the number of transverse septa ranged from 1 to 6. The isolates formed both beaked and non-beaked conidia. Maximum and minimum conidial breadth was recorded as  $13.98 \pm 1.74 \mu\text{m}$  and  $8.73 \pm 2.49 \mu\text{m}$ , respectively. The ITS regions (650bp) were used for the confirmation of all isolates through PCR assay and sequences were submitted with the GenBank accession number MN944580 to MN944587. The maximum 17.8% pathogen frequency was recorded for *A. alternata* isolates. For multigene identification, BT gene was amplified through PCR assay, and sequences were submitted at NCBI with GenBank accession numbers MT003125 to MT003132.

### Morpho-molecular identification of twig dieback

Dark brown to black colonies in six isolates of *L. theobromae* were observed in PDA and Pycnidia were dark brown to black. Two celled thin-walled oval conidia averaging 25.3  $\mu\text{m}$  length by 12.9  $\mu\text{m}$  width were observed. Six isolates were obtained from Tret (TR-73 and TR-80), Wah-Cantt (PK7), Taxila (PAK40), and Choa-Saidan Shah (PAK8). The amplification of 650bp of ITS regions confirmed these isolates in PCR and sequences were submitted at NCBI with GenBank accession number KR259516 to KR259519 and KR092217 to KR092220. The frequency was 13.3%.

### Morpho-molecular identification of *Phomopsis* fruit rot

Five isolates of *Phomopsis* sp. were recorded from Khan-Pur (PS1), Taxila (PS2 and PS5), and Wah-Cantt (PS3 and PS4). The isolates were yielding colonies with a dark mycelium and abundant black, globose pycnidia 70 to 300  $\mu\text{m}$  in diameter. Hyphae were septate and conidia were hyaline, one-celled, ellipsoid to ovoid (average  $5-11 \times 2-4 \mu\text{m}$ ). The amplification of ITS regions (650bp) in PCR assay confirmed the molecular identification and sequences were submitted with the GenBank accession number MN892356 to MN892360. The frequency was computed as 11.1%.

### Morpho-molecular identification of *Aspergillus* fruit rots

Four isolates of *A. flavis* were obtained from Taxila (AF1 and AF2), Wah-Cantt (AF3), and Khan-Pur (AF4). The isolates were exhibiting yellow colonies which later became bright to dark yellow green. Colonies were granular, flat, often with radial grooves, and covered by a layer of brown to black conidial heads. The conidial heads were radiate (400 to 800  $\mu\text{m}$ ), hyaline conidiophore stipes were coarsely roughened, and conidia (10 spores per sample) were globose to sub-globose (3 to 6  $\mu\text{m}$  in diameter). The 650bp amplification of the ITS region was recorded from each isolate in PCR assay and GenBank accession numbers were obtained as MN893384 to MN893387. The frequency was recorded as 17.8%.

### Morph-molecular identification of *Chaetomium* fruit rots

After 1 week of incubation at  $25 \pm 2^\circ\text{C}$  with a 12-h photoperiod, colonies from two isolates of *Chaetomium globosum* were pale yellow with olivaceous green aerial mycelium, later

producing a greenish pigment that diffused into the medium. Ascospores were olivaceous green, globose or ovoidal (140 to 201  $\mu\text{m} \times 113$  to 229  $\mu\text{m}$ ), with numerous non-branched brownish setae (470  $\mu\text{m}$  in length). The ascospore wall was brown and composed of septate hyphae. Terminal and lateral setae were abundant, brown, and tapering toward the tips (3.9 to 5.9  $\mu\text{m}$ ). Asci were clavate (23 to 35  $\mu\text{m} \times 10$  to 15  $\mu\text{m}$ ) with eight limoniform, bilaterally flattened, brown ascospores (6.1 to 8.2  $\mu\text{m} \times 5.9$  to 7.8  $\mu\text{m}$  in size). Isolates were obtained from Taxila (CG1) and Wah-Cantt (CG1). The molecular identification was confirmed with the amplification of the ITS regions and GenBank accession numbers were obtained as MN891844 and MN891845. The frequency was 4.4%.

### Morpho-molecular identification of *Botrytis* fruit rot

Light to the dark gray colony was recorded in the single isolate of *Botrytis cinerea* obtained from Taxila (BC). The single-celled conidia were globose or ellipsoidal shaped, ranging from 5.92 to 11.15  $\mu\text{m} \times 5.74$  to 7.30  $\mu\text{m}$ . More or less straight conidiophores directly emerged from the mycelia ranging from 594 to 3,284  $\mu\text{m}$  in length. Conidiophores were septate, branched at the apex, and producing grape clusters shape conidia. The frequency of *B. cinerea* was 2.2%. The ITS regions (650bp) were amplified through PCR assay and the sequence was submitted with GenBank Accession number MN891765.

### Morpho-molecular identification and frequency *Pestalotiopsis* fruit rot

White fungal colony of sparse aerial mycelium with acervuli containing black, slimy spore masses on the surface of PDA was observed in the single isolate of *Pestalotiopsis mangiferae* obtained from Taxila (PM). The colony reached 8 cm diameter after 7-days of culture on PDA at 24°C. Conidia produced in the culture were five-celled, narrow fusiform, straight or slightly curved, with a tapering base and 2 to 4 hyaline appendages (apical appendages measured 15 to 34  $\mu\text{m}$  long and a single basal appendage were 5 to 9  $\mu\text{m}$  long). Conidia were 24 to 32  $\times$  5 to 8  $\mu\text{m}$  with median cells 15 to 20  $\mu\text{m}$  and two hyaline, cylindrical to conical apical cells. The ITS regions were confirmed with the amplification of 650bp in PCR assay and the GenBank accession number was MN888956. The frequency was recorded as 2.2%.


### Pathogenicity tests

Disease symptoms were recorded on artificially inoculated leaves, twigs and fruits, while no symptoms were found on healthy fruits and plants.

### Sequencing and sequence analysis

Initially, the ITS sequences from all fungal pathogens associated with loquat were used for the sequence analysis. The tree with the highest log-likelihood is presented in Fig 1A. The tree was drawn to scale, with branch lengths measured in the number of substitutions per site. The isolates exhibiting the maximum genetic similarity were found in a similar cluster. The combined sequence analysis of ITS and GAPDH of *C. lunata* isolates was conducted and all isolates were recorded in *C. lunata* cluster with reference isolates CIMAP SB-8213 (Fig 1B). The other reference isolates of *C. coatesiae* were recorded in *C. coatesiae* cluster. The combined sequence analysis of ITS and BT gene from all *A. alternata* isolates were recorded in *A. alternata* cluster with reference isolates GZU-BCEC154 due to maximum genetic homology between the isolates. The combined sequences of reference isolates of *A. panax* were recorded in *A. panax* cluster (Fig 1C).





**Fig 1. Evolutionary history of ITS region from all isolates (a) and multigene sequence analysis of *Curvularia* (b) and *Alternaria* (c) isolates.**

<https://doi.org/10.1371/journal.pone.0257951.g001>

### Discussion

Pakistan has the largest area under loquat production in southern Asia [11] and it is considered a major fruit crop of the country for local and export markets. Loquat fruit has a very short postharvest life and about 500 tones reduction in the loquat yield has been attributed to different biotic and abiotic factors [4]. An inappropriate post-harvest handling, poor storage conditions, unavailability of suitable transport vehicles, and inadequate marketing strategies are responsible for changing the physiological state of the fruit and seriously affecting its quality. Nevertheless, information on diversity of fungal pathogens associated with loquat is insufficient. The current study describes information regarding diversity of fungal pathogens associated with leaf spots, fruit rot, and twig dieback of loquat in Pakistan. The study was conducted to identify the isolated fungi based on morphological as well as molecular approaches.

Ascomycota and Deuteromycota are two phyla associated with loquat. The identified pathogens belonged to the families *Botryosphaeriaceae*, *Cerambycidae*, *Nectriaceae*, *Pleosporaceae*, *Sclerotiniaceae*, and *Venturiaceae*. The pathogens associated with *Pleosporaceae* cause leaf spots and fruit rot, while the remaining were associated with fruit rot. The occurrence of *A. alternata*, *C. lunata*, *L. theobromae*, *A. flavis*, *B. cinerea*, *C. globosum*, *P. mangiferae* and *Phomopsis* sp. Was confirmed in the major loquat growing areas of Pakistan. The *A. flavus*, *B. cinerea*, *P. mangiferae*, *C. globosum* and *Phomopsis* sp. were minor pathogenic species. The *C. lunata* and *A. alternata* were majorly associated with leaf spots and fruit rot and *L. theobromae* with twig dieback. This may indicate that variation in environmental conditions results in different fungal species [25].

Earlier, gray mold and black spot rot have been reported from loquat fruit [26]. *A. alternata* has been reported to cause decay of loquat in Palestina [27], China [28], Taiwan [26], Greece [29], and Iran [30]. This study also represents the first attempt to characterize the virulence of these fungal species. The frequency of associated pathogens was determined in each growing

location because they depend on local pre-harvest, harvest, and postharvest factors. The samples were collected based on the visual disease symptoms as they play a significant role in diagnosis [9]. The symptoms from a single pathogen were somewhat like other locations and a relationship between symptoms and the different fungal pathogens isolated was observed to a certain extent.

The symptoms associated with *Alternaria* leaf spots and fruit rot of loquat in Pakistan were similar to the symptoms recorded in different locations of Taiwan [26], Greece [29], Iran [30], and Pakistan [19]. The leaf spots associated with *C. lunata* were similar to the previously described symptoms of loquat in Pakistan [12]; however, this pathogen was isolated first time from the fruit. Circular to elongated sunken spots were recorded on the loquat fruit infected with *Pestalotiopsis* sp. and the symptoms were similar to the description of *Pestalotiopsis* fruit rot of loquat in China [31]. The occurrence of gray mold caused by *B. cinerea* was also recorded on loquat [32] and it is an aggressive airborne pathogen and can cause massive losses in the field conditions [33]. Circular to elongated, sunken spots were recorded over the surface of loquat fruit infected with *P. mangiferae* and similar symptoms were observed in Argentina [34]. *Aspergillus* sp. [35], *Chaetomium* sp. [36], and *L. theobromae* [11] were also reported from loquat.

Symptomology is not an appropriate criterion because symptom development might be due to several biotic as well as abiotic factors. To reduce this doubt, physiological and microscopic characters were further adapted for the identification of fungal pathogens associated with loquat. These characters are limited to describe species boundaries due to high level of cryptic speciation and extreme morphological homoplasy. The species identification in the genus is confusing and a high level of expertise is mandatory to distinguish these at species level based on morphological characters. To remove this confusion, new molecular approaches, i.e., sequence analysis of the universal ITS region was further proposed for the identification of fungal pathogens at species level [22]. *A. flavus* is common in crops producing most important mycotoxins [37]. This is the first evidence of *A. flavus*, *B. cinerea*, *P. mangiferae*, *C. globosum* and *Phomopsis* sp. decay in loquat growing areas of Pakistan, and morpho-molecular identification was used to confirm of these pathogens in Spain [32, 38].

Traditional diagnostic methods for detection and identification of fungal pathogens were based on specific media, temperature, and lighting for sporulation and morphological characterization. *Alternaria* and *Curvularia* isolates were difficult to differentiate based on morphological characteristics and sequence analysis of the ITS regions. The multi-gene phylogenetic analyses can positively separate them. The multiple sequence analysis of *Curvularia* sp. (Fig 1B) and *Alternaria* sp., (Fig 1C) confirmed that all isolates from a single species were recorded in a similar cluster. In pathogenicity tests, the rot was recorded on wounded and non-wounded fruits, while no symptoms were recorded on negative control. All the fungal pathogens isolated from fruit rot, leaf spots, and twig dieback of loquat were pathogenic with different virulence categories. The spreading of the diseases in loquat growing areas of Pakistan probably occurs by the transportation of contaminated loquat fruit and it could help to disperse the fungal pathogens to other locations. The dispersal might also occur through the movement of plant material and fruit from neighboring countries due to the inadequate quarantine department facility in the country.

## Conclusion

In conclusion, this study identified fungal species isolated from loquat trees with symptoms of leaf spots, fruit rot, and twig dieback. The present study depicted that the fruit rot of loquat can occur as a result of infection by *A. flavus*, *B. cinerea*, *C. globosum*, *P. mangiferae*, *Phomopsis*

sp., whereas *A. alternata*, *C. lunata* and *L. theobromae* was infecting twigs. The development of new loquat cultivars should consider the diversity of fungal species associated with the fruit rot. The current study provides a framework for phylogenetic analysis of fungal pathogens associated with fruit rot, leaf spots, and twig dieback of loquat. The morphological characteristics, multigene sequence analysis, and pathogenicity testing can assist the Pakistan loquat industry to implement control strategies to minimize economic losses and attention should be devoted to the treatment of loquat growing areas to reduce the incidence of fungal infections. The proper management will increase the production of loquat fruit and improve the economic condition of the growers.

## Author Contributions

**Conceptualization:** Sana Batool, Naseeb Ullah, Muhammad Rafiq, Saqer S. Alotaibi, Ahmed M. El-Shehawi, Yunzhou Li, Ali Tan Kee Zuan, Mohammad Javed Ansari.

**Data curation:** Sana Batool, Muhammad Rafiq.

**Formal analysis:** Muhammad Fahim Abbas, Sidra Mubeen, Naseeb Ullah.

**Funding acquisition:** Saqer S. Alotaibi, Ahmed M. El-Shehawi, Yunzhou Li, Ali Tan Kee Zuan.

**Investigation:** Sana Batool, Naseeb Ullah, Khalida Zafar.

**Methodology:** Muhammad Fahim Abbas, Khalida Zafar.

**Project administration:** Sidra Mubeen, Khalida Zafar, Muhammad Rafiq, Abdullah M. Al-Sadi.

**Resources:** Sidra Mubeen, Azziz-ud-Din.

**Software:** Sobia Khaliq, Azziz-ud-Din.

**Supervision:** Abdullah M. Al-Sadi.

**Validation:** Sobia Khaliq.

**Visualization:** Sobia Khaliq, Azziz-ud-Din.

**Writing – original draft:** Muhammad Fahim Abbas.

**Writing – review & editing:** Abdullah M. Al-Sadi, Saqer S. Alotaibi, Ahmed M. El-Shehawi, Yunzhou Li, Ali Tan Kee Zuan, Mohammad Javed Ansari.

## References

1. Hussain A, Abbasi N, Akhtar A. Fruit characteristics of different loquat genotypes cultivated in Pakistan; 2006. pp. 287–292.
2. Khan I. The history of loquat growing and future prospects of its commercial cultivation and marketing in Pakistan; 2003. pp. 25–26.
3. Hussain A, Abbasi NA, Hafiz IA, Ahmad Z, Naqvi SSJPB (2009) Molecular characterization and genetic relationship among loquat (*Eriobotrya japonica* Lindl.) genotypes of Pakistan assessed by RAPD markers. 41: 2437–2444.
4. GOP (2016) Fruits, vegetables and condiments statistics of Pakistan. Govt of Pak Mini. Food Agri. Livest: 1–10.
5. Cai C, Xu C, Shan L, Li X, Zhou C, et al. (2006) Low temperature conditioning reduces postharvest chilling injury in loquat fruit. *Post Biol Tech* 41: 252–259.
6. Cao S, Cai Y, Yang Z, Joyce DC, Zheng Y (2014) Effect of MeJA treatment on polyamine, energy status and anthracnose rot of loquat fruit. *Food Chem* 145: 86–89. <https://doi.org/10.1016/j.foodchem.2013.08.019> PMID: 24128452

7. Liu A, Chen W, Gu H, Shi J, Li J. Biological characteristic of pathogenic fungus causing anthracnose of loquat fruit; 2006. pp. 465–470.
8. Yahia EM (2011) Postharvest biology and technology of tropical and subtropical fruits: fundamental issues: Elsevier.
9. Batool M, Mukhtar T, Butt H (2014) First report of *Diplocarpon mespili* on loquat (*Eriobotrya japonica*) in Pakistan. *Österreichische Zeitschrift für Pilzkunde* 23: 143–147.
10. Hafeez R, Akhtar N, Shoaib A, Bashir U, Haider M, et al. (2015) First report of *Geotrichum candidum* from Pakistan causing postharvest sour rot in loquat (*Eriobotrya japonica*). *Anim Plant Sci* 25: 1737–1740.
11. Butt H, Mukhtar T, Batool M, Aslam MN, Kayani MZ, et al. (2015) Occurrence of leaf scab caused by *Spilocaea pyracanthae* on loquat (*Eriobotrya japonica*) in the Pothohar region of Pakistan. *Mycopath* 13: 29–32.
12. Abbas M, Naz F, Tariq A, Mumtaz A, Irshad G, et al. (2016) First report of *Curvularia lunata* causing leaf spots on loquat from Pakistan. *J Plant Pathol* 98.
13. Abbas MF, Naz F, Rauf CA, Khan MA (2016) Cultural, morphological, pathogenic and molecular characterization of *Alternaria mali* associated with necrotic leaf spot of loquat. *Int J Biosci* 9: 271–281.
14. Mehmood N, Riaz A, Jabeen N, Anwaar S, Hussain S, et al. (2017) First report of *Fusarium solani* causing fruit rot of strawberry in Pakistan. *Plant Dis* 101: 1681–1681.
15. Abbas M, Naz F, Rauf CA, Mehmood N, Zhang X, et al. (2017) First report of *Fusarium solani* causing fruit rot of loquat (*Eriobotrya japonica*) in Pakistan. *Plant Dis* 101: 839.
16. Naz F, Abbas M, Rauf C, Tariq A, Mumtaz A, et al. (2017) First report of *Colletotrichum gloeosporioides* causing anthracnose on loquat in Pakistan. *Plant Dis* 101: 1550–1550.
17. Abbas MF, Naz F (2018) First report of *Diplodia seriata* causing fruit rot of loquat in Pakistan. *J Plant Pathol* 100: 325–325.
18. Abbas MF, Naz F, Batool S, Naeem M, Qamar MI (2018) First evidence of *Mucor* rot infecting loquat (*Eriobotrya japonica* L.) in Pakistan. *Mycopath* 16.
19. Abbas MF, Naz F, Batool S, Qamar MI, Moosa A (2020) Morpho-molecular identification of *Alternaria alternata* causing leaf spot on loquat in Pakistan. *J Plant Pathol*: 1–1. <https://doi.org/10.1007/s42161-020-00644-w> PMID: 32904534
20. Leslie JF, Summerell BA (2008) *The Fusarium laboratory manual*: John Wiley & Sons.
21. Arif M, Atta S, Bashir MA, Khan MI, Hussain A, et al. (2021) The impact of Fosetyl-Aluminium application timing on Karnal bunt suppression and economic returns of bread wheat (*Triticum aestivum* L.). *Plos one* 16: e0244931. <https://doi.org/10.1371/journal.pone.0244931> PMID: 33428646
22. White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *CR protocols: a guide to methods applications* 18: 315–322.
23. Abbas MF, Rafiq M, Al-Sadi AM, Alfarraj S, Alharbi SA, et al. (2021) Molecular characterization of leaf spot caused by *Alternaria alternata* on buttonwood (*Conocarpus erectus* L.) and determination of pathogenicity by a novel disease rating scale. *Plos one* 16: e0251471. <https://doi.org/10.1371/journal.pone.0251471> PMID: 33984023
24. Tamura K, Stecher G, Peterson D, Filipksi A, Kumar S (2013) MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* 30: 2725–2729. <https://doi.org/10.1093/molbev/mst197> PMID: 24132122
25. Saikkonen K (2007) Forest structure and fungal endophytes. *fUNG Biol Rev* 21: 67–74.
26. Ko Y, Liu C, Chen S, Chen C, Yao K, et al. (2010) First report of fruit rot of loquat caused by an *Alternaria* sp. in Taiwan. *Plant Dis* 94: 481–481. <https://doi.org/10.1094/PDIS-94-4-0481B> PMID: 30754488
27. Batta Y (2005) Control of *Alternaria* spot disease on loquat (*Eriobotrya japonica* Lindl.) using detached fruits and leaf-disk assay. *An-Najah University Journal for Research Natural*.
28. Gu H, Liu A, Chen W, Feng S, Shi J. Development and control of postharvest diseases of loquat fruit; 2006. pp. 437–444.
29. Tziros GT (2013) *Alternaria alternata* causes leaf spot and fruit rot on loquat (*Eriobotrya japonica*) in Greece. *Aust Plant Dis Not* 8: 123–124.
30. Mirhosseini HA, Babaeizad V, Basavand E (2015) Identification and detection of agent of loquat leaf spot and fruit rot in north of Iran. *J New Biol Repor* 4: 135–138.
31. Chen Y, Wang W-X, Zhang A-F, Yang X, Xu Y-L (2013) First report of *Pestalotiopsis theae* on Loquat (*Eriobotrya japonica*) in Anhui province of China. *Plant Dis* 97: 558–558. <https://doi.org/10.1094/PDIS-08-12-0710-PDN> PMID: 30722250

32. Palou L, Sánchez-Torres P, Montesinos-Herrero C, Taberner V (2016) Incidence and etiology of post-harvest fungal diseases of loquat fruit (*Eriobotrya japonica* (Thunb.) Lindl. cv. 'Algerie') in Alacant province (Spain). *Eur J Plant Pathol* 146: 847–860.
33. Williamson B, Tudzynski B, Tudzynski P, van Kan JA (2007) *Botrytis cinerea*: the cause of grey mould disease. *Mol Plant Pathol* 8: 561–580. <https://doi.org/10.1111/j.1364-3703.2007.00417.x> PMID: 20507522
34. Perelló A, Larran S (1999) First report of *Pestalotiopsis guepini* on loquat in Argentina. *Plant Dis* 83: 695–695. <https://doi.org/10.1094/PDIS.1999.83.7.695A> PMID: 30845627
35. Rana O, Upadhyaya J (1972) A new fruit rot of loquat (*Eriobotrya japonica* L.). *Sci Cult* 38: 35–36.
36. Wang X, Lombard L, Groenewald J, Li J, Videira S, et al. (2016) Phylogenetic reassessment of the *Chaetomium globosum* species complex. *Mol Phyl Evol Fungi* 36: 83. <https://doi.org/10.3767/003158516X689657> PMID: 27616789
37. Frisvad JC, Hubka V, Ezekiel C, Hong S-B, Nováková A, et al. (2019) Taxonomy of *Aspergillus* section *Flavi* and their production of aflatoxins, ochratoxins and other mycotoxins. *Stud in Mycol* 93: 1–63.
38. González-Domínguez E, León M, Armengol J, Berbegal M (2015) A nested polymerase chain reaction protocol for in planta detection of *Fusicladium eriobotryae*, causal agent of loquat scab. *J Phytopath* 163: 415–418.
39. Jeon SJ, Nguyen TTT, Lee HB (2015) Phylogenetic status of an unrecorded species of *Curvularia*, *C. spicifera*, based on current classification system of *Curvularia* and *Bipolaris* group using multi loci. *Mycologia* 43: 210–217.