


RESEARCH ARTICLE

Poor neutralization and rapid decay of antibodies to SARS-CoV-2 variants in vaccinated dialysis patients

Jessica Bassi¹ , Olivier Giannini^{2,3} , Chiara Silacci-Fregni¹, Laura Pertusini⁴, Paolo Hitz², Tatiana Terrot⁵, Yves Franzosi⁵, Francesco Muoio¹ , Christian Saliba¹, Marcel Meury⁶, Exequiel A. Dellota, Jr.⁶, Josh R. Dillen⁶, Patrick Hernandez^{6*} , Nadine Czudnochowski⁶, Elisabetta Cameroni¹, Nicola Beria⁴, Mariangela Ventresca⁴, Alberto Badellino⁴, Soraya Lavorato-Hadjeres⁴, Elisabetta Lecchi⁴, Tecla Bonora⁴, Matteo Mattiolo⁴, Guido Trinci⁴, Daniela Garzoni⁴, Giuseppe Bonforte⁴, Valentina Forni-Ogna^{4*} , Davide Giunzioni⁴, Lorenzo Berwert⁴, Ravindra K. Gupta^{7,8}, Paolo Ferrari^{2,4,9} , Alessandro Ceschi^{2,5,10,11}, Pietro Cippà^{3,4,12}, Davide Corti¹, Antonio Lanzavecchia¹, Luca Piccoli^{1*} 

1 Humabs BioMed SA, A Subsidiary of Vir Biotechnology, Bellinzona, Switzerland, **2** Faculty of Biomedical Sciences, Università della Svizzera italiana, Lugano, Switzerland, **3** Department of Medicine, Ente Ospedaliero Cantonale, Bellinzona, Switzerland, **4** Division of Nephrology, Ente Ospedaliero Cantonale, Lugano, Switzerland, **5** Clinical Trial Unit, Ente Ospedaliero Cantonale, Lugano, Switzerland, **6** Vir Biotechnology, San Francisco, California, United States of America, **7** Cambridge Institute of Therapeutic Immunology & Infectious Disease (CITIID), Cambridge, United Kingdom, **8** Department of Medicine, University of Cambridge, Cambridge, United Kingdom, **9** Clinical School, University of New South Wales, Sydney, Australia, **10** Division of Clinical Pharmacology and Toxicology, Institute of Pharmacological Science of Southern Switzerland, Ente Ospedaliero Cantonale, Lugano, Switzerland, **11** Department of Clinical Pharmacology and Toxicology, University Hospital Zurich, Zurich, Switzerland, **12** Faculty of Medicine, University of Zurich, Zurich, Switzerland

 These authors contributed equally to this work.

* patrick.hernandez1@gmail.com (PH); valentina.forniozna@eoc.ch (VFO); lpiccoli@vir.bio (LP)



OPEN ACCESS

Citation: Bassi J, Giannini O, Silacci-Fregni C, Pertusini L, Hitz P, Terrot T, et al. (2022) Poor neutralization and rapid decay of antibodies to SARS-CoV-2 variants in vaccinated dialysis patients. *PLoS ONE* 17(2): e0263328. <https://doi.org/10.1371/journal.pone.0263328>

Editor: Etsuro Ito, Waseda University: Waseda Daigaku, JAPAN

Received: October 21, 2021

Accepted: January 14, 2022

Published: February 10, 2022

Copyright: © 2022 Bassi et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The minimal data set underlying the results described in our paper is provided as [Supplementary Information](#) (source data of each figure). Due to health and data protection laws relating to the demographic and clinical information contained in the paper, we will not be able to fully comply with the requirement to share demographic and clinical data of patients in this study.

Funding: The project was partially funded by the Swiss Kidney Foundation (to O.G.). R.K.G. received

Abstract

Patients on dialysis are at risk of severe course of SARS-CoV-2 infection. Understanding the neutralizing activity and coverage of SARS-CoV-2 variants of vaccine-elicited antibodies is required to guide prophylactic and therapeutic COVID-19 interventions in this frail population. By analyzing plasma samples from 130 hemodialysis and 13 peritoneal dialysis patients after two doses of BNT162b2 or mRNA-1273 vaccines, we found that 35% of the patients had low-level or undetectable IgG antibodies to SARS-CoV-2 Spike (S). Neutralizing antibodies against the vaccine-matched SARS-CoV-2 and Delta variant were low or undetectable in 49% and 77% of patients, respectively, and were further reduced against other emerging variants. The fraction of non-responding patients was higher in SARS-CoV-2-naïve hemodialysis patients immunized with BNT162b2 (66%) than those immunized with mRNA-1273 (23%). The reduced neutralizing activity correlated with low antibody avidity. Patients followed up to 7 months after vaccination showed a rapid decay of the antibody response with an average 21- and 10-fold reduction of neutralizing antibodies to vaccine-matched SARS-CoV-2 and Delta variant, which increased the fraction of non-responders to 84% and 90%, respectively. These data indicate that dialysis patients should be prioritized

support from the G2P-UK National Virology consortium funded by MRC/UKRI (grant ref. MR/W005611/1).

Competing interests: I have read the journal's policy and the authors of this manuscript have the following competing interests: J.B., C.S.-F., F.M., C.S., M.Me., E.A.D.J., N.C., E.C., D.C. A.L., and L.Pi. are employees of Vir Biotechnology Inc. and may hold shares in Vir Biotechnology Inc. R.K.G. has received consulting fees from Johnson and Johnson and GlaxoSmithKline for educational activities. The other authors declare no competing interests. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

for additional vaccination boosts. Nevertheless, their antibody response to SARS-CoV-2 must be continuously monitored to adopt the best prophylactic and therapeutic strategy.

Introduction

Patients with chronic kidney disease (CKD), in particular those with end-stage kidney disease (ESKD) on dialysis, are highly predisposed to infections, which are the major cause of morbidity and the second cause of mortality in this vulnerable population [1, 2]. Infection by SARS-CoV-2, the causative agent of the Coronavirus disease-2019 (COVID-19), has posed a new threat for CKD patients, who were found to have a greater risk of severe COVID-19 course [3, 4]. Early reports indicated a case fatality ranging from 10 to 30% in patients on hemodialysis (HD) [5–10].

The rapid development of COVID-19 vaccines has provided an important strategy to prevent SARS-CoV-2 infection and, especially, severe COVID-19 course. In particular, mRNA-based vaccines developed by Pfizer/BioNTech (BNT-162b2) and Moderna (mRNA-1273) have demonstrated high safety and efficacy in healthy and at-risk individuals, including patients with chronic diseases, cancer and solid organ transplantation [11–14]. However, immunosuppressed patients, in particular those with hematological malignancies, autoimmune diseases and solid organ transplantations, were shown to mount a low antibody response to these vaccines [15–20]. Because of their immunological frailty, patients on dialysis were prioritized in international COVID-19 vaccination programs [21]. Recent studies in HD patients showed a delayed and lower serological response to vaccines and a rapid decline of anti-SARS-CoV-2 antibodies [19, 22–24]. These findings suggest an overall diminished vaccine response to SARS-CoV-2 in ESKD patients that is reminiscent of the low response observed after vaccination against Hepatitis B virus (HBV) and seasonal Influenza virus [25, 26]. In addition, the rapid increase in cases of infection by SARS-CoV-2 Delta variant (B.1.617.2 lineage) since April 2021 provided a new potential challenge for dialysis patients, especially those with a sub-optimal response to the vaccine [27].

At this stage of the pandemic, it is urgent to identify populations not developing sufficient levels of neutralizing antibodies against circulating SARS-CoV-2 variants, which therefore may be at-risk of developing severe COVID-19. In this study, we provide evidence of a poor neutralizing and rapidly decaying antibody response to mRNA-vaccines in the dialysis population, which supports the strategy of prioritizing these patients for an additional boost and other therapeutic strategies.

Material and methods

Study participants and ethics statement

Blood samples were obtained from 143 dialysis patients and 48 healthcare workers under study protocols approved by the local Institutional Review Board (Canton Ticino Ethics Committee, Switzerland). All the participants were recruited from the four public hospitals of the Ente Ospedaliero Cantonale (EOC) in Ticino (Southern Switzerland). All subjects provided written informed consent for the use of blood and blood components (such as PBMCs, sera or plasma).

Isolation of plasma

Blood samples were collected from all the participants before, 2–3 weeks and up to 7 months after the second dose. An additional sample was collected 2–3 weeks after the first COVID-19

vaccine dose from 26 patients and all the 48 healthcare workers. Plasma was isolated from blood draw performed using BD tubes containing Ficoll (BD, CPT Ficoll, Cat. No. 362780) and stored at +4°C until use.

Cell lines

Cell lines used in this study were obtained from ATCC (Vero E6 TMPRSS2) or ThermoFisher Scientific (Expi CHO cells, Expi293F™ and HEK293F cells). Vero E6 TMPRSS2 cells were grown in DMEM supplemented with 10% HyClone (FBS). Expi293F cells were grown in Expi293 Expression Medium.

Production of recombinant glycoproteins

The SARS-CoV-2 RBD WT construct was synthesized by GenScript into phCMV1, with a sequence encoding an N-terminal mu-phosphatase signal peptide, an ‘ETGT’ linker, SARS-CoV-2 S residues 328–531, a linker sequence, an Avi tag, a twin Strep tag and a 8xHis-tag. Recombinant ACE2 (UniProt Q9BYF1, residues 19–615 with a C-terminal thrombin cleavage site-TwinStrep-10xHis-GGG-tag, and N-terminal signal peptide) and RBD WT constructs were transiently transfected into Expi293F cells following manufacturer’s instructions. Supernatants were clarified by centrifugation and affinity purified using StrepTrap HP columns (Cytiva). Monomeric ACE2 was isolated by size exclusion chromatography using a Superdex 200 Increase 10/300 GL column (Cytiva). The SARS-CoV-2 stabilized Spike WT (D614G) construct was synthesized by GenScript into pCDNA3.1 with an N-terminal mu-phosphatase signal peptide, 2P stabilizing mutation [28, 29], a TEV cleavage site and a C-terminal foldon, 8x His-tag, Avi tag and C-tag [30] and expressed in HEK293F 293 cells following manufacturer’s instructions. Supernatants were clarified by centrifugation and affinity purified using a 5 mL C-tag affinity matrix column.

VSV spike mutants and pseudovirus generation

Amino acid substitutions were introduced into the D614G pCDNA_SARS-CoV-2_S plasmid as previously described [31]. SARS-CoV-2 S glycoprotein-encoding-plasmids used to produce SARS-CoV-2 VSV, namely the Wuhan prototype (D614, referred as WT) and variants Alpha (B.1.1.7), Beta (B.1.351), Gamma (P.1), Delta (B.1.617.2), Epsilon (B.1.429), Kappa (B.1.617.1) and Lambda (C.37), were obtained using a multistep based on overlap extension PCR (oePCR) protocol [32].

Replication defective VSV pseudovirus [33] expressing SARS-CoV-2 Spike proteins corresponding to the different VOC were generated as previously described [34].

Plasma pseudovirus neutralization assay

Vero E6-TMPRSS2 were grown in DMEM supplemented with 10% FBS and seeded into white bottom 96 well plates (PerkinElmer, 6005688), as previously described [35]. Conditions were tested in duplicate wells in each plate and at least six wells per plate contained untreated infected cells (defining the 0% of neutralization, “MAX RLU” value) and infected cells in the presence of S2E12 and S2X259 mAbs at 50 µg/mL each (defining the 100% of neutralization, “MIN RLU” value). Average of Relative light units (RLUs) of untreated infected wells (MAX RLU_{ave}) was subtracted by the average of MIN RLU (MIN RLU_{ave}) and used to normalize percentage of neutralization of individual RLU values of experimental data according to the following formula: $(1 - (RLU_x - MIN\ RLU_{ave}) / (MAX\ RLU_{ave} - MIN\ RLU_{ave})) \times 100$. Data were analyzed and visualized with Prism (Version 9.1.0). Each neutralization experiment included a

technical duplicate. The loss or gain of neutralization potency across Spike variants was calculated by dividing the variant ID₅₀ by the parental ID₅₀.

Enzyme-linked immunosorbent assay (ELISA)

Spectraplate-384 with high protein binding treatment (custom made from Perkin Elmer) were coated overnight at 4°C with 5 µg/mL RBD or 1 µg/mL SARS-CoV-2 S protein in PBS. The day after plates were washed and blocked with Blocker Casein in PBS (Thermo Fisher Scientific, 37528) supplemented with 0.05% Tween 20 (Sigma Aldrich), 1h RT. Serial dilutions of plasma samples were then added to plates for 1h RT. Alkaline Phosphatase-conjugated goat anti-human IgG, IgM or IgA (Southern Biotech) were added to plates and incubated for 1h RT. 4-NitroPhenyl Phosphate (pNPP, Sigma-Aldrich, N2765-100TAB) substrate was then added and plates were read after 1 h (IgG) or 2 h (IgA and IgM) at 405 nm with a BioTek plate reader. Data were plotted and analyzed with GraphPad Prism software (version 9.1.0). For chaotropic ELISA, after incubation with plasma, plates were washed and incubated with a 1 M solution of sodium thiocyanate (NaSCN, Sigma 251410) for 1h RT. Avidity Index was calculated as the ratio (%) of the ED₅₀ in presence and the ED₅₀ in absence of NaSCN.

Blockade of RBD binding to human ACE2

Plasma were diluted in PBS and mixed with SARS-CoV-2 RBD mouse Fc-tagged antigen (Sino Biological, 40592-V05H, final concentration 20 ng/mL) and incubated for 30 min at 37°C. The percentage of inhibition was calculated as follows: $(1 - (\text{OD sample} - \text{OD neg ctr}) / (\text{OD pos ctr} - \text{OD neg ctr})) \times 100$.

Statistical analysis

The study was designed to have 80% power to detect a minimum 25% difference in total incidence of cases with poor neutralizing antibody response (i.e., low or undetectable plasma antibody titers) or in average neutralizing titers between dialysis patients and healthy controls as well as within the HD subgroups. Comparisons of means between two groups of unpaired data were made with Mann-Whitney rank test. Comparisons of means between multiple groups of unpaired data were made with Kruskal-Wallis rank test and corrected with Dunn's test. Comparisons of means between multiple groups of matched data were made with Friedman rank test and corrected with Dunn's test. Relative risks of poor neutralizing response in selected groups of patients were calculated from 2x2 contingency tables using two-sided Fisher's exact test. Statistical significance is set as $P < 0.05$ and P-values are indicated with: ns = non-significant; * = 0.033; ** = 0.002; *** < 0.001. ED₅₀ and ID₅₀ titers were calculated from the interpolated value from the log(agonist) and the log(inhibitor), respectively, versus response, using variable slope (four parameters) non-linear regression. Data were plotted and analyzed with GraphPad Prism software (version 9.1.0).

Results

Naïve dialysis patients produce low or undetectable levels of antibodies after two vaccine doses

At the beginning of the COVID-19 vaccination campaign in January 2021, 143 dialysis patients (130 HD, of whom 68% on hemodiafiltration, and 13 on peritoneal dialysis, PD) were enrolled in this study from four dialysis units of the Ente Ospedaliero Cantonale in Ticino, Switzerland, and received two doses of SARS-CoV-2 mRNA vaccines (83% with BNT162b2 from Pfizer-BioNTech and 17% with mRNA-1273 from Moderna) between January and May 2021. Socio-

demographic data, dialysis features, comorbidities, therapies and information about previous vaccinations and infections are summarized in [Table 1](#). Of note, twenty-four (17%) dialysis patients were previously infected with SARS-CoV-2, of whom 18 (75%) had severe COVID-19 requiring hospitalization (13) or admission to an intensive care unit (5). A group of 48 health-care workers, who received two doses of BNT162b2 vaccine between April and June 2021, were included as healthy controls (HC). Twenty-four HC were previously infected with SARS-CoV-2 reporting mild symptoms that did not require hospitalization.

Plasma samples were collected 2 to 3 weeks after the first and the second dose and the levels of plasma IgG, IgA and IgM specific for SARS-CoV-2 Spike (S) were measured by ELISA. A single vaccine dose induced detectable IgG in 34.6% of dialysis patients compared to 89.6% of HC, with high antibody levels in all HD and HC participants, who had been previously infected with SARS-CoV-2, and in 29.2% of HD and 79.2% of HC, who had not been exposed to the virus ([S1 Fig](#)). Importantly, although seroconversion was observed in 94.4% of dialysis patients after the second vaccine dose, 35% of them had still low or undetectable levels of anti-SARS-CoV-2 S antibodies, compared to 100% of HC showing high levels of S-specific IgG ([Fig 1A and 1B](#)).

The larger fraction of participants with low or no antibodies (36.9%) was represented by the HD group, with naïve patients immunized with BNT162b2 showing the lowest levels of plasma IgG (50.6%) compared to those who received mRNA-1273 (9.1%) ([Fig 1A and 1B](#)). The lower IgG level in naïve HD patients was the result of a slower kinetics of antibodies induced by the vaccines as compared to naïve HC and previously infected participants ([Fig 1C](#)). Of note, some previously infected HD patients produced higher amounts of antibodies compared to HC ([Fig 1A](#)), a finding that is consistent with a more severe COVID-19 course in these patients [[36–38](#)]. Plasma samples from the 13 PD patients were collected only after the second vaccine dose and all the participants showed detectable S-specific IgG with average plasma levels that were comparable to those of HC ([Fig 1A and 1B](#)). S-specific IgG levels highly correlated with RBD-specific IgG, suggesting that, similarly to natural infection, antibodies to this domain dominated the response induced by vaccination in both patients and controls ([Fig 1D and S2 Fig](#), panel a). Finally, we observed that two vaccine doses induced, in the majority of the participants, undetectable levels of IgM and detectable levels of IgA specific for SARS-CoV-2 S, which were higher in previously infected HC and HD participants vaccinated with BNT162b2 and in naïve HD patients vaccinated with mRNA-1273 compared to naïve HD vaccinated with BNT162b2, and correlated with the corresponding IgG levels ([Fig 1E–1G and S2 Fig](#), panels b and c).

Poor neutralization of Wuhan prototypic SARS-CoV-2 and related risk factors in naïve dialysis patients

To determine the neutralizing activity of vaccine-induced antibodies, we used plasma samples to perform an *in vitro* neutralization assay with pseudotyped vesicular stomatitis virus (VSV) that expresses the Wuhan wild-type (WT, D614) SARS-CoV-2 S glycoprotein ([Fig 2A and S3 Fig](#), panel a). Compared to HC, 49% of dialysis patients had no or low neutralizing activity against the vaccine-matched SARS-CoV-2 strain ([Fig 2B](#)). In particular, while most of previously infected HD patients (87%) had moderate to high neutralizing antibody titers, which were similar or higher than those of HC, naïve HD patients had a heterogeneous response with 57% of them characterized by a poor neutralizing activity, which was completely absent in the majority (60.7%) of these patients ([Fig 2A](#)). Of note, the fraction of non-responding patients was higher in naïve HD patients immunized with BNT162b2 (65.9%) than those immunized with mRNA-1273 (22.7%) ([Fig 2B](#)). Despite higher plasma S-specific IgG titers ([Fig 1A and](#)

Table 1. Socio-demographic and clinical data of enrolled dialysis patients and healthy controls.

	Dialysis patients	Hemodialysis (HD)	Peritoneal dialysis (PD)	Healthy controls (HC)
Enrolled participants	143	130	13	48
Age, median years (min-max, IQR)	76 (29–97, 67–83)	76 (29–97, 68–83)	71 (32–82, 55–78)	42 (24–67, 32–49)
Gender, female, n (%)	48 (34%)	43 (33%)	5 (38%)	35 (73%)
BMI, median Kg/m ² (min-max, IQR)	25.9 (17.6–58.8, 22.9–29.2)	26.0 (17.6–58.8, 22.8–29.4)	25.7 (18–36.8, 23.3–29.0)	25.0 (17–3, 21–28)
Smoker, n (%)	21 (15%)	18 (14%)	3 (23%)	13 (27%)
Dialysis treatment, n (%)				
HD, n (%)	130 (91%)	130 (100%)		
HDF, n (%)		89 (68%)		
PD, n (%)	13 (9%)		13 (100%)	
Dialysis vintage, median months (min-max, IQR)	42 (11–268, 25–80)	46 (11–268, 26–87)	28 (11–70, 16–44)	
Dialysis access, native AVF, n (%)	96 (67%)	96 (74%)		
Comorbidities, n (%)				
Diabetes	58 (41%)	56 (43%)	2 (15%)	2 (4%)
Hypertension	121 (85%)	109 (84%)	12 (92%)	5 (10%)
Heart failure	19 (13%)	18 (14%)	1 (8%)	-
Coronary heart disease	46 (32%)	41 (32%)	5 (38%)	-
Peripheral arterial disease	29 (20%)	28 (22%)	1 (8%)	-
Pulmonary disease	42 (29%)	38 (29%)	4 (31%)	-
Chronic liver disease	12 (8%)	12 (9%)	-	-
Gastrointestinal disease	27 (19%)	24 (18%)	3 (23%)	-
Hemato-oncological disease	12 (8%)	12 (9%)	1 (8%)	-
Autoimmune disease	11 (8%)	9 (7%)	2 (15%)	1 (3%)
Kidney transplant	10 (7%)	10 (8%)	-	-
Neurodegenerative disease	4 (3%)	4(3%)	-	-
Psychiatric disease	13 (9%)	13 (10%)	-	-
Charlson comorbidity index, median (min-max, IQR)	8 (3–21, 7–11)	9 (3–21, 7–11)	6 (5–11, 6–8)	-
Previous SARS-CoV-2 infection ¹ , n (%)	24 (17%)	23 (18%)	1 (8%)	24 (50%)
mild or no symptoms	6 (25%)	5 (22%)	1 (100%)	24 (50%)
severe course with hospitalization	18 (75%)	18 (78%)	-	-
ICU admission	5 (28%)	5 (28%)	-	-
Diagnosis-vaccine interval, median months (min-max, IQR)	3.5 (1.1–9.7, 2.9–7.7)	3.3 (1.1–9.7, 2.9–7.7)	9.3	4.8 (3.5–12.8, 4.7–5.5)
COVID-19 vaccine received				
Pfizer-BioNTech, n (%)	118 (83%)	107 (82%)	11 (85%)	48 (100%)
Moderna, n (%)	25 (17%)	23 (18%)	2 (15%)	-
Dose 1—dose 2 interval, median days (min-max, IQR)	28 (20–38, 26–28)	28 (20–28, 26–28)	24 (21–28, 22–28)	38 (23–31, 28–28)
Sampling after dose 1, median days (min-max, IQR)	15 (9–33, 13–26)	15 (9–33, 13–26)	-	13 (12–16, 12–14)
Sampling after dose 2, median days (min-max, IQR)	18 (13–27, 17–19)	18 (13–27, 17–19)	18 (16–25, 17–21)	13 (10–21, 12–14)
2020 seasonal flu vaccination, n (%)	101 (70%)	94 (72%)	7 (54%)	29 (60%)
Pharmacological treatment				
Oral anticoagulation	27 (19%)	25 (19%)	2 (15%)	-
Antiaggregant drugs	124 (43%)	124 (95%)	-	-
ACE inhibitors/ARBs	70 (49%)	58 (45%)	12 (92%)	3 (6%)
Calcium antagonists	61 (43%)	51 (39%)	10 (77%)	-
Beta blockers	88 (62%)	79 (61%)	9 (69%)	2 (4%)
Statins	80 (56%)	71 (55%)	9 (69%)	2 (4%)
Oral antidiabetic drugs	22 (15%)	22 (17%)	-	2 (4%)

(Continued)

Table 1. (Continued)

	Dialysis patients	Hemodialysis (HD)	Peritoneal dialysis (PD)	Healthy controls (HC)
Insulin	37 (26%)	35 (27%)	2 (15%)	-
Immunosuppressive drugs ²	24 (17%)	22 (17%)	2 (15%)	-
Oral/intravenous steroids	17 (12%)	16 (12%)	1 (8%)	-
Calcineurin inhibitors ³	8 (6%)	8 (6%)	-	-
Vitamin D derivatives	112 (78%)	102 (78%)	10 (77%)	-

IQR, interquartile range. BMI, body mass index. HDF, hemodiafiltration. AVF, arteriovenous fistula. PCR, polymerase chain reaction. ICU, intensive care unit. ACE, angiotensin-converting enzyme. ARB, angiotensin receptor blockers.

¹with positive PCR and/or serology.

²include any of steroids, cyclosporine, tacrolimus, rituximab, mycophenolate mofetil and azathioprine.

³include cyclosporine and tacrolimus.

<https://doi.org/10.1371/journal.pone.0263328.t001>

1B), 60% of naïve PD patients had a similar poor neutralizing response compared to naïve HD patients, with 33.3% of those, who received BNT162b2 vaccine, showing no neutralizing activity (Fig 2A and 2B). The poor neutralizing activity was confirmed also by the low capability of plasma antibodies to inhibit binding of RBD to human ACE2 receptor (S3 Fig, panel b). In some previously infected dialysis patients, we observed neutralizing and ACE2-inhibiting antibody titers that were higher than those of HC and correlated with S- and RBD-specific IgG levels (Fig 2A and S3 Fig, panels c-e).

We next performed a sub-analysis to understand which socio-demographic and clinical data were associated to a significant risk for dialysis patients of being poor responders, here defined as having low or no neutralizing antibody titers after two mRNA-vaccine doses. Among all patients, dialysis mode (HD or PD) was not identified as risk factor, whereas vaccination with BNT162b2 and no previous SARS-CoV-2 infection represented the major risk factors for having a poor neutralizing antibody response (relative risk of 2.75 and 4.50, 95% confidence interval [CI] of 1.37 to 6.30 and 1.78 to 13.1, respectively) (Table 2). Within the larger naïve HD group, socio-demographic factors, including gender, age, body mass index, smoke and dialysis features could not be identified as risk factors (Table 3). Among comorbidities, we found that patients with heart failure history had an 1.68-increased relative risk of poor response (95% CI, 1.16 to 2.17). As expected, patients in therapy with immunosuppressive drugs had a 1.53 relative risk of poor response to the vaccine (95% CI, 1.05 to 2.02) with, in particular, calcineurin inhibitors accounting for the highest relative risk (1.87; 95% CI, 1.18 to 2.29). Vaccination with BNT162b2 was confirmed as a major factor with a relative risk of poor response of 2.88 (95% CI, 1.47 to 6.55), which increased to 3.60 (95% CI, 1.79 to 8.23) in 22 naïve HD patients who were matched to 22 naïve HD patients immunized with mRNA-1273 by age, gender, dialysis vintage and comorbidities. Among HD patients immunized with BNT162b2, those older than 80 and with heart failure history had an increased risk of poor response of 1.57 (95% CI, 1.18 to 2.08) and 1.55 (95% CI, 1.11 to 1.96), respectively (Table 3).

Loss of neutralization of Delta and other SARS-CoV-2 variants in dialysis patients

We next addressed the question whether vaccine-induced antibodies that neutralize wild-type SARS-CoV-2 could also neutralize current circulating SARS-CoV-2 variants, in particular the Delta variant (B.1.617.2 lineage) (Fig 2C). Most dialysis patients (76.9%) had plasma antibodies with low or undetectable neutralizing activity against the Delta variant, with 48.0% and 80.0% of naïve HD and PD patients with detectable levels of neutralizing antibodies against




Fig 1. Comparison of mRNA vaccine-induced plasma antibody titers against SARS-CoV-2 between healthy controls and dialysis patients. a) Plasma IgG titers (ED50) to SARS-CoV-2 S after two doses of Pfizer/BioNTech (P/B) or Moderna (M) vaccines in previously infected (square) and naïve (circle) healthy controls (HC, red), hemodialysis (HD, blue) and peritoneal dialysis (PD, orange) patients. Grey areas indicate non-specific IgG titers <50, a cut-off that was determined on non-specific binding to uncoated ELISA plates. An additional cut-off of 1'000, determined from the lowest titers in HC after two doses, was used to distinguish low (50–1'000) from high (>1'000) IgG titers. Statistical significance is set as $P < 0.05$ and P-values are indicated with asterisks (* = 0.033; ** = 0.002; *** < 0.001). b) Percentages of participants with high, low or no plasma SARS-CoV-2 S-specific IgG after two doses of mRNA-vaccine. Total number of participants within each cohort of HC and dialysis patients (DP) and within each subgroup is shown at the top of each bar. c) Kinetics of plasma IgG titers to SARS-CoV-2 S measured before vaccination (T0), after one (T1) or two (T2) vaccine doses. Each line connects samples from the same individual. White squares indicate HD patients who were not yet infected at T0 sampling. d) Correlation analysis between plasma IgG titers to SARS-CoV-2 S and RBD in all the plasma samples collected after the second vaccine dose. Spearman $r = 0.96$ ($p < 0.001$). e) Plasma IgM titers to SARS-CoV-2 S after two vaccine doses in HC, HD and PD patients. Grey areas indicate non-specific IgM titers <50. f) Plasma IgA titers to SARS-CoV-2 S after two vaccine doses in HC, HD and PD patients. Grey areas indicate non-specific IgA titers <50. g) Correlation analysis between plasma IgG and IgA titers to SARS-CoV-2 S in all the plasma samples collected after the second vaccine dose. Spearman $r = 0.68$ ($p < 0.001$).

<https://doi.org/10.1371/journal.pone.0263328.g001>

(P.1), Epsilon (B.1.429) and Kappa (B.1.617.1) SARS-CoV-2 variants in 28 HC, 47 HD and 5 PD patients. g) Fold change analysis of neutralizing titers against different variants. Numbers in each cell of panel g indicate the average fold change values of WT ID50 to variant ID50 in all the cohorts analyzed including only participants with ID50 neutralizing titers against WT SARS-CoV-2 greater than 80. Shown are data from $n = 2$ independent experiments.

<https://doi.org/10.1371/journal.pone.0263328.g002>

WT SARS-CoV-2 showing complete loss of neutralization against this variant (Fig 2D and S4 Fig, panel a). Almost all the participants, including HC, had a greater than 2-fold reduction in the neutralizing activity against Delta variant, with naïve HD patients showing up to 17-fold (average 4.7) loss in neutralization, as compared to previously infected patients that showed a maximum 5-fold loss (average 3.0) (Fig 2E). Similarly, naïve PD patients showed up to 10-fold (average 5.4) loss of neutralization against Delta variant (Fig 2E).

The loss of neutralization against some of the other circulating SARS-CoV-2 variants was higher compared to Delta, in particular for naïve HD patients, who showed an average 7.3-, 7.6- and 6.8-fold reduction in their neutralizing activity against Beta (B.1.351), Epsilon (B.1.429) and Kappa (B.1.617.1) variants (Fig 2F and 2G and S4 Fig, panels b-d). Neutralization of Alpha (B.1.1.7), Gamma (P.1) and Lambda (C.37) was also reduced in HD patients, but with minor loss compared to Delta (1.8-, 3.3- and 3.1-fold reduction, respectively) (Fig 2F and 2G and S4 Fig, panels e-h). Previously infected HD patients and naïve PD patients also showed a drastic loss of neutralization against different variants, which was comparable to that of naïve HD patients, but higher than that of HC (Fig 2F and 2G and S4 Fig). Complete loss of neutralization of Beta, Epsilon and Kappa variants was observed in 37–43.3% of HD and 75% of PD patients with detectable neutralizing antibodies against WT SARS-CoV-2 (S4 Fig, panel a).

The reduced neutralizing activity correlates with a low antibody avidity

To further characterize the poor neutralizing activity of vaccine-induced antibodies, we determined the avidity of plasma antibodies by measuring their binding to SARS-CoV-2 S in presence of sodium thiocyanate, a chaotropic agent that induces dissociation of the antibody from the antigen in case of low affinity. While most of the previously infected HD patients (87%) showed moderate to high-avidity antibodies, a high fraction of naïve HD showed low- (42%) or no (9%) avidity antibodies and most of the naïve PD patients (75%) also showed low-avidity antibodies (Fig 3A). As expected, antibodies of all HC that were previously infected with SARS-CoV-2 showed high avidity (index >50%), whereas naïve HC showed moderate avidity (23–50%), consistent with the different duration of affinity maturation ongoing in the two groups (Fig 3A). The avidity indexes correlated with SARS-CoV-2 S-specific antibody and neutralizing titers (Fig 3B and S5 Fig, panel a). In particular, we observed larger fractions of HD patients with moderate to high avidity among patients with higher neutralizing antibody titers (Fig 3C). The lower avidity titers were paralleled by higher fractions of dialysis patients

Table 2. Analysis of relative risk of poor neutralizing response to COVID-19 vaccination in dialysis patients.

Variable	Analyzed group	Reference group	n_{an}	n_{ref}	Relative Risk	95% CI	P-value	Significant [§]
Dialysis mode	HD/HDF	PD	130	13	1.07	0.661–2.16	>0.9999	ns
Previous SARS-CoV-2 infection	No	Yes	119	24	4.50	1.78–13.1	<0.001	***
mRNA vaccine	BNT162b2	mRNA-1273	118	25	2.75	1.37–6.30	0.002	**

CI, confidence interval.

[§]ns = non-significant; * = 0.033

** = 0.002

*** < 0.001.

<https://doi.org/10.1371/journal.pone.0263328.t002>

Table 3. Analysis of relative risk of poor neutralizing response to COVID-19 vaccination in naive HD patients.

Variable	Analyzed group	Reference group	All naive HD patients					Naive HD patients immunized with BNT162b2						
			n _{na}	n _{ref}	Relative Risk	95% CI	P-value	Significant ⁵	n _{na}	n _{ref}	Relative Risk	95% CI	P-value	Significant ⁵
Socio-demographic data	Gender	Female	36	70	1.07	0.738–1.50	0.8367	ns	30	54	1.11	0.789–1.51	0.6338	ns
	Age	>80 years	46	60	1.00	0.702–1.39	0.8456	ns	25	59	1.57	1.18–2.08	0.0054	**
	Body Mass Index (BMI)	<18.5	4	40	1.90	0.934–2.67	0.1217	ns	3	32	1.68	0.715–2.37	0.279	ns
Dialysis features	Body Mass Index (BMI)	18.5–24.9	36	40	0.95	0.606–1.48	>0.999	ns	29	32	1.05	0.687–1.58	>0.999	ns
		25–29.9	26	40	1.25	0.811–1.87	0.3224	ns	20	32	1.26	0.833–1.87	0.3701	ns
		>30	15	91	1.36	0.864–1.84	0.2604	ns	14	70	1.25	0.811–1.65	0.361	ns
	Smoker	Yes	61	45	1.11	0.791–1.59	0.692	ns	50	34	1.02	0.747–1.44	>0.999	ns
	Residual diuresis	Oligoanuria	43	63	1.20	0.852–1.67	0.3231	ns	34	50	1.06	0.755–1.44	0.8171	ns
	Dialysis vintage	>5 years	63	63	1.42	0.969–2.23	0.0942	ns	59	25	1.13	0.815–1.71	0.6165	ns
	Dialysis mode	HDF	72	34	1.42	0.885–1.93	0.2186	ns	55	28	1.14	0.828–1.68	0.4704	ns
	HD duration	>4 hours	70	35	1.26	0.470–1.64	0.504	ns	82	2	0.65	0.138–1.90	0.5425	ns
	Vascular access	Graft/catheter	104	2	0.56	0.295–1.29	0.5034	ns	8	66	0.75	0.318–1.24	0.44	ns
	Treatment adequacy (Kt/V)	<1.2	10	82	0.71	0.636–3.82	>0.999	ns	81	3	0.98	0.651–3.18	>0.999	ns
Comorbidities	Anticoagulant therapy	Yes	102	4	1.14	1.16–2.17	0.0119	*	14	70	1.55	1.11–1.96	0.0281	*
	Heart failure	Yes	15	91	1.68	0.899–1.76	0.2053	ns	28	56	1.24	0.886–1.66	0.2302	ns
	Coronary heart disease	Yes	33	73	1.28	0.436–1.58	>0.999	ns	6	76	1.02	0.453–1.48	>0.999	ns
	Autoimmune disease	Yes	7	99	1.01	0.643–1.49	0.7962	ns	69	15	0.98	0.707–1.60	>0.999	ns
	Hypertension	Yes	88	18	0.91	0.631–1.40	>0.999	ns	22	63	0.96	0.630–1.33	>0.999	ns
	Peripheral arterial disease	Yes	25	81	0.99	0.952–1.88	0.117	ns	39	45	1.29	0.942–1.79	0.167	ns
	Diabetes mellitus	Yes	49	57	1.33	0.754–1.52	0.6801	ns	28	56	0.97	0.668–1.33	>0.999	ns
	Pulmonary disease	Yes	35	71	1.09	0.689–1.70	0.5464	ns	10	74	0.89	0.302–2.12	>0.999	ns
	Cerebrovascular disease	Yes	12	94	1.21	0.531–1.88	0.6312	ns	4	84	1.15	0.458–1.60	>0.999	ns
	Neurodegenerative disease	Yes	4	102	1.34	0.774–1.82	0.3424	ns	8	76	1.16	0.624–1.58	0.7084	ns
Therapies	Oncologic disease	Yes	11	95	1.33	0.203–1.14	0.1719	ns	8	76	0.55	0.198–1.05	0.1176	ns
	Psychiatric disorder	Yes	9	97	0.57	0.774–1.82	0.3424	ns	11	73	1.13	0.661–1.54	0.7405	ns
	Liver disease	Yes	11	95	1.33	0.589–1.38	>0.999	ns	18	66	1.02	0.653–1.41	>0.999	ns
	Gastrointestinal disease	Yes	22	84	0.96	0.956–2.08	0.1339	ns	8	76	1.39	0.821–1.77	0.2526	ns
	Kidney transplant	Yes	8	98	1.62	0.983–2.01	0.078	ns	44	40	1.36	0.994–1.94	0.0679	ns
	Charlson comorbidity index	>8	57	49	1.38	0.715–1.57	0.631	ns	16	68	1.06	0.668–1.45	>0.999	ns
	Oral anticoagulants	Yes	21	85	1.12	0.819–1.72	0.5378	ns	58	26	1.00	0.734–1.47	>0.999	ns
	Antiaggregant drugs	Yes	69	37	1.16	0.799–1.57	0.559	ns	14	70	1.09	0.791–1.50	0.6507	ns
	ACE inhibitors/ARBs ¹	Yes	50	56	1.12	0.559–1.16	0.3183	ns	31	53	0.77	0.512–1.06	0.1546	ns
	Calcium antagonists	Yes	42	64	0.82	0.919–1.96	0.1609	ns	55	29	1.18	0.852–1.75	0.3471	ns
Therapies	Beta blockers	Yes	66	40	1.31	0.775–1.54	0.6964	ns	46	38	0.99	0.724–1.38	>0.999	ns
	Statins	Yes	58	48	1.08	0.728–1.64	0.5957	ns	14	70	1.25	0.811–1.65	0.361	ns
	Oral antidiabetic drugs	Yes	17	89	1.18	0.804–1.62	0.5144	ns	24	60	1.22	0.855–1.63	0.3137	ns
	Insulin	Yes	30	76	1.17	1.05–2.02	0.0405	*	18	66	1.25	0.852–1.66	0.2712	ns
	Immunosuppressive drugs ²	Yes	19	87	1.53	0.839–1.87	0.2238	ns	13	71	1.21	0.765–1.62	0.5276	ns
	Oral/intravenous steroids	Yes	12	94	1.38	1.18–2.29	0.018	*	6	78	1.59	0.955–1.93	0.0886	ns
	Calcineurin inhibitors ³	Yes	7	99	1.87	0.434–1.37	0.7595	ns	12	72	0.74	0.367–1.15	0.3252	ns
	Antibiotics	Yes	12	94	0.87	0.107–1.43	0.578	ns	3	81	0.50	0.092–1.23	0.2727	ns
	Antiviral drugs	Yes	3	103	0.58	0.903–2.41	0.163	ns	66	18	1.23	0.848–2.07	0.4033	ns
	Vitamin D derivatives	Yes	83	23	1.39	0.531–1.88	0.6312	ns	2	82	1.55	0.525–1.86	0.5425	ns
Nonsteroidal anti-inflam. drugs	Yes	4	102	1.34										

(Continued)

Table 3. (Continued)

Variable	Analyzed group	Reference group	All naive HD patients						Naive HD patients immunized with BNT162b2						
			n _{an}	n _{ref}	Relative Risk	95% CI	P-value	Significant ⁵	n _{an}	n _{ref}	Relative Risk	95% CI	P-value	Significant ⁵	
Vaccinations and infections	mRNA vaccine	mRNA-1273	84	22	2.88	1.47–6.55	0.0005	***	-	-	-	-	-	-	
	mRNA vaccine (matched) ⁴	mRNA-1273	22	22	3.60	1.79–8.23	0.0002	***	-	-	-	-	-	-	
	Flu vaccination	Yes	19	86	1.04	0.631–1.49	>0.999	ns	17	66	0.99	0.617–1.38	>0.999	ns	
	HBV vaccination	Yes	30	56	1.18	0.798–1.69	0.494	ns	26	43	1.16	0.812–1.62	0.4381	ns	
	HBV vaccine response	Non-responder	15	40	1.40	0.812–2.22	0.2393	ns	10	32	1.69	1.04–2.54	0.0608	ns	
	Non-SARS-CoV-2 infections	Yes	42	64	1.17	0.825–1.62	0.4261	ns	35	49	1.01	0.717–1.37	>0.999	ns	
	Non-SARS-CoV-2 resp. infection	Yes	5	101	1.44	0.667–1.94	0.3854	ns	6	78	1.24	0.574–1.65	0.655	ns	
	HBV infection	Yes	16	90	0.99	0.571–1.46	>0.999	ns	15	69	0.90	0.525–1.29	0.7655	ns	
	HCV infection	Yes	5	101	1.44	0.667–1.94	0.3854	ns	4	84	1.57	0.790–1.89	0.2932	ns	
	Bacteriemia/Fungemia	Yes	No	16	90	1.13	0.676–1.59	0.7854	ns	12	72	1.02	0.586–1.43	>0.999	ns

¹ACE, angiotensin-converting enzyme. ARB, angiotensin-receptor blocker.

²include any of steroids, cyclosporine, tacrolimus, rituximab, mycophenolate mofetil and azathioprine.

³include cyclosporine and tacrolimus.

⁴patients were matched by age (± 5 years), gender, dialysis vintage (± 3 or > 10 years), comorbidity index (± 3)

⁵ns = non-significant; * = 0.033; ** = 0.002; *** < 0.001.

<https://doi.org/10.1371/journal.pone.0263328.t003>




Fig 3. Role of avidity of plasma antibodies in the neutralization of wild-type SARS-CoV-2 and Delta variant. a) Avidity index of plasma antibodies to WT SARS-CoV-2 S in previously infected (square) and naïve (circle) healthy controls (HC), hemodialysis (HD) and peritoneal dialysis (PD) patients after two doses of mRNA vaccine. Grey areas indicate no avidity measured of plasma antibodies with ED50 titers lower than 50. A cut-off of 23%, determined from the lowest avidity index in naïve HC, and a cut-off of 50%, determined from the lowest avidity index in previously infected HC cohort, were used to distinguish low (0–23%) from moderate (23–50%) or high (>50%) avidity. Statistical significance is set as $P < 0.05$ and P-values are indicated with asterisks (* = 0.033; ** = 0.002; *** < 0.001). Shown are data from n = 2 independent experiments. b) Correlation analysis between plasma IgG avidity index to WT SARS-CoV-2 S and neutralization of WT SARS-CoV-2 in all the plasma samples collected after the second vaccine dose. c) Percentages of participants having plasma antibodies with high, moderate, low or no avidity to WT SARS-CoV-2 S after two doses of mRNA vaccine. Participants are shown as a total (bold) or divided by level of neutralization of WT SARS-CoV-2 (no, low, moderate, high). Total number of participants within each group is shown at the top of each bar. d) Percentages of participants having plasma antibodies with high, moderate, low or no neutralization of Delta SARS-CoV-2 after two doses of mRNA vaccine. Participants are shown as a total (bold) or divided by level of avidity to WT SARS-CoV-2 S (no, low, moderate, high). Total number of participants within each group is shown at the top of each bar.

<https://doi.org/10.1371/journal.pone.0263328.g003>

showing low- or non-neutralizing antibodies against the Delta and other SARS-CoV-2 variants (Fig 3D and S5 Fig, panels b-f).

Rapid decay of neutralizing antibody in 7 months after vaccination

We followed 47 HC and 133 dialysis patients (121 HD and 12 PD) up to 7 months after vaccination (mean 5 months) and measured plasma titers and avidity of antibodies to SARS-CoV-2 S and their neutralizing activity against the vaccine-matched SARS-CoV-2 and Delta variant (Table 4). We found that 12.8% and 77.4% of HC and dialysis patients had low or undetectable S-specific IgG with a 4.2- and 5.2 fold- average reduction of IgG titers compared to the peak of

Table 4. Follow-up analysis of SARS-CoV-2 S IgG titer, wild-type SARS-CoV-2 and Delta neutralization, and avidity index.

	Nr.	Months after 2 nd dose	S IgG titer	Fold reduction		WT neutralization	Fold reduction		Delta neutralization	Fold reduction		Avidity index	Variation (%)	
				Nr. ^b	Mean (min-max)		Responders (%) ^c	Nr. ^d		Mean (min-max)	Responders (%) ^e		Nr. ^d	Mean (min-max)
Healthy controls	47	4.6 (4.5–4.9)	41 (87.2)	47	4.2 (1.6–14.9)	28 (59.6)	47	12.7 (0.2–100)	13 (27.7)	45	6.2 (0.4–28.4)	17 (36.2)	47	1.2 (-29.6–36)
Prev. infected ^{P/B}	24	4.7 (4.5–4.9)	24 (100)	24	4.2 (1.6–14.9)	22 (91.7)	24	12.7 (0.2–100)	12 (50)	22	6.2 (0.4–28.4)	5 (20.8)	24	1.2 (-29.6–36)
Naïve ^{P/B}	23	4.6 (4.5–4.9)	17 (73.9)	23	3.3 (1.6–9)	6 (26.1)	23	4.7 (0.2–17.4)	1 (4.3)	23	2.6 (0.6–8.2)	12 (52.2)	23	-4.4 (-29.6–14)
Dialysis patients	133	5.3 (3.4–6.8)	30 (22.6)	126	5.2 (1.6–14.9)	22 (16.5)	87	21.1 (3.4–100)	14 (10.5)	48	9.8 (0.4–28.4)	51 (38.3)	103	7.1 (-12.8–36)
Hemodialysis	121	5.3 (3.4–6.8)	28 (23.1)	114	10.6 (0.5–83.2)	21 (17.4)	79	11.7 (0.1–100)	13 (10.7)	45	13.3 (1.1–100)	43 (35.5)	93	2.9 (-45.6–40.9)
Prev. infected	21	5.1 (4–6.7)	16 (76.2)	20	9.8 (1.1–72.1)	16 (76.2)	18	10.9 (1–100)	11 (52.4)	18	12 (1.1–100)	5 (23.8)	19	1.9 (-45.6–40.9)
Pfizer-BioNTech	20	5.2 (4–6.7)	15 (75)	19	5.1 (1.8–15.2)	15 (75)	17	9 (1.9–45.4)	10 (50)	17	15.7 (1.1–67)	4 (20)	18	-3.2 (-15.7–7)
Moderna	1	4.0 (4.0–4.0)	1 (100)	1	5.2 (1.8–15.2)	1 (100)	1	9.3 (1.9–45.4)	1 (100)	1	16.4 (1.1–67)	1 (100)	1	-3.7 (-15.7–7)
Naïve	100	5.3 (3.4–6.8)	12 (12)	94	3.4 (3.4–3.4)	5 (5)	61	4.2 (4.2–4.2)	2 (2)	27	3.1 (3.1–3.1)	38 (38)	74	5.3 (5.3–5.3)
Pfizer-BioNTech	80	5.3 (3.4–6.8)	8 (10)	74	10.8 (1.1–72.1)	2 (2.5)	42	11.5 (1–100)	1 (1.3)	14	9.5 (2.2–100)	26 (32.5)	54	3.2 (-45.6–40.9)
Moderna	20	5.5 (3.7–6.2)	4 (20)	20	11 (1.1–72.1)	3 (15)	19	10 (1–100)	1 (5)	13	12 (2.2–100)	12 (60)	20	1.1 (-45.6–40.9)
Peritoneal dialysis	12	5.4 (4.3–6.2)	2 (16.7)	12	9.8 (2.2–20.6)	1 (8.3)	8	14.8 (1.1–59.8)	1 (8.3)	3	6.8 (2.4–37)	8 (66.7)	10	8.8 (-12.5–31)
Prev. infected ^{P/B}	1	5.3 (5.3–5.3)	0 (0)	1	18.8 (0.5–83.2)	0 (0)	1	19.3 (0.1–100)	0 (0)	1	32.8 (5.6–82.8)	1 (100)	1	12.9 (-13.2–31.4)
Naïve	11	5.4 (4.3–6.2)	2 (18.2)	11	83.2 (83.2–83.2)	1 (9.1)	7	100 (100–100)	1 (9.1)	2	82.8 (82.8–82.8)	7 (63.6)	9	21.2 (21.2–21.2)
Pfizer-BioNTech	9	5.5 (4.8–6)	2 (22.2)	9	12.9 (0.5–30.4)	1 (11.1)	5	7.8 (0.1–23.5)	1 (11.1)	1	7.8 (5.6–9.9)	6 (66.7)	7	12 (-13.2–31.4)
Moderna	2	5.2 (4.3–6.2)	0 (0)	2	14.5 (0.5–30.4)	0 (0)	2	7.1 (0.1–23.5)	0 (0)	1	9.9 (9.9–9.9)	1 (50)	2	14 (-13.2–31.4)

(Continued)

Table 4. (Continued)

	Nr.	Months after 2 nd dose	S IgG titer	Fold reduction		WT neutralization	Fold reduction		Delta neutralization	Fold reduction		Avidity index	Variation (%)	
		Mean (min-max)	Responders (%) ^a	Nr. ^b	Mean (min-max)	Responders (%) ^c	Nr. ^d	Mean (min-max)	Responders (%) ^e	Nr. ^d	Mean (min-max)	Responders (%) ^e	Nr. ^f	Mean (min-max)
Total	180	5.1 (3.4–6.8)	71 (39.4)	173	5.8 (5.1–6.5)	50 (27.8)	134	9.7 (8–11.5)	27 (15)	93	5.6 (5.6–5.6)	68 (37.8)	150	4.9 (0.8–9.1)

^{P/B}Pfizer-BioNTech.

^aED50>1'000.

^bparticipants with ED50 titers >50 at 2 weeks after the second dose were counted.

^cID50>200.

^dparticipants with neutralizing ID50 titers >80 at 2 weeks after the second dose were counted.

^eavidity index variation >5%.

^fparticipants with detectable titers (ED50>50) at 2 weeks and up to 7 months after the second dose were counted.

<https://doi.org/10.1371/journal.pone.0263328.t004>

the antibody response after the second vaccine dose (Fig 4A and S6 Fig, panel a). This titer decay was paralleled by a 12.7- and 21-fold average reduction of neutralizing antibodies to wild-type SARS-CoV-2 in HC and dialysis patients, which resulted in 40% and 84% of non-responders, respectively (Fig 4C and S6 Fig, panel c). In the case of antibodies neutralizing the Delta variant, we observed a 6.2- and 9.8-average reduction in HC and dialysis patients, which increased the fraction of non-responders to 72% and 90%, respectively (Fig 4D and S6 Fig, panel d). In line with the analysis at 2 weeks after the second dose, the most affected patients were those in hemodialysis, naïve to SARS-CoV-2 and vaccinated with BNT-162b2, with 97.5% and 99% of non-responders to WT SARS-CoV-2 and Delta variant, respectively (Table 4). Interestingly, only 36.2% and 38.8% of HC and dialysis patients showed an increase in the avidity index of SARS-CoV-2 S IgG, which, however, maintained its correlation with the neutralizing activity (Fig 4B and S6 Fig, panels b and e). Importantly, despite the overall waning of the antibody response during the follow-up, no new cases of SARS-CoV-2 infection were diagnosed in both HC and dialysis patients after vaccination.

Discussion

In this phase of the COVID-19 pandemic where SARS-CoV-2 variants are rapidly spreading worldwide, we urgently need to understand the efficacy of vaccines administered so far in high-risk immunocompromised populations, in particular patients with chronic diseases [39, 40]. In this study, we provide evidence that dialysis patients do not develop sufficient levels of neutralizing antibodies against SARS-CoV-2 variants after two doses of mRNA vaccines.

The analysis of plasma SARS-CoV-2-specific IgG showed an overall response to mRNA vaccine in 94.4% dialysis patients, which is comparable to data shown in other reports (70.5–96%) [19, 22, 41, 42]. Nevertheless, we observed reduced antibody levels after both the first and the second dose in HD patients who were naïve to SARS-CoV-2, a finding consistent with a delayed immune response in this population [19, 43, 44]. These data suggest that measuring serum antibody titers to SARS-CoV-2 in dialysis patients could help clinicians to identify vaccine non-responders and guide clinical decision-making. However, currently available serological tests detect antibodies against the Spike of the Wuhan SARS-CoV-2 and do not define




Fig 4. Follow-up analysis of SARS-CoV-2 S IgG, neutralizing titers and avidity index. a) Plasma IgG titers (ED50) to SARS-CoV-2 S after two doses of Pfizer/BioNTech (P/B, red, light blue and dark orange) or Moderna (M, dark blue and light orange) vaccines in samples collected from previously infected (square) and naïve (circle) healthy controls (HC, red), hemodialysis (HD, blue) and peritoneal dialysis (PD, orange) patients 2 weeks and up to 7 months (indicated mean 5 months) after vaccination. Cut-offs are shown as explained in Fig 1. Statistical significance is set as $P < 0.05$ and P-values are indicated with asterisks (* = 0.033; ** = 0.002; *** < 0.001). b) Avidity index of plasma antibodies to WT SARS-CoV-2 S. Cut-offs are shown as explained in Fig 3C and 3D) Neutralizing antibody titers (ID50) against pseudotyped VSV viruses harboring wild-type (D614) SARS-CoV-2 S (c) or Delta variant (d). Cut-offs are shown as explained in Fig 2.

<https://doi.org/10.1371/journal.pone.0263328.g004>

the level of neutralizing activity against the virus, which is considered a more relevant serological correlate of protection [45].

In this study, we found that 49% of dialysis patients had low or undetectable levels of neutralizing antibodies against the vaccine-matched SARS-CoV-2 with a significant reduction in neutralizing titers against different circulating variants in dialysis patients, which is consistent with an immune evasion by the virus [27, 35, 46]. Based on these data, we calculated that the fraction of dialysis patients non-responding to the vaccine further increased to 77% in the case of the Delta variant, suggesting that the great majority of the patients immunized with two doses of an mRNA vaccine may not develop a protective antibody response against SARS-CoV-2 variants. Importantly, follow-up data collected up to 7 months after vaccination showed a rapid decay of neutralizing antibodies to both wild-type SARS-CoV-2 and Delta variant, which increased the fraction of non-responders to 84% and 90%, respectively. These data suggest that dialysis patients have an increased risk of infection with possible severe outcomes over time from vaccination, an association which has been recently described in a US veteran

cohort followed up to 6 months after vaccination [47]. However, we did not diagnose new cases of SARS-CoV-2 infection during the follow-up, which might be explained by the lowest incidence observed in Switzerland in the spring-summer season, and by the strict preventive measures adopted by the dialysis patients that might have compensated for the waning of the vaccine-induced antibodies.

Despite the huge discrepancy between the fraction of non-responders based on antibody levels and their neutralizing activity, we observed a high correlation between these titers, which may help to define proper cut-offs in the serological tests used in clinical settings to identify non-responding patients. Furthermore, the correlation observed between avidity and neutralization suggests that vaccine-induced antibodies require high avidity to neutralize SARS-CoV-2. Therefore, the identification of non-responders may be guided by a SARS-CoV-2 avidity test, as successfully implemented for other infectious diseases [48–52].

We identified that having never been exposed to SARS-CoV-2 is the major factor that increases the risk of a poor antibody response after vaccination, especially in HD patients. Indeed, naïve individuals were shown to develop lower titers of neutralizing antibodies to SARS-CoV-2 variants compared to previously infected individuals after two mRNA vaccine doses [35], indicating that vaccination may boost the neutralizing antibody response in dialysis patients after infection [53]. This effect can be explained by the higher antibody avidity that previously infected patients gained over time compared to naïve patients, who instead showed a delayed and low-avidity response. These findings are consistent with a slower affinity maturation of SARS-CoV-2 S-specific B cells, which appears to be a peculiar immunological feature of the dialysis population.

A recent report by Carr et al [54], showed that naïve HD patients vaccinated with BNT162b2 developed higher neutralizing antibodies against SARS-CoV-2 variants compared to Astra Zeneca—Oxford University AZD1222 vaccine, suggesting that mRNA vaccines provide a higher level of protection in dialysis patients compared to adenovirus-based vaccines. However, a difference in efficacy may exist among mRNA vaccines, as we found that being immunized with BNT162b2 instead of mRNA-1273 was the second most relevant risk factor for a poor response in the dialysis population, especially in naïve HD patients. This difference can be explained by the lower mRNA amount provided by one dose of BNT162b2 compared to mRNA-1273 [11, 12]. Indeed, the mRNA-1273 vaccine was found to be more reactogenic [55] and to induce higher antibody titers in healthy individuals as well as in the more vulnerable elderly population [56–59].

Similarly to other studies analyzing the factors that were associated to low antibody levels after COVID-19 vaccination in dialysis patients, we also found a few risk factors of poor neutralizing antibody response to mRNA vaccines, including age, heart failure and immunosuppression [14, 20, 22, 23, 43, 60–62]. However, other factors, such as gender and dialysis vintage [43], dialysis adequacy measured by Kt/V [20], comorbidities [14] and non-responsiveness to Hepatitis B vaccination [20, 23] were not identified as risk factors in our study.

Our study has some limitations including a missing control group of patients matched by age, gender and comorbidities, without ESKD, while our control group was composed by younger individuals with few or no comorbidities. Another limitation was the low number of PD patients that did not allow to make subgroup analyses of risk factors of poor response. In addition, our patients were unbalanced in terms of type of mRNA vaccine received (BNT162b2 or mRNA-1273), although we had sufficient statistical power to make a comparison, even in a subgroup of patients matched by age, gender and comorbidity index.

In conclusion, our study demonstrates, at the functional level, that mRNA vaccines induce a poor neutralizing and rapidly decaying antibody response against SARS-CoV-2 variants in dialysis patients, in particular in naïve HD patients immunized with BNT162b2. Our findings

support the need of an additional boost, preferentially with a high-dose mRNA vaccine, in this population [62–67], which, however, need to be continuously monitored with proper serological tests that measure not only the serum antibody levels, but also their neutralizing activity, either directly or indirectly through an avidity test. Finally, our data suggest that some patients may not respond efficiently even after an additional boost and, therefore, in case of SARS-CoV-2 infection, they should be considered for other therapeutic strategies, including early immunotherapy with monoclonal antibodies.

Supporting information

S1 Fig. Plasma antibody titers to SARS-CoV-2 after one dose of mRNA vaccine. a) Plasma IgG titers (ED50) to SARS-CoV-2 S after one dose of Pfizer/BioNTech (P/B) or Moderna (M) vaccines in previously infected (square) and naïve (circle) healthy controls (HC, red) and hemodialysis (HD, blue) patients. Grey areas indicate non-specific IgG titers <50, a cut-off that was determined on non-specific binding to uncoated ELISA plates. An additional cut-off of 1'000, determined from the lowest titers in previously infected HC, was used to distinguish low (50–1'000) from high (>1'000) IgG titers. Statistical significance is set as $P < 0.05$ and P-values are indicated with asterisks (* = 0.033; ** = 0.002; *** < 0.001). b) Percentages of participants with high, low or no plasma IgG to SARS-CoV-2 S after one dose of mRNA-vaccine. Total number of participants within each cohort of HC and dialysis patients (DP) and within each subgroup is shown at the top of each bar. (TIF)

S2 Fig. Kinetics of plasma antibody titers to SARS-CoV-2 in healthy controls and dialysis patients. a) Plasma IgG, IgM and IgA titers to SARS-CoV-2 RBD. b) Plasma IgM and IgA titers (ED50) to SARS-CoV-2 S measured before vaccination (T0), after one (T1) or two (T2) vaccine doses. c) Plasma IgG, IgM and IgA titers to SARS-CoV-2 Nucleoprotein (N). (TIF)

S3 Fig. Correlation of plasma IgG titers, ACE2 inhibition and neutralization against wild-type SARS-CoV-2. a) Neutralization of WT SARS-CoV-2 pseudotyped VSV by four representative plasma samples showing high, moderate, low or no neutralization. b) Inhibition of RBD binding to human ACE2 by plasma antibodies in previously infected (square) and naïve (circle) healthy controls (HC), hemodialysis (HD) and peritoneal dialysis (PD) patients after two doses of mRNA vaccine. Grey areas indicate no ACE2 inhibition by plasma antibodies with BD80 titers lower than 10. A cut-off of 80, determined from the lowest inhibition capacity in HC, was used to distinguish low (10–80) from high (>80) inhibition. Statistical significance is set as $P < 0.05$ and P-values are indicated with asterisks (* = 0.033; ** = 0.002; *** < 0.001). Shown are data from $n = 2$ independent experiments. c) Correlation analysis between plasma IgG titers index and WT SARS-CoV-2 neutralization in all the plasma samples collected after the second vaccine dose. d) Correlation analysis between ACE2 inhibition and plasma RBD IgG titers index in all the plasma samples collected after the second vaccine dose. e) Correlation analysis between ACE2 inhibition and WT SARS-CoV-2 neutralization in all the plasma samples collected after the second vaccine dose. (TIF)

S4 Fig. Comparison of neutralizing antibody titers against wild-type SARS-CoV-2 and other variants. a) Fraction of participants with neutralizing titers against WT SARS-CoV-2 ($ID_{50} > 40$) showing complete loss of neutralization against Alpha (B.1.1.7), Beta (B.1.351), Gamma (P.1), Delta (B.1.617.2), Epsilon (B.1.429), Kappa (B.1.617.1) and Lambda (C.37) variants. b-g) Fold change analysis of neutralizing titers against WT and Beta (b), Epsilon (c),

Kappa (d), Alpha (e), Gamma (f) and Lambda (g) SARS-CoV-2 in 28 HC, 36 HD and 5 PD patients with ID50 neutralizing titers against WT SARS-CoV-2 greater than 80. h) Side-by-side comparison of neutralizing titers against WT and Lambda (C.37) SARS-CoV-2 variant in 28 HC, 47 HD and 5 PD patients.

(TIF)

S5 Fig. Correlation of avidity and neutralization of plasma antibodies to wild-type SARS-CoV-2 and other variants. a) Correlation analysis between plasma IgG titers index and SARS-CoV-2 S IgG avidity in all the plasma samples collected after the second vaccine dose. b-f) Percentages of participants having plasma antibodies with high, moderate, low or no neutralization of Alpha (b), Beta (c), Gamma (d), Epsilon (e) and Kappa (f) SARS-CoV-2 after two doses of mRNA vaccine. Participants are shown as a total (bold) or divided by level of avidity to WT SARS-CoV-2 S (no, low, moderate, high). Total number of participants within each group is shown at the top of each bar.

(TIF)

S6 Fig. Comparison of variation of SARS-CoV-2 S IgG, neutralizing titers and avidity index. a-d) Fold change analysis of SARS-CoV-2 S IgG (a), avidity index (b), neutralizing titers against WT SARS-CoV-2 (c) and Delta variant (d) measured in samples collected at 2 weeks and up to 7 months (average 5 months) after the second vaccine dose. n indicates the number of analyzed participants as explained in Table 4. e) Correlation analysis between plasma IgG avidity index to WT SARS-CoV-2 S and neutralization of WT SARS-CoV-2 in all the 180 plasma samples collected up to 7 months after the second vaccine dose.

(TIF)

S1 File.

(ZIP)

Acknowledgments

We would like to thank all the nurses of the Ente Ospedaliero Cantonale, who helped collecting data and blood samples: Gianna Croppi, Sara Candolfi, Marie-Eve Brodeur, Chiara Tettamanti, Giorgia Esposito, Fabio Capocasale, Elisa Capitale, Marianna Vitini, Pedrag Lazarevic, Gaetano Donato, Fabienne Poncetta, Chiara Mattei. We thank Sarita Prosperi and Barbara Guggiari-De Rosa for organizational support. A special thanks to all the dialysis patients who accepted to participate to this study.

Author Contributions

Conceptualization: Olivier Giannini, Soraya Lavorato-Hadjeres, Paolo Ferrari, Alessandro Ceschi, Pietro Cippà, Davide Corti, Antonio Lanzavecchia, Luca Piccoli.

Data curation: Jessica Bassi, Olivier Giannini, Chiara Silacci-Fregni, Laura Pertusini, Paolo Hitz, Tatiana Terrot, Yves Franzosi, Nicola Beria, Mariangela Ventresca, Alberto Badellino, Elisabetta Lecchi, Tecla Bonora, Matteo Mattiolo, Guido Trinci, Daniela Garzoni, Giuseppe Bonforte, Valentina Forni-Ogna, Davide Giunzioni, Lorenzo Berwert, Luca Piccoli.

Formal analysis: Jessica Bassi, Chiara Silacci-Fregni, Francesco Muoio, Luca Piccoli.

Funding acquisition: Olivier Giannini, Ravindra K. Gupta.

Investigation: Jessica Bassi, Olivier Giannini, Nicola Beria, Mariangela Ventresca, Alberto Badellino, Soraya Lavorato-Hadjeres, Elisabetta Lecchi, Tecla Bonora, Matteo Mattiolo, Guido Trinci, Daniela Garzoni, Giuseppe Bonforte, Valentina Forni-Ogna, Davide

Giunzioni, Lorenzo Berwert, Paolo Ferrari, Pietro Cippà, Davide Corti, Antonio Lanzavecchia, Luca Piccoli.

Methodology: Jessica Bassi, Olivier Giannini, Chiara Silacci-Fregni, Laura Pertusini, Francesco Muoio, Christian Saliba, Marcel Meury, Exequiel A. Dellota, Jr., Josh R. Dillen, Patrick Hernandez, Nadine Czudnochowski, Elisabetta Cameroni, Nicola Beria, Mariangela Ventresca, Alberto Badellino, Soraya Lavorato-Hadjeres, Elisabetta Lecchi, Tecla Bonora, Matteo Mattiolo, Guido Trinci, Daniela Garzoni, Giuseppe Bonforte, Valentina Forni-Ogna, Davide Giunzioni, Lorenzo Berwert, Ravindra K. Gupta, Luca Piccoli.

Project administration: Olivier Giannini, Laura Pertusini, Luca Piccoli.

Resources: Olivier Giannini, Christian Saliba, Marcel Meury, Exequiel A. Dellota, Jr., Josh R. Dillen, Patrick Hernandez, Nadine Czudnochowski, Elisabetta Cameroni, Ravindra K. Gupta, Alessandro Ceschi, Davide Corti.

Software: Laura Pertusini, Paolo Hitz, Tatiana Terrot, Yves Franzosi.

Supervision: Olivier Giannini, Laura Pertusini, Pietro Cippà, Davide Corti, Antonio Lanzavecchia, Luca Piccoli.

Validation: Jessica Bassi, Chiara Silacci-Fregni, Ravindra K. Gupta, Luca Piccoli.

Visualization: Luca Piccoli.

Writing – original draft: Jessica Bassi, Olivier Giannini, Luca Piccoli.

Writing – review & editing: Olivier Giannini, Paolo Ferrari, Alessandro Ceschi, Pietro Cippà, Davide Corti, Antonio Lanzavecchia, Luca Piccoli.

References

1. Zoccali C, Vanholder R, Massy ZA, Ortiz A, Sarafidis P, Dekker FW, et al. The systemic nature of CKD. *Nat Rev Nephrol.* 2017; 13(6):344–58. Epub 2017/04/25. <https://doi.org/10.1038/nrneph.2017.52> PMID: 28435157.
2. Dalrymple LS, Go AS. Epidemiology of acute infections among patients with chronic kidney disease. *Clin J Am Soc Nephrol.* 2008; 3(5):1487–93. Epub 2008/07/25. <https://doi.org/10.2215/CJN.01290308> PMID: 18650409; PubMed Central PMCID: PMC4571152.
3. Henry BM, Lippi G. Chronic kidney disease is associated with severe coronavirus disease 2019 (COVID-19) infection. *Int Urol Nephrol.* 2020; 52(6):1193–4. Epub 2020/03/31. <https://doi.org/10.1007/s11255-020-02451-9> PMID: 32222883; PubMed Central PMCID: PMC7103107.
4. Cippa PE, Cugnata F, Ferrari P, Brombin C, Ruinelli L, Bianchi G, et al. A data-driven approach to identify risk profiles and protective drugs in COVID-19. *Proc Natl Acad Sci U S A.* 2021; 118(1). Epub 2020/12/12. <https://doi.org/10.1073/pnas.2016877118> PMID: 33303654; PubMed Central PMCID: PMC7817222.
5. Anand S, Montez-Rath M, Han J, Bozeman J, Kerschmann R, Beyer P, et al. Prevalence of SARS-CoV-2 antibodies in a large nationwide sample of patients on dialysis in the USA: a cross-sectional study. *Lancet.* 2020; 396(10259):1335–44. Epub 2020/09/25. [https://doi.org/10.1016/S0140-6736\(20\)32009-2](https://doi.org/10.1016/S0140-6736(20)32009-2) PMID: 32987007; PubMed Central PMCID: PMC7518804.
6. Corbett RW, Blakey S, Nitsch D, Loucaidou M, McLean A, Duncan N, et al. Epidemiology of COVID-19 in an Urban Dialysis Center. *J Am Soc Nephrol.* 2020; 31(8):1815–23. Epub 2020/06/21. <https://doi.org/10.1681/ASN.2020040534> PMID: 32561681; PubMed Central PMCID: PMC7460899.
7. Jager KJ, Kramer A, Chesnaye NC, Couchoud C, Sanchez-Alvarez JE, Garneata L, et al. Results from the ERA-EDTA Registry indicate a high mortality due to COVID-19 in dialysis patients and kidney transplant recipients across Europe. *Kidney Int.* 2020; 98(6):1540–8. Epub 2020/09/27. <https://doi.org/10.1016/j.kint.2020.09.006> PMID: 32979369; PubMed Central PMCID: PMC7560263.
8. Hilbrands LB, Duivenvoorden R, Vart P, Franssen CFM, Hemmelder MH, Jager KJ, et al. COVID-19-related mortality in kidney transplant and dialysis patients: results of the ERACODA collaboration. *Nephrol Dial Transplant.* 2020; 35(11):1973–83. Epub 2020/11/06. <https://doi.org/10.1093/ndt/gfaa261> PMID: 33151337; PubMed Central PMCID: PMC7665620.

9. The Renal Association B, UK. UK Renal Registry (2020) COVID-19 surveillance report for renal centres in the UK: All regions and centres.
10. Robinson BM, Guedes M, Alghonaim M, Cases A, Dasgupta I, Gan L, et al. Worldwide Early Impact of COVID-19 on Dialysis Patients and Staff and Lessons Learned: A DOPPS Roundtable Discussion. *Kidney Med.* 2021; 3(4):619–34. Epub 2021/05/20. <https://doi.org/10.1016/j.xkme.2021.03.006> PMID: 34007963; PubMed Central PMCID: PMC8120787.
11. Baden LR, El Sahly HM, Essink B, Kotloff K, Frey S, Novak R, et al. Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N Engl J Med.* 2021; 384(5):403–16. Epub 2020/12/31. <https://doi.org/10.1056/NEJMoa2035389> PMID: 33378609; PubMed Central PMCID: PMC7787219.
12. Polack FP, Thomas SJ, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N Engl J Med.* 2020; 383(27):2603–15. Epub 2020/12/11. <https://doi.org/10.1056/NEJMoa2034577> PMID: 33301246; PubMed Central PMCID: PMC7745181.
13. Dagan N, Barda N, Kepten E, Miron O, Perchik S, Katz MA, et al. BNT162b2 mRNA Covid-19 Vaccine in a Nationwide Mass Vaccination Setting. *N Engl J Med.* 2021; 384(15):1412–23. Epub 2021/02/25. <https://doi.org/10.1056/NEJMoa2101765> PMID: 33626250; PubMed Central PMCID: PMC7944975.
14. Butt AA, Omer SB, Yan P, Shaikh OS, Mayr FB. SARS-CoV-2 Vaccine Effectiveness in a High-Risk National Population in a Real-World Setting. *Ann Intern Med.* 2021. Epub 2021/07/20. <https://doi.org/10.7326/M21-1577> PubMed Central PMCID: PMC8381771. PMID: 34280332
15. Malard F, Gaugler B, Gozlan J, Bouquet L, Fofana D, Siblany L, et al. Weak immunogenicity of SARS-CoV-2 vaccine in patients with hematologic malignancies. *Blood Cancer J.* 2021; 11(8):142. Epub 2021/08/12. <https://doi.org/10.1038/s41408-021-00534-z> PMID: 34376633; PubMed Central PMCID: PMC8353615.
16. Louapre C, Ibrahim M, Maillart E, Abdi B, Papeix C, Stankoff B, et al. Anti-CD20 therapies decrease humoral immune response to SARS-CoV-2 in patients with multiple sclerosis or neuromyelitis optica spectrum disorders. *J Neurol Neurosurg Psychiatry.* 2021. Epub 2021/08/04. <https://doi.org/10.1136/jnnp-2021-326904> PubMed Central PMCID: PMC8331322. PMID: 34341142
17. Boyarsky BJ, Werbel WA, Avery RK, Tobian AAR, Massie AB, Segev DL, et al. Antibody Response to 2-Dose SARS-CoV-2 mRNA Vaccine Series in Solid Organ Transplant Recipients. *JAMA.* 2021; 325(21):2204–6. Epub 2021/05/06. <https://doi.org/10.1001/jama.2021.7489> PMID: 33950155; PubMed Central PMCID: PMC8100911.
18. Ikizler TA, Coates PT, Rovin BH, Ronco P. Immune response to SARS-CoV-2 infection and vaccination in patients receiving kidney replacement therapy. *Kidney Int.* 2021; 99(6):1275–9. Epub 2021/05/02. <https://doi.org/10.1016/j.kint.2021.04.007> PMID: 33931226; PubMed Central PMCID: PMC8055920.
19. Rincon-Arevalo H, Choi M, Stefanski AL, Halleck F, Weber U, Szelinski F, et al. Impaired humoral immunity to SARS-CoV-2 BNT162b2 vaccine in kidney transplant recipients and dialysis patients. *Sci Immunol.* 2021; 6(60). Epub 2021/06/17. <https://doi.org/10.1126/sciimmunol.abj1031> PMID: 34131023.
20. Danthu C, Hantz S, Dahlem A, Duval M, Ba B, Guibbert M, et al. Humoral Response after SARS-CoV-2 mRNA Vaccination in a Cohort of Hemodialysis Patients and Kidney Transplant Recipients. *J Am Soc Nephrol.* 2021. Epub 2021/06/18. <https://doi.org/10.1681/ASN.2021040490> PMID: 34135083
21. Public Health England, Immunization against infection disease: the Green book: Public Health England; 2021.
22. Grupper A, Sharon N, Finn T, Cohen R, Israel M, Agbaria A, et al. Humoral Response to the Pfizer BNT162b2 Vaccine in Patients Undergoing Maintenance Hemodialysis. *Clin J Am Soc Nephrol.* 2021. Epub 2021/04/08. <https://doi.org/10.2215/CJN.03500321> PMID: 33824157.
23. Zitt E, Davidovic T, Schimpf J, Abbassi-Nik A, Mutschlechner B, Ulmer H, et al. The Safety and Immunogenicity of the mRNA-BNT162b2 SARS-CoV-2 Vaccine in Hemodialysis Patients. *Front Immunol.* 2021; 12:704773. Epub 2021/07/06. <https://doi.org/10.3389/fimmu.2021.704773> PMID: 34220867; PubMed Central PMCID: PMC8242233.
24. Alcazar-Arroyo R, Portoles J, Lopez-Sanchez P, Zalamea F, Furaz K, Mendez A, et al. Rapid decline of anti-SARS-CoV-2 antibodies in patients on haemodialysis: the COVID-FRIAT study. *Clin Kidney J.* 2021; 14(7):1835–44. Epub 2021/07/03. <https://doi.org/10.1093/ckj/sfab048> PMID: 34211708; PubMed Central PMCID: PMC7989535.
25. Krueger KM, Ison MG, Ghossein C. Practical Guide to Vaccination in All Stages of CKD, Including Patients Treated by Dialysis or Kidney Transplantation. *Am J Kidney Dis.* 2020; 75(3):417–25. Epub 2019/10/06. <https://doi.org/10.1053/j.ajkd.2019.06.014> PMID: 31585683.
26. Chaves SS, Daniels D, Cooper BW, Malo-Schlegel S, Macarthur S, Robbins KC, et al. Immunogenicity of hepatitis B vaccine among hemodialysis patients: effect of revaccination of non-responders and duration of protection. *Vaccine.* 2011; 29(52):9618–23. Epub 2011/11/03. <https://doi.org/10.1016/j.vaccine.2011.10.057> PMID: 22044739.

27. Mlcochova P, Kemp S, Dhar MS, Papa G, Meng B, Ferreira I, et al. SARS-CoV-2 B.1.617.2 Delta variant replication and immune evasion. *Nature*. 2021. Epub 2021/09/07. <https://doi.org/10.1038/s41586-021-03944-y> PMID: 34488225
28. Kirchdoerfer RN, Wang N, Pallesen J, Wrapp D, Turner HL, Cottrell CA, et al. Stabilized coronavirus spikes are resistant to conformational changes induced by receptor recognition or proteolysis. *Sci Rep*. 2018; 8(1):15701. Epub 2018/10/26. <https://doi.org/10.1038/s41598-018-34171-7> PMID: 30356097; PubMed Central PMCID: PMC6200764.
29. Pallesen J, Wang N, Corbett KS, Wrapp D, Kirchdoerfer RN, Turner HL, et al. Immunogenicity and structures of a rationally designed prefusion MERS-CoV spike antigen. *Proc Natl Acad Sci U S A*. 2017; 114(35):E7348–E57. Epub 2017/08/16. <https://doi.org/10.1073/pnas.1707304114> PMID: 28807998; PubMed Central PMCID: PMC5584442.
30. Walls AC, Park YJ, Tortorici MA, Wall A, McGuire AT, Velesler D. Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell*. 2020; 181(2):281–92 e6. Epub 2020/03/11. <https://doi.org/10.1016/j.cell.2020.02.058> PMID: 32155444; PubMed Central PMCID: PMC7102599.
31. Gregson J, Rhee SY, Datir R, Pillay D, Perno CF, Derache A, et al. Human Immunodeficiency Virus-1 Viral Load Is Elevated in Individuals With Reverse-Transcriptase Mutation M184V/I During Virological Failure of First-Line Antiretroviral Therapy and Is Associated With Compensatory Mutation L74I. *J Infect Dis*. 2020; 222(7):1108–16. Epub 2019/11/28. <https://doi.org/10.1093/infdis/jiz631> PMID: 31774913; PubMed Central PMCID: PMC7459140.
32. Forloni M, Liu AY, Wajapeyee N. Creating Insertions or Deletions Using Overlap Extension Polymerase Chain Reaction (PCR) Mutagenesis. *Cold Spring Harb Protoc*. 2018;2018(8). Epub 2018/08/03. <https://doi.org/10.1101/pdb.prot097758> PMID: 30068588.
33. Takada A, Robison C, Goto H, Sanchez A, Murti KG, Whitt MA, et al. A system for functional analysis of Ebola virus glycoprotein. *Proc Natl Acad Sci U S A*. 1997; 94(26):14764–9. Epub 1998/02/07. <https://doi.org/10.1073/pnas.94.26.14764> PMID: 9405687; PubMed Central PMCID: PMC25111.
34. Riblett AM, Blomen VA, Jae LT, Altamura LA, Doms RW, Brummelkamp TR, et al. A Haploid Genetic Screen Identifies Heparan Sulfate Proteoglycans Supporting Rift Valley Fever Virus Infection. *J Virol*. 2016; 90(3):1414–23. Epub 2015/11/20. <https://doi.org/10.1128/JVI.02055-15> PMID: 26581979; PubMed Central PMCID: PMC4719632.
35. McCallum M, Bassi J, De Marco A, Chen A, Walls AC, Di Iulio J, et al. SARS-CoV-2 immune evasion by the B.1.427/B.1.429 variant of concern. *Science*. 2021; 373(6555):648–54. Epub 2021/07/03. <https://doi.org/10.1126/science.abi7994> PMID: 34210893.
36. Piccoli L, Park YJ, Tortorici MA, Czudnochowski N, Walls AC, Beltramello M, et al. Mapping Neutralizing and Immunodominant Sites on the SARS-CoV-2 Spike Receptor-Binding Domain by Structure-Guided High-Resolution Serology. *Cell*. 2020; 183(4):1024–42 e21. Epub 2020/09/30. <https://doi.org/10.1016/j.cell.2020.09.037> PMID: 32991844; PubMed Central PMCID: PMC7494283.
37. Legros V, Denolly S, Vogrig M, Boson B, Siret E, Rigai J, et al. A longitudinal study of SARS-CoV-2-infected patients reveals a high correlation between neutralizing antibodies and COVID-19 severity. *Cell Mol Immunol*. 2021; 18(2):318–27. Epub 2021/01/08. <https://doi.org/10.1038/s41423-020-00588-2> PMID: 33408342; PubMed Central PMCID: PMC7786875.
38. Roltgen K, Powell AE, Wirz OF, Stevens BA, Hogan CA, Najeeb J, et al. Defining the features and duration of antibody responses to SARS-CoV-2 infection associated with disease severity and outcome. *Sci Immunol*. 2020; 5(54). Epub 2020/12/09. <https://doi.org/10.1126/sciimmunol.abe0240> PMID: 33288645; PubMed Central PMCID: PMC7857392.
39. Klinger AS. Targeting Zero Infections in Dialysis: New Devices, Yes, but also Guidelines, Checklists, and a Culture of Safety. *J Am Soc Nephrol*. 2018; 29(4):1083–4. Epub 2018/03/07. <https://doi.org/10.1681/ASN.2018020132> PMID: 29507046; PubMed Central PMCID: PMC5875966.
40. Windpessl M, Bruchfeld A, Anders HJ, Kramer H, Waldman M, Renia L, et al. COVID-19 vaccines and kidney disease. *Nat Rev Nephrol*. 2021; 17(5):291–3. Epub 2021/02/10. <https://doi.org/10.1038/s41581-021-00406-6> PMID: 33558753; PubMed Central PMCID: PMC7869766.
41. Schrezenmeier E, Bergfeld L, Hillus D, Lippert JD, Weber U, Tober-Lau P, et al. Immunogenicity of COVID-19 Tozinameran Vaccination in Patients on Chronic Dialysis. *Front Immunol*. 2021; 12:690698. Epub 2021/07/20. <https://doi.org/10.3389/fimmu.2021.690698> PMID: 34276681; PubMed Central PMCID: PMC8284337.
42. Stumpf J, Siepman T, Lindner T, Karger C, Schwobel J, Anders L, et al. Humoral and cellular immunity to SARS-CoV-2 vaccination in renal transplant versus dialysis patients: A prospective, multicenter observational study using mRNA-1273 or BNT162b2 mRNA vaccine. *Lancet Reg Health Eur*. 2021:100178. Epub 2021/07/29. <https://doi.org/10.1016/j.lanpep.2021.100178> PMID: 34318288; PubMed Central PMCID: PMC8299287.

43. Lacson E, Argyropoulos C, Manley H, Aweh G, Chin A, Salman L, et al. Immunogenicity of SARS-CoV-2 Vaccine in Dialysis. *J Am Soc Nephrol*. 2021. Epub 2021/08/06. <https://doi.org/10.1681/ASN.2021040432> PMID: 34348908.
44. Anderegg MA, Liu M, Saganas C, Montani M, Vogt B, Huynh-Do U, et al. De novo vasculitis after mRNA-1273 (Moderna) vaccination. *Kidney Int*. 2021; 100(2):474–6. Epub 2021/06/05. <https://doi.org/10.1016/j.kint.2021.05.016> PMID: 34087251; PubMed Central PMCID: PMC8166777.
45. Khoury DS, Cromer D, Reynaldi A, Schlub TE, Wheatley AK, Juno JA, et al. Neutralizing antibody levels are highly predictive of immune protection from symptomatic SARS-CoV-2 infection. *Nat Med*. 2021. Epub 2021/05/19. <https://doi.org/10.1038/s41591-021-01377-8> PMID: 34002089.
46. Speer C, Benning L, Tollner M, Nusschag C, Kalble F, Reichel P, et al. Neutralizing antibody response against variants of concern after vaccination of dialysis patients with BNT162b2. *Kidney Int*. 2021; 100(3):700–2. Epub 2021/07/16. <https://doi.org/10.1016/j.kint.2021.07.002> PMID: 34265359; PubMed Central PMCID: PMC8274271.
47. Cohn BA, Cirillo PM, Murphy CC, Krigbaum NY, Wallace AW. SARS-CoV-2 vaccine protection and deaths among US veterans during 2021. *Science*. 2021:eabm0620. Epub 2021/11/04. <https://doi.org/10.1126/science.abm0620> PMID: 34735261.
48. Hazell SL. Clinical utility of avidity assays. *Expert Opin Med Diagn*. 2007; 1(4):511–9. Epub 2007/12/01. <https://doi.org/10.1517/17530059.1.4.511> PMID: 23496357.
49. Gaspar EB, De Gaspari E. Avidity assay to test functionality of anti-SARS-Cov-2 antibodies. *Vaccine*. 2021; 39(10):1473–5. Epub 2021/02/15. <https://doi.org/10.1016/j.vaccine.2021.02.003> PMID: 33581919; PubMed Central PMCID: PMC7857056.
50. Bauer G, Struck F, Schreiner P, Staschik E, Soutschek E, Motz M. The challenge of avidity determination in SARS-CoV-2 serology. *J Med Virol*. 2021; 93(5):3092–104. Epub 2021/02/11. <https://doi.org/10.1002/jmv.26863> PMID: 33565617; PubMed Central PMCID: PMC8013859.
51. Teimouri A, Mohtasebi S, Kazemirad E, Keshavarz H. Role of *Toxoplasma gondii* IgG Avidity Testing in Discriminating between Acute and Chronic Toxoplasmosis in Pregnancy. *J Clin Microbiol*. 2020; 58(9). Epub 2020/04/24. <https://doi.org/10.1128/JCM.00505-20> PMID: 32321784; PubMed Central PMCID: PMC7448626.
52. Genco F, Sarasini A, Parea M, Prestia M, Scudeller L, Meroni V. Comparison of the LIAISON(R)XL and ARCHITECT IgG, IgM, and IgG avidity assays for the diagnosis of *Toxoplasma*, cytomegalovirus, and rubella virus infections. *New Microbiol*. 2019; 42(2):88–93. Epub 2019/04/18. PMID: 30994178.
53. Chan L, Fuca N, Zeldis E, Campbell KN, Shaikh A. Antibody Response to mRNA-1273 SARS-CoV-2 Vaccine in Hemodialysis Patients with and without Prior COVID-19. *Clin J Am Soc Nephrol*. 2021; 16(8):1258–60. Epub 2021/05/26. <https://doi.org/10.2215/CJN.04080321> PMID: 34031182.
54. Carr EJ, Wu M, Harvey R, Wall EC, Kelly G, Hussain S, et al. Neutralising antibodies after COVID-19 vaccination in UK haemodialysis patients. *The Lancet*. 2021. [https://doi.org/10.1016/S0140-6736\(21\)01854-7](https://doi.org/10.1016/S0140-6736(21)01854-7) PMID: 34391504
55. Chapin-Bardales J, Gee J, Myers T. Reactogenicity Following Receipt of mRNA-Based COVID-19 Vaccines. *JAMA*. 2021; 325(21):2201–2. Epub 2021/04/06. <https://doi.org/10.1001/jama.2021.5374> PMID: 33818592.
56. Puranik A, Lenehan PJ, Silvert E, Niesen MJM, Corchado-Garcia J, O'Horo JC, et al. Comparison of two highly-effective mRNA vaccines for COVID-19 during periods of Alpha and Delta variant prevalence. *medRxiv*. 2021. Epub 2021/08/18. <https://doi.org/10.1101/2021.08.06.21261707> PMID: 34401884.
57. Abe KT, Hu Q, Mozafarihashjin M, Samson R, Manguiat K, Robinson A, et al. Neutralizing antibody responses to SARS-CoV-2 variants in vaccinated Ontario long-term care home residents and workers. *medRxiv*. 2021. <https://doi.org/10.1101/2021.08.06.21261721v2>
58. Steensels D, Pierlet N, Penders J, Mesotten D, Heylen L. Comparison of SARS-CoV-2 Antibody Response Following Vaccination With BNT162b2 and mRNA-1273. *JAMA*. 2021; 326(15):1533–5. Epub 2021/08/31. <https://doi.org/10.1001/jama.2021.15125> PMID: 34459863; PubMed Central PMCID: PMC8406205.
59. Richards NE, Keshavarz B, Workman LJ, Nelson MR, Platts-Mills TAE, Wilson JM. Comparison of SARS-CoV-2 Antibody Response by Age Among Recipients of the BNT162b2 vs the mRNA-1273 Vaccine. *JAMA Netw Open*. 2021; 4(9):e2124331. Epub 2021/09/03. <https://doi.org/10.1001/jamanetworkopen.2021.24331> PMID: 34473262; PubMed Central PMCID: PMC8414189.
60. Jahn M, Korth J, Dorsch O, Anastasiou OE, Sorge-Hadicke B, Tyczynski B, et al. Humoral Response to SARS-CoV-2-Vaccination with BNT162b2 (Pfizer-BioNTech) in Patients on Hemodialysis. *Vaccines (Basel)*. 2021; 9(4). Epub 2021/05/01. <https://doi.org/10.3390/vaccines9040360> PMID: 33918085; PubMed Central PMCID: PMC8070660.

61. Clarke CL, Predecki M, Dhutia A, Gan J, Edwards C, Prout V, et al. Longevity of SARS-CoV-2 immune responses in hemodialysis patients and protection against reinfection. *Kidney Int.* 2021; 99(6):1470–7. Epub 2021/03/29. <https://doi.org/10.1016/j.kint.2021.03.009> PMID: 33774082; PubMed Central PMCID: PMC7992297.
62. Longlune N, Nogier MB, Miedouge M, Gabilan C, Cartou C, Seigneuric B, et al. High immunogenicity of a messenger RNA based vaccine against SARS-CoV-2 in chronic dialysis patients. *Nephrol Dial Transplant.* 2021. Epub 2021/06/01. <https://doi.org/10.1093/ndt/gfab193> PMID: 34057463; PubMed Central PMCID: PMC8195197.
63. Ducloux D, Colladant M, Chabannes M, Yannarakis M, Courivaud C. Humoral response after 3 doses of the BNT162b2 mRNA COVID-19 vaccine in patients on hemodialysis. *Kidney Int.* 2021; 100(3):702–4. Epub 2021/07/04. <https://doi.org/10.1016/j.kint.2021.06.025> PMID: 34216675; PubMed Central PMCID: PMC8243640.
64. Espi M, Charmetant X, Barba T, Mathieu C, Pelletier C, Koppe L, et al. A prospective observational study for justification, safety, and efficacy of a third dose of mRNA vaccine in patients receiving maintenance hemodialysis. *Kidney Int.* 2021. Epub <https://doi.org/10.1016/j.kint.2021.10.040> PMID: 34856313; PubMed Central PMCID: PMC8628628.
65. Dekervel M, Henry N, Torreggiani M, Pouteau LM, Imiela JP, Mellaza C, et al. Humoral response to a third injection of BNT162b2 vaccine in patients on maintenance haemodialysis. *Clin Kidney J.* 2021; 14(11):2349–55. Epub 20210813. <https://doi.org/10.1093/ckj/sfab152> PMID: 34754430; PubMed Central PMCID: PMC8573007.
66. Massa F, Cremoni M, Gerard A, Grabsi H, Rogier L, Blois M, et al. Safety and cross-variant immunogenicity of a three-dose COVID-19 mRNA vaccine regimen in kidney transplant recipients. *EBioMedicine.* 2021; 73:103679. Epub 20211108. <https://doi.org/10.1016/j.ebiom.2021.103679> PMID: 34763205; PubMed Central PMCID: PMC8573385.
67. Robert T, Lano G, Giot M, Fourie T, de Lamballeri X, Jehel O, et al. Humoral response after SARS-COV2 vaccination in patient undergoing maintenance hemodialysis: loss of immunity, third dose and non-responders. *Nephrol Dial Transplant.* 2021. Epub 20211013. <https://doi.org/10.1093/ndt/gfab299> PMID: 34643714.