

STUDY PROTOCOL

The co-occurrence of the SAVA syndemic, depression and anxiety as barriers to antiretroviral therapy adherence among sub-Saharan Africa population groups: A scoping review protocol

Anton Delport^{1*}, Hanani Tabana¹, Lucia Knight^{1,2}, Edwin Wouters³

1 School of Public Health, University of the Western Cape, Bellville, South Africa, **2** Division of Social and Behavioural Sciences, University of Cape Town, Cape Town, South Africa, **3** Department of Sociology, University of Antwerp, Antwerpen, Belgium

✉ These authors contributed equally to this work.

* adelport@uwc.ac.za



OPEN ACCESS

Citation: Delport A, Tabana H, Knight L, Wouters E (2022) The co-occurrence of the SAVA syndemic, depression and anxiety as barriers to antiretroviral therapy adherence among sub-Saharan Africa population groups: A scoping review protocol. PLoS ONE 17(9): e0274614. <https://doi.org/10.1371/journal.pone.0274614>

Editor: Emily Chenette, PLOS (Public Library of Science), UNITED KINGDOM

Received: December 13, 2021

Accepted: September 1, 2022

Published: September 20, 2022

Copyright: © 2022 Delport et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: No datasets were generated or analysed during the current study. All relevant data from this study will be made available upon study completion. This study will make use of secondary data that are available to the public and does not require additional ethical approval.

Funding: Mr. Anton Delport received a PhD studentship through the Flemish Interuniversity Council (VLIR), as well as a partial scholarship through the University of the Western Cape, School

Abstract

Introduction

The scale-up of access to antiretroviral therapy has transformed HIV from an acute, terminal disease to a manageable chronic illness. Yet, sustaining high levels of antiretroviral therapy adherence remain a challenge, especially in the sub-Saharan Africa region which is disproportionately affected by HIV. This protocol proposes a scoping review to explore literature reporting on the antiretroviral therapy adherence levels among people who experience substance abuse and violence (SAVA) syndemics, as well as mood disorders such as anxiety and depression among people living with HIV in sub-Saharan Africa.

Methods and analysis

This proposed scoping review will follow Arksey and O'Malley's methodological framework for conducting scoping reviews as refined by Levac *et al.* The review will follow the Joanna Briggs Institute's manual for conducting scoping reviews. Literature searches will be conducted using six databases: Academic search complete; APA PsycArticles; CINAHL; MEDLINE; SocINDEX and Web of science. Title screening will see the "Participant, Concept, Context" framework applied to identify relevant literature and will not include the appraisal of search results. Data charting will follow an adapted version of Trico and colleagues' PRISMA-ScR and results will be mapped descriptively and in tabular format. Furthermore, results will be discussed within the syndemics model of health, and summarised as a biosocial conceptual model.

Ethics and dissemination

The study will make use of secondary data that are readily available to the public and will not require ethical approval. We intend to publish our results in a peer-reviewed journal and disseminate our findings at relevant conferences and seminars.

of Public Health (SoPH), linked to a grant received by SoPH from the Belgian Development Cooperation, through the Institute of Tropical Medicine Antwerp. Grant number: #ZA2018TEA474A102 (VLIR); #FWO G035018N

Competing interests: The authors have declared that no competing interests exist.

Introduction

The human immunodeficiency virus (HIV) remains a major global health epidemic with an estimated 38 million people living with HIV (PLWH), 1.7 million of whom were newly infected in 2020 [1]. Sub-Saharan Africa (SSA) is disproportionately affected by HIV as the region currently holds almost two thirds of the global HIV burden [2] and more than 60% of new infections [3]. The scale-up of access to antiretroviral therapy (ART) have transformed HIV from an acute, terminal disease to a manageable chronic illness [4, 5]. Rigid ART adherence behaviour has been shown to significantly decrease viral loads in PLWH to undetectable levels, effectively eliminating the risk of transmission [6]. Furthermore, good adherence to ART, meaning, taking antiretroviral medication as prescribed by a health practitioner, prevents immunological damage and improves life expectancy [7, 8].

It is for these reasons that it is essential to reach and sustain high levels of good ART adherence, however, it remains challenging for SSA—one of the poorest regions in the world [9]. There are social challenges which potentially not only induce the spread of HIV, but also hamper ART adherence. For example, the robust link between poverty and violence [10] and sexual and gender-based violence (SGBV) places women and young girls in particular at risk [11]. Studies have found that indirect exposure to violence [12] or being the recipient of violence [13] could significantly hinder an individual's capacity to adhere to ART [14].

Violence is often accompanied by substance use [15] with a particular high prevalence among women [16]. The literature highlights the intersecting, co-occurring and mutually reinforcing factors of substance use and violence as determinants of HIV risk [17] and non-adherence behaviour [18]. This distinguished phenomenon is known as the SAVA syndemic, a mnemonic that stands for Substance Abuse, Violence and HIV/AIDS [19]. This syndemic model closely considers the details of biosocial factors and how these through their interaction perpetuate adverse health outcomes [19] and impede sustainable ART success.

When studying the link between the SAVA syndemic and ART adherence, an important correlate linked to ART adherence factors come into play; namely mood disorders. Mood disorders, in terms of multimorbidity of disease, often occur as an outcome of-, or as a pretext to violence [19] and/or substance use [20]. This relationship of adverse health outcomes has been well documented in the literature; with depression and anxiety being the most common mood disorders associated with violence [21] and substance use [22]. Furthermore, depression and anxiety are typical mood disorders found among PLWH, and are considered to be key barriers to ART adherence and hamper positive, or 'good' adherence behaviour [23]. Thus, it can be argued that when assessing the impact of the SAVA syndemic on ART adherence levels, the influence that stems from co-occurring mood disorders, and in particular depression and anxiety, should also be accounted for.

Study rationale

Health problems rarely exist independently, and exploring the interplay between health problems, how they cluster and mutually impact ART adherence behaviour, is crucial in understanding and managing the spread of HIV. It is important to recognize that the SAVA syndemic is further confounded by mood disorders which typically manifest as anxiety and depression. The presence of both the SAVA syndemic and mood disorders have been found to be independent- and co-occurring as mutually re-enforcing barriers to ART adherence. Contemporary literature fails to adequately address the existence of mood disorders as co-occurring health burdens, and should be included when assessing the impact of the SAVA syndemic on ART adherence behaviour. This scoping review will attempt to bridge this gap by exploring

the most recent literature around this topic and offer new insights into the biosocial factors that reinforce poor ART adherence levels.

Theoretical underpinning of the study

The intended study aims to investigate secondary data around adverse biosocial factors common among PLWH, and how these factors impact individual capacity to manage HIV disease progression. Because of the aforementioned, we will discuss our results through the syndemics model of health [19]. This theoretical lens regards adverse biosocial factors as interacting, mutually occurring and reinforcing diseases; which flourishes under negative social and environmental factors [19]. The theory considers individual epidemics as sustained within communities due to harmful social connections, interactions and/or conditions [24]. It accounts for multiple interrelated systems that may contribute [24] to PLWHs poor levels of adherence behaviour by unpacking the syndemic interactions of substance use, IPV and poor mental health.

Aims and objectives

The study aims to explore literature reporting on substance use, violence, anxiety and depression among PLWH in SSA. We intend to identify the characteristics of these factors and map the outcomes as thematic domains of mutually occurring health problems which may synergistically impede ART adherence behaviour among SSA populations. The following objectives will guide the study:

1. Map the prevalence of the SAVA syndemic among PLWH on ART in SSA
2. Map the coexistence of the SAVA syndemic and depression/anxiety among PLWH on ART in SSA
3. Explore impact of the SAVA syndemic on ART adherence in SSA
4. Explore the impact of the combination of SAVA and depression/anxiety on ART adherence in SSA

Methods

Scoping review methodology

The broad nature of our research question and the large volume of literature published around our topic identified a scoping review as a fitting methodology for conducting this review [25, 26]. The documentation process of systematic searching, screening and synthesis of literature will offer results that are trustworthy by using a method that is replicable [25, 26]. The study's design will be guided by Arksey and O'Malley's (2005) [25] methodological framework for conducting scoping reviews as refined by Levac *et al.* (2010) [27]. The Joanna Briggs Institute's (JBI, updated 2020) [28] manual for conducting scoping reviews will guide this review, whilst data charting will follow the Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) reporting checklist (see Fig 1) [29]. Presented below is a non-linear framework which consists of five iterative stages to guide this study. It should be noted that the framework has an optional sixth stage which speaks to consulting with stakeholders however, this stage will be excluded for the purpose of this study as we will not consult with any stakeholders [25].

Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) Checklist

SECTION	ITEM	PRISMA-ScR CHECKLIST ITEM	REPORTED ON PAGE #
TITLE			
Title	1	Identify the report as a scoping review.	
ABSTRACT			
Structured summary	2	Provide a structured summary that includes (as applicable): background, objectives, eligibility criteria, sources of evidence, charting methods, results, and conclusions that relate to the review questions and objectives.	
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known. Explain why the review questions/objectives lend themselves to a scoping review approach.	
Objectives	4	Provide an explicit statement of the questions and objectives being addressed with reference to their key elements (e.g., population or participants, concepts, and context) or other relevant key elements used to conceptualize the review questions and/or objectives.	
METHODS			
Protocol and registration	5	Indicate whether a review protocol exists; state if and where it can be accessed (e.g., a Web address); and if available, provide registration information, including the registration number.	
Eligibility criteria	6	Specify characteristics of the sources of evidence used as eligibility criteria (e.g., years considered, language, and publication status), and provide a rationale.	
Information sources*	7	Describe all information sources in the search (e.g., databases with dates of coverage and contact with authors to identify additional sources), as well as the date the most recent search was executed.	
Search	8	Present the full electronic search strategy for at least 1 database, including any limits used, such that it could be repeated.	
Selection of sources of evidence†	9	State the process for selecting sources of evidence (i.e., screening and eligibility) included in the scoping review.	
Data charting process‡	10	Describe the methods of charting data from the included sources of evidence (e.g., calibrated forms or forms that have been tested by the team before their use, and whether data charting was done independently or in duplicate) and any processes for obtaining and confirming data from investigators.	
Data items	11	List and define all variables for which data were sought and any assumptions and simplifications made.	
Critical appraisal of individual sources of evidence§	12	If done, provide a rationale for conducting a critical appraisal of included sources of evidence; describe the methods used and how this information was used in any data synthesis (if appropriate).	
Synthesis of results	13	Describe the methods of handling and summarizing the data that were charted.	

SECTION	ITEM	PRISMA-ScR CHECKLIST ITEM	REPORTED ON PAGE #
RESULTS			
Selection of sources of evidence	14	Give numbers of sources of evidence screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally using a flow diagram.	
Characteristics of sources of evidence	15	For each source of evidence, present characteristics for which data were charted and provide the citations.	
Critical appraisal within sources of evidence	16	If done, present data on critical appraisal of included sources of evidence (see item 12).	
Results of individual sources of evidence	17	For each included source of evidence, present the relevant data that were charted that relate to the review questions and objectives.	
Synthesis of results	18	Summarize and/or present the charting results as they relate to the review questions and objectives.	
DISCUSSION			
Summary of evidence	19	Summarize the main results (including an overview of concepts, themes, and types of evidence available), link to the review questions and objectives, and consider the relevance to key groups.	
Limitations	20	Discuss the limitations of the scoping review process.	
Conclusions	21	Provide a general interpretation of the results with respect to the review questions and objectives, as well as potential implications and/or next steps.	
FUNDING			
Funding	22	Describe sources of funding for the included sources of evidence, as well as sources of funding for the scoping review. Describe the role of the funders of the scoping review.	

JBI = Joanna Briggs Institute; PRISMA-ScR = Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews.
 * Where sources of evidence (see second footnote) are compiled from, such as bibliographic databases, social media platforms, and Web sites.
 † A more inclusive/heterogeneous term used to account for the different types of evidence or data sources (e.g., quantitative and/or qualitative research, expert opinion, and policy documents) that may be eligible in a scoping review as opposed to only studies. This is not to be confused with information sources (see first footnote).
 ‡ The frameworks by Arksey and O'Malley (6) and Levac and colleagues (7) and the JBI guidance (4, 5) refer to the process of data extraction in a scoping review as data charting.
 § The process of systematically examining research evidence to assess its validity, results, and relevance before using it to inform a decision. This term is used for items 12 and 19 instead of "risk of bias" (which is more applicable to systematic reviews of interventions) to include and acknowledge the various sources of evidence that may be used in a scoping review (e.g., quantitative and/or qualitative research, expert opinion, and policy document).

From: Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med*. 2018;169:467-473. doi:10.7326/M18-0850



Fig 1. PRISMA-ScR [29].

<https://doi.org/10.1371/journal.pone.0274614.g001>

Stage 1: Identifying the research question

As part of this study’s conceptual process, an informal literature search was conducted by the primary investigator (PI) to advise the development of the preliminary research question. The yield of the initial literature search was reviewed to generate a sense around the knowledge-gaps in the literature. Following the literature review, in accordance with the recommendations made by the JBI [28], we applied the “Population, Concept, and Context” (PCC) framework as a guide to frame our primary research question (see Table 1). It was important at this stage for the research team to define the key concepts and population [30]. We agreed on the following concepts as central to our study: *SAVA syndemic, violence, substance use, depression, anxiety, and ART adherence behaviour*.

Our population of interest is PLWH who reside in the SSA region. We define substance use as any use or abuse of harmful or hazardous psychoactive substances [31], this includes

Table 1. Population- concept-context framework [28].

Components	Definition	Study
Population	Important characteristics (including age and other qualifying factors)	People living with HIV who have initiated ART (no age restriction, includes at-risk populations)
Concepts	A clear definition of the key concepts (including the scope and breadth of the search)	ART adherence behaviour, violence, substance use, anxiety, depression, SAVA syndemic
Context	Broad (including cultural, geographic, gender, racial or population factors)	Sub-Saharan Africa (impoverished settings)

<https://doi.org/10.1371/journal.pone.0274614.t001>

alcohol, street drugs and prescription or over-the-counter drugs. It was agreed that violence was a wide-ranging factor with many facets, with violence conceptualised as inclusive of inter- and/or intrapersonal abuse i.e. interpersonal violence, sexual and gender-based violence (SGBV), community-based violence, emotional abuse, self-harm or any other vicarious experiences of violence or harm.

Similarly, poor mental health is a broad concept that includes multiple conditions and criteria. It has been found that depression is the most widespread mood disorder found among PLWH [23], and is often experienced in conjunction with substance use [32–34] and/or violence [35]. Anxiety was identified as an additional key mood disorder due to its regular co-occurrence with depression [36] and strong association with HIV [37], substance use [38] and violence, specifically domestic and SGBV [39]. Similar to Kumar and colleagues (2009) [40], we defined ART adherence as the accuracy to which a client's behaviour compares with their recommended healthcare regime as discussed and agreed upon with their health care provider. Sub-Saharan Africa was chosen as the context of our study due to its disproportionately high burden of HIV in relation to the rest of the world [41]. Our preliminary research question concluded as: “*What is the impact of concurrent occurring SAVA syndemics and mood disorders, manifested as anxiety and depression, on ART adherence behaviour among PLWH in the SSA region?*”

Although scoping reviews do not require their explicit outcomes to be stated [26], we intend to address our primary research question through answering a series of secondary research questions (see Table 2 for intended outcomes), namely: (i) *How prevalent is the SAVA syndemic among PLWH on ART in SSA?*; (ii) *How prevalent is the coexistence of the SAVA syndemic and depression and/or anxiety among PLWH on ART in SSA?*; (iii) *What is the impact of the SAVA syndemic on ART adherence in SSA?*; (iv) *What is the impact of depression and/or anxiety on ART?*

Stage 2: Identify relevant studies

The PI developed the search strategy which was refined in collaboration with the research team. Six contextually relevant electronic databases were identified for the study and will be accessed through the University of the Western Cape's library databases: *Academic search*

Table 2. Research questions and intended outcomes.

Primary research question	Primary outcomes
What is the impact of concurrent occurring SAVA syndemics and mood disorders, manifested as anxiety and depression, on ART adherence behaviour among PLWH in the SSA region?	A conceptual syndemics model to inform the conceptualisation around the co-occurrence of SAVA syndemics and poor mental health among PLWH who have initiated ART (-who reside in the SSA region).
Secondary research questions	Secondary outcomes
How prevalent is the SAVA syndemic among PLWH on ART in SSA?	Prevalence levels/descriptives of mutually occurring SAVA syndemics among PLWH who have initiated ART (-who reside in the SSA region).
How prevalent is the coexistence of the SAVA syndemic and depression and/or anxiety among PLWH on ART in SSA?	Prevalence levels/descriptives of mutually occurring SAVA syndemics and depression and/or anxiety among PLWH who have initiated ART (-who reside in the SSA region).
What is the impact of the SAVA syndemic on ART adherence in SSA?	Tabular mapping detailing the impact which SAVA syndemics has on ART adherence among PLWH who have initiated ART (-who reside in the SSA region).
What is the impact of depression and/or anxiety on ART adherence?	Tabular mapping detailing the impact of depression and/or anxiety on ART adherence among PLWH who have initiated ART (-who reside in the SSA region).

<https://doi.org/10.1371/journal.pone.0274614.t002>

complete (EbscoHost); *APA PsycArticles* (EbscoHost); *CINAHL* (EbscoHost); *MEDLINE* (EbscoHost); *SocINDEX* (EbscoHost); and *Web of Science*. Our Boolean search string was informed by our preliminary literature review—see [Table 3](#) for the selected databases, search field and Boolean search string.

However, we acknowledge that our search string may change as we continue to refine our literature search as recommended by Levac *et al.* (2006) [30]. This study will include qualitative, quantitative and/or mixed-methods peer-reviewed publications, with literature published only in Afrikaans, English, Dutch, and French due to the language limitation of the research team. Furthermore, age will not be considered a limiter as HIV, violence, substance use and mood disorders, nor the impact thereof, are limited by age. Only literature published from the year 2010 till 2022 will be included in the search, we opted for a time limit that fell within the last decade to include findings that coincide with the widespread roll-out of ART, the test-and-treat initiative, and pre- and postexposure prophylaxis (PEP). This timeframe was identified to limit the potential for research that may be dated due to the aforementioned medical advancements. In addition, reference lists of articles included for full-text review will be hand searched to maintain the breadth of this review's inclusivity [30]. All literature searches will be conducted by the research team.

Stage 3: Study selection

Adhering to the recommendations made by Levac *et al.* (2006) [30, 42], identifying articles will follow a transparent, iterative review process before finalising the inclusion and exclusion criteria. The team will meet regularly throughout the screening process to discuss the results and update the criteria should it be required. A refined literature search will be led by the PI and the yield of the search exported to Mendeley for removal of duplicate references. The reviewing process will consist of two phases: (1) title/abstract screening and (2) full text screening. In the first stage, the identified references will be sorted alphabetically by author and divided equally among the three reviewers, i.e. each reviewer will screen a third of the titles. Reference titles will be screened using the PCC mnemonic as a guideline, it should be noted that the inclusion criteria may be refined post-hoc [30]. Furthermore, in the event of a reference with an ambiguous title, the abstracts will be screened using the PCC framework [30, 42]. References that were excluded during phase one of Stage 3, will undergo a second round of review by an independent research assistant. The second phase of this stage involves the screening of the 'full-text' of the references that have been identified for inclusion during phase one. This includes reading the content of the identified references, and compare it to the scope of this

Table 3. Database, field and boolean search strings.

Databases	Field	Boolean combinations
<i>Academic search complete</i>	Title	HIV AND (violence OR abuse OR gender-based violence OR GBV OR intimate partner violence OR IPV AND depress* OR anxiety OR mood disorder*) OR
<i>APA PsycArticles</i>	Abstract	(substance* OR drug* OR addict* OR people who inject drugs OR PWID* OR
<i>CINAHL</i>	Title	SAVA AND depress* OR anxiety OR mood disorder*) AND (association OR
<i>MEDLINE</i>	Title	relationship) AND (antiretroviral OR antiretroviral therapy OR HAART AND
<i>SocINDEX</i>	Title	adhere*) AND (sub Saharan Africa OR SSA OR Angola OR Benin OR Botswana
<i>Web of science</i>	Title	OR Burkina Faso OR Burundi OR Cabo Verde OR Cameroon OR Central African
		Republic OR Chad OR Comoros OR Congo OR Cote d'Ivoire OR Equatorial
		Guinea OR Eritrea OR Ethiopia OR Gabon OR Gambia OR Ghana OR Guinea
		OR Guinea-Bissau OR Kenya OR Lesotho OR Liberia OR Madagascar OR Malawi
		OR Mali OR Mauritania OR Mauritius OR Mozambique OR Namibia OR Niger
		OR Nigeria OR Rwanda OR Sao Tome and Principe OR Senegal OR Seychelles
		OR Sierra Leone OR Somalia OR South Africa OR South Sudan OR Sudan OR
		Tanzania OR Togo OR Uganda OR Zambia OR Zimbabwe)

<https://doi.org/10.1371/journal.pone.0274614.t003>

study using a rubric conceptualised on the PCC approach and the PRISMA-ScR [30] reporting criteria [30, 42]. The literature elimination process will be captured and detailed using the PRISMA flowchart (Fig 2).

Stage 4: Charting the data

Articles that are included will have their data extracted using an adapted version of the JBI's evidence details, characteristics and results extraction tool to address the objectives of this study; and follow Tricco *et al's* (2018) 27-item PRISMA-ScR [30] reporting guideline to guide and chart the contextual factors that are relevant for this review. All customisations of the extraction instrument will strictly follow the guidelines set out by the JBI [28]. It is noteworthy to highlight that the extraction tool will be further refined following an iterative process meaning, at specific intervals the team will regroup and review the information captured by the instrument and reflect the relevance of the contents in relation to the study's aims and objectives [30].




Fig 2. PRISMA flowchart.

<https://doi.org/10.1371/journal.pone.0274614.g002>

Extraction of the data will be done by the research team who will pilot a data charting exercise using the adapted instrument to familiarise themselves with, and review the rigor of the instrument. For this exercise, two publications will be randomly selected from the list of identified articles, with each researcher independently charting these articles [30]. This exercise will ensure that there is a mutual understanding of the data charting process and that the process remain aligned with the purpose of this study [30]. In the event that the exercise does not offer conclusive consensus around extracting the data, the tool will be amended as per recommendation and the exercise repeated.

Stage 5: Collating, summarising and reporting the results

The study's final stage will follow the PRISMA-ScR as a guideline to report our findings, this will occur across a three-phased step-wise process as suggested by Levac *et al.* (2006) [30]. Although it is acceptable for scoping reviews to report only on simple frequency counts i.e. concepts, populations, and characteristics [28], this study will further provide results as a summary of coded categories i.e. barriers to ART adherence which will be determined using descriptive qualitative content analysis [28]. Qualitative and quantitative data, i.e. frequency counts, will be descriptively mapped and coded into thematic domains that aligns with substance use, violence, and poor mental health in relation to ART adherence and where noted, document the interaction between these factors. During the second step we will follow the PRISMA-ScR guideline to report our results and outcomes as identified in step one, and map these details accordingly [30]. In step three we intend to provide an overview of our results and present this in a tabular descriptive format, as well as reflect on potential research gaps as identified through our literature review [28]. Lastly, the findings will be contextualised within the syndemics model of health, and summarised as a biosocial conceptual model to objectify the impact of co-occurring SAVA syndemics and poor mental health on ART adherence among PLWH residing in the SSA region (see Table 2: intended outcomes).

Ethics and dissemination

This study will make use of secondary data that are available to the public and does not require additional ethics approval. We intend to publish our results in a peer-reviewed journal and disseminate our findings at appropriate conferences and seminars. We acknowledge that the limitations of this proposed scoping review are found within the parameters of our search criteria and database limiters including time span (2010–2022), language (Afrikaans, English, Dutch, and French only), exclusion of white or green papers, grey literature and any literature that may not have been identified within our selected databases.

Discussion

This scoping review marks the first phase towards developing a model that will represent the synergistic and mutually reinforcing impact substance use, violence and mood disorders have on ART adherence behaviour. We foresee that the results will illuminate a broader foundation from which to build our understanding regarding the clustering and impact of the synergistic co-occurrence of the SAVA syndemic and mood disorders; and how these adverse health clusters perpetuate poor ART adherence behaviour, specifically among SSA populations.

Acknowledgments

Open access: This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy,

redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

Author Contributions

Conceptualization: Anton Delport.

Methodology: Anton Delport.

Supervision: Hanani Tabana.

Writing – original draft: Anton Delport.

Writing – review & editing: Hanani Tabana, Lucia Knight, Edwin Wouters.

References

1. UNAIDS. Global HIV Statistics 2020. Ending the AIDS epidemic. 2020;(June):1–3.
2. Magadi MA. The disproportionate high risk of HIV infection among the urban poor in sub-Saharan Africa. *AIDS Behav.* 2013; 17(5):1645–54. <https://doi.org/10.1007/s10461-012-0217-y> PMID: 22660933
3. Chilaka VN, Konje JC. HIV in pregnancy—An update. *European Journal of Obstetrics and Gynecology and Reproductive Biology.* 2021; 256:484–91. <https://doi.org/10.1016/j.ejogrb.2020.11.034> PMID: 33246666
4. Chetty V, Hanass-Hancock J. A rehabilitation model as key to comprehensive care in the era of HIV as a chronic disease in South Africa. *AIDS Care—Psychological and Socio-Medical Aspects of AIDS/HIV.* 2016; 28:132–9. <https://doi.org/10.1080/09540121.2016.1146204> PMID: 27002771
5. Wouters E, Masquillier C, le Roux Booysen F. The Importance of the Family: A Longitudinal Study of the Predictors of Depression in HIV Patients in South Africa. *AIDS Behav.* 2016; 20(8):1591–602. <https://doi.org/10.1007/s10461-016-1294-0> PMID: 26781870
6. Patel RR, Curoe KA, Chan PA. Undetectable Equals Untransmittable: A Game Changer for HIV Prevention. *Clin Chem.* 2020; 66(3):406–7. <https://doi.org/10.1093/clinchem/hvz010> PMID: 32109297
7. Soriano V, Barreiro P, de Mendoza C. Long-acting antiretroviral therapy. *Nat Mater.* 2020; 19 (August):823–9. <https://doi.org/10.1038/s41563-020-0731-7> PMID: 32704135
8. Gueler A, Moser A, Calmy AI, Gunthard CA, Battegay M, Cavassini M, et al. Life expectancy in HIV-positive persons in Switzerland: matched comparison with general population. *AIDS.* 2017; 31:427–36. <https://doi.org/10.1097/QAD.0000000000001335> PMID: 27831953
9. Bicaba Z, Brixiová Z, Ncube M. No Title. *Journal of African Development.* 2017; 19(2):93–110.
10. Crutchfield D Robert; Wadsworth T. Poverty and Violence. In: J HW and H, editor. *International Handbook of Violence Research.* Kluwer Academic Publishers; 2003. p. 67–82.
11. Artz L, Klazinga L, Müller A. Sexual and gender-based violence and HIV in South Africa: An HIV facility-based study. *South African Medical Journal.* 2020; 110(5):377–81. <https://doi.org/10.7196/SAMJ.2020.v110i5.13942> PMID: 32657721
12. Kacanek D, Malee K, Mellins CA, Tassiopoulos K, Smith R, Grant M, et al. Exposure to Violence and Virologic and Immunological Outcomes Among Youth With Perinatal HIV in the Pediatric HIV/AIDS Cohort Study. *Journal of Adolescent Health.* 2016; 59(1):30–7. <https://doi.org/10.1016/j.jadohealth.2016.03.004> PMID: 27089837
13. Merrill KG, Campbell JC, Decker MR, McGready J, Burke VM, Mwansa JK, et al. Prevalence of physical and sexual violence and psychological abuse among adolescents and young adults living with HIV in Zambia. *PLoS One.* 2020; 15(6):1–17. <https://doi.org/10.1371/journal.pone.0235203> PMID: 32584889
14. Carter A, Roth EA, Ding E, Milloy MJ, Kestler M, Jabbari S, et al. Substance Use, Violence, and Antiretroviral Adherence: A Latent Class Analysis of Women Living with HIV in Canada. *AIDS Behav.* 2018; 22(3):971–85. <https://doi.org/10.1007/s10461-017-1863-x> PMID: 28733919
15. Hicks PL, Mulvey KP, Chander G, Fleishman JA, Josephs JS, Korthuis PT, et al. The impact of illicit drug use and substance abuse treatment on adherence to HAART. *AIDS Care—Psychological and Socio-Medical Aspects of AIDS/HIV.* 2007; 19(9):1134–40. <https://doi.org/10.1080/09540120701351888> PMID: 18058397

16. Nydegger LA, Dickson-Gomez J, Ko TK. Structural and syndemic barriers to PrEP adoption among Black women at high risk for HIV: a qualitative exploration. *Cult Health Sex*. 2020; 0(0):1–15. <https://doi.org/10.1080/13691058.2020.1720297> PMID: 32212993
17. Hatcher AM, Gibbs A, McBride RS, Rebombo D, Khumalo M, Christofides NJ. Gendered syndemic of intimate partner violence, alcohol misuse, and HIV risk among peri-urban, heterosexual men in South Africa. *Soc Sci Med*. 2019;(xxxx):112637. <https://doi.org/10.1016/j.socscimed.2019.112637> PMID: 31708236
18. Rogers BG, Lee JS, Steven A. Encyclopedia of AIDS. *Encyclopedia of AIDS*. 2016;(May 2020).
19. Singer M, Bulled N, Ostrach B, Mendenhall E. Syndemics 1 Syndemics and the biosocial conception of health [Internet]. Vol. 389, www.thelancet.com. 2017. Available from: www.thelancet.com
20. Steinkamp JM, Goldblatt N, Borodovsky JT, LaVertu A, Kronish IM, Marsch LA, et al. Technological interventions for medication adherence in adult mental health and substance use disorders: A systematic review. *JMIR Ment Health*. 2019 Mar 1; 6(3). <https://doi.org/10.2196/12493> PMID: 30860493
21. Chandan JS, Thomas T, Bradbury-Jones C, Russell R, Bandyopadhyay S, Nirantharakumar K, et al. Female survivors of intimate partner violence and risk of depression, anxiety and serious mental illness. *The British Journal of Psychiatry*. 2019;(June):1–6.
22. Fojo AT, Lesko CR, Calkins KL, Moore RD, Mary E, Hutton HE, et al. *HHS Public Access*. 2020; 23(3):580–91.
23. Nel A, Kagee A. Common mental health problems and antiretroviral therapy adherence. *AIDS Care—Psychological and Socio-Medical Aspects of AIDS/HIV*. 2011; 23(11):1360–5. <https://doi.org/10.1080/09540121.2011.565025> PMID: 22022846
24. Wilson PA, Nanin J, Amesty S, Wallace S, Cherenack EM, Fullilove R. Using Syndemic Theory to Understand Vulnerability to HIV Infection among Black and Latino Men in New York City. Vol. 91, *Journal of Urban Health*. Springer Science and Business Media Deutschland GmbH; 2014. p. 983–98. <https://doi.org/10.1007/s11524-014-9895-2> PMID: 25155096
25. Arksey H, O'Malley L. Scoping studies: Towards a methodological framework. *International Journal of Social Research Methodology: Theory and Practice*. 2005; 8(1):19–32.
26. Tricco AC, Lillie E, Zarin W, O'Brien K, Colquhoun H, Kastner M, et al. A scoping review on the conduct and reporting of scoping reviews. *BMC Med Res Methodol*. 2016; 16(1):1–10. <https://doi.org/10.1186/s12874-016-0116-4> PMID: 26857112
27. Levac Danielle; Colquhoun Heather; O'Brien KK. Scoping studies: advancing the methodology. *Implementation science*. 2010; 5(69):1–9. <https://doi.org/10.1186/1748-5908-5-69> PMID: 20854677
28. Aromataris E. MZ, (Eds.). *JBIC Manual for Evidence Synthesis*. [Internet]. JBI. 2020. Available from: <https://wiki.jbi.global/display/MANUAL/Chapter+1+3A+Scoping+reviews>
29. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA extension for scoping reviews (PRISMA-ScR): Checklist and explanation. *Ann Intern Med*. 2018; 169(7):467–73. <https://doi.org/10.7326/M18-0850> PMID: 30178033
30. Levac Danielle, Heather Colquhoun1 KKO. Scoping studies: advancing the methodology. *Representing and Intervening*. 2012;1–18.
31. World Health Organisation. Management of substance abuse [Internet]. Available from: https://www.who.int/substance_abuse/terminology/abuse/en/
32. Singer M, Bulled N, Ostrach B, Mendenhall E. Syndemics and the biosocial conception of health. *The Lancet* [Internet]. 2017; 389(10072):941–50. Available from: [https://doi.org/10.1016/S0140-6736\(17\)30003-X](https://doi.org/10.1016/S0140-6736(17)30003-X) PMID: 28271845
33. Thomas JL, Lewis JB, Martinez I, Cunningham SD, Siddique M, Tobin JN, et al. Associations between intimate partner violence profiles and mental health among low-income, urban pregnant adolescents. *BMC Pregnancy Childbirth*. 2019; 19(1):1–8.
34. Abraham HD, Fava M. Order of Onset of Substance Abuse and Depression in a Sample of Depressed Outpatients. *Compr Psychiatry*. 1999; 40(1):44–50. [https://doi.org/10.1016/s0010-440x\(99\)90076-7](https://doi.org/10.1016/s0010-440x(99)90076-7) PMID: 9924877
35. Fazel S, Wolf A, Chang Z, Larsson H, Goodwin GM, Lichtenstein P. Depression and violence: A Swedish population study. *Lancet Psychiatry*. 2015; 2(3):224–32. [https://doi.org/10.1016/S2215-0366\(14\)00128-X](https://doi.org/10.1016/S2215-0366(14)00128-X) PMID: 26236648
36. Koeter MWJ, Den Brink W Van. The relationship between depression and anxiety: Construction of a prototypical anxiety and depression scale. *Psychol Med*. 1992; 22(3):597–606. <https://doi.org/10.1017/s0033291700038058> PMID: 1410085
37. Conall O'Cleirigh Sarah E, Valentine, Pinkston Megan, Herman Debra, C. Andres Bedoya, Janna R. Gordon and SAS. The unique challenges facing HIV-infected patients who smoke cigarettes: HIV

viremia, ART adherence, engagement in HIV care, and concurrent substance use. *Physiol Behav.* 2015; 19(1):178–185.

38. Parcesepe AM, Bernard C, Agler R, Ross J, Yotebieng M, Bass J, et al. Mental health and HIV: research priorities related to the implementation and scale up of “treat all” in sub-Saharan Africa. *J Virus Erad.* 2018; 4(Suppl 2):16–25. PMID: [30515310](https://pubmed.ncbi.nlm.nih.gov/30515310/)
39. Ahmadzad-Asl M, Davoudi F, Zarei N, Mohammad-Sadeghi H, Rasoulian M. Domestic violence against women as a risk factor for depressive and anxiety disorders: findings from domestic violence household survey in Tehran, Iran. *Arch Womens Ment Health.* 2016; 19(5):861–9. <https://doi.org/10.1007/s00737-016-0626-4> PMID: [26984712](https://pubmed.ncbi.nlm.nih.gov/26984712/)
40. Kumar A, Jones DD, Hanna MA, Soediono B, Bartocci AC. From access to adherence: the challenges to antiretroviral treatment. *J Chem Inf Model.* 2009; 53(3):556–81.
41. Larson JS. The World Health Organization’s definition of health: Social versus spiritual health. *Soc Indic Res.* 1996; 38(2):181–92.
42. Colquhoun HL, Levac D, O’Brien KK, Straus S, Tricco AC, Perrier L, et al. Scoping reviews: Time for clarity in definition, methods, and reporting. *J Clin Epidemiol.* 2014; 67(12):1291–4. <https://doi.org/10.1016/j.jclinepi.2014.03.013> PMID: [25034198](https://pubmed.ncbi.nlm.nih.gov/25034198/)