

RESEARCH ARTICLE

An X-STRs analysis of the Iraqi Sorani Kurds

Balnd M. Albarzinji¹, Shams Hadi², Bahez Ismael¹, Ahmed Barqee¹, Abdullah Hadi², Hayder Lazim^{3*}

1 Kurdistan Institution for Strategic Studies and Scientific Research (KISSR), Sulaymaniyah, Iraq, **2** University of Central Lancashire Medical School, Preston, United Kingdom, **3** Faculty of Health, Social Care and Medicine (FHSCM), School of Medicine, Edge Hill University, Ormskirk, United Kingdom

* alazawihayder@yahoo.com, lazimh@edgehill.ac.uk

Abstract

A database for the Iraqi Sorani Kurds, specifically focused on the 12 X-short tandem repeat (STR) loci, has been developed to facilitate forensic and population genetics investigations. The present study involved genotyping 117 unrelated individuals from the Sorani Kurds ethnic group using the Investigator Argus X-12 QS kit. The analysis revealed that the DXS10135 locus exhibited the highest degree of polymorphism, as indicated by a polymorphism information content (PIC) value of 0.94565 and a gene diversity (GD) value of 0.95623. Conversely, the DXS8378 locus displayed the lowest level of polymorphism, with a PIC value of 0.61026 and a GD value of 0.68170. Notably, two individuals were found to possess a rare allele (allele = 6) at the DXS8378 locus, which was not included in the allelic ladder of the kit. Furthermore, a significant linkage disequilibrium (LD) ($p < 0.05/117$) was observed between the DXS10103 and DXS10101 loci on linkage group 3 (LG3). The ancestral composition of the five primary geographic regions, namely Africa, Middle East, East Asia, Europe, and South America, was determined through the utilization of the F_{ST}/F_{ST}^{max} ratio. The findings of this analysis revealed that the Middle Eastern populations exhibited the lowest F_{ST}/F_{ST}^{max} ratio, measuring at 0.23243, indicating a relatively lower ancestral diversity. Conversely, the European populations showcased the highest F_{ST}/F_{ST}^{max} ratio, measuring at 0.27122, indicative of a greater ancestral diversity within this region. Additionally, the allelic richness indicators, namely distinctive and private alleles, indicated that Africa and the Middle East displayed the highest levels, while Far East Asia exhibited the lowest. This analysis supports the hypothesis of repeated founder effects during outward migrations, as evidenced by both the ancestry variability and the allelic richness. Consequently, the findings of this study have important implications for forensic genetics and population genetics research, particularly in relation to the consideration of genetic predispositions within specific ethnic groups.

OPEN ACCESS

Citation: M. Albarzinji B, Hadi S, Ismael B, Barqee A, Hadi A, Lazim H (2023) An X-STRs analysis of the Iraqi Sorani Kurds. PLoS ONE 18(11): e0294973. <https://doi.org/10.1371/journal.pone.0294973>

Editor: Naji Arafat Mahat, Universiti Teknologi Malaysia - Main Campus Skudai: Universiti Teknologi Malaysia, MALAYSIA

Received: August 1, 2023

Accepted: November 10, 2023

Published: November 27, 2023

Copyright: © 2023 M. Albarzinji et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its [Supporting Information](#) files.

Funding: BMA, The funding was partially provided by Kurdistan Institution for Strategic Studies and Scientific Research (project number 11, 2022-2023). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. There was no additional external funding received for this study.

Introduction

In many mammal species, including humans, the X chromosome is one of the two chromosomes that determines sex. It is only found in one copy in males, sharing this characteristic with both the Y chromosome and the mitochondrial DNA (mtDNA) [1].

Competing interests: The authors have declared that no competing interests exist.

The estimated length of the X chromosome is 155 million base pairs (Mb), or about 5% of the estimated size of the human genome (3,200 Mb). The X chromosome of the male gender is predominantly transmitted to females as a single unit. In contrast, the two copies in females recombine, like autosomes, thus reorganising genetic variation in each generation and increasing haplotype diversity. The newly rearranged chromosome is then passed on to both male and female offspring [1–4].

In forensic genetics, Short Tandem Repeat (STR) markers on the X and Y chromosomes have been developed as valuable tools that can complement the analysis of autosomal STR and mtDNA [5]. The major advantage of X chromosome short tandem repeats (X-STRs) is the analysis of female traces in male backgrounds and complicated kinship deficiency cases [6]. Moreover, X chromosome markers are also used in population analyses to understand the genetic diversity of a population and consequently can be helpful in demographic studies such as migration [7]. One of the commercial kits that have been used to analyse X-STRs in many populations worldwide is the Investigator Argus X-12 QS kit, which comprises 12 loci organized into four linkage groups (LGs) [8].

The X chromosome is crucial in understanding the differences in population-genetic studies between male and female individuals. Several factors influence the genetic distances found between males and females, including breeding and death rates, polygamy, and patrilocality. Notably, an important aspect contributing to the gender differences is the uneven distribution of the X chromosome, with females harbouring it for twice as long as males. The enormous genetic information encoded inside the X chromosome has the potential to provide crucial insights that cannot be obtained by analysing either the Y chromosome or mtDNA. Despite having substantially more genetic information than the Y chromosome or mtDNA, the X chromosome gets little attention in population genetic studies [1, 3].

Iraq is a nation located in the Arabian Peninsula, with a population of approximately 40 million people. It shares borders with several countries, including the Arabian Gulf, Kuwait, Saudi Arabia, Jordan, Syria, Turkey, and Iran. The population of Iraq consists of five distinct ethnic groups, with the Arabs and Kurds being the predominant groups. The Kurds, who inhabit a mountainous region spanning Iraq, Iran, Syria, Turkey, and Armenia, have different dialects such as Kurmanji, Sorani, Kirmashani, Zazaki, and Gorani. However, there is limited published data available on the comprehensive diversity of the population in Iraq and the genetic diversity of the Kurdish population [9–11]. This study focuses on the Sorani group, which is part of the Central Kurdish group in the northeastern Iraqi province of Sulaymaniyah. The Sulaymaniyah province, with a population of 779,000, is geographically situated amidst Iran to the east and Iraqi Arabs to the south [9].

For the first time, a representative database of X-STRs loci for the central Sorani Kurdish population of northern Iraq was presented to the scientific community. This study also looked at potential demographic differences based on genetic distances and allele frequencies between single populations and continental regions. Further, this study included a global investigation of the X chromosome's ancestral variability, allelic richness, and various populations' genetic structures.

Materials and methods

Sampling and DNA extraction

Buccal swab samples were collected from 117 Sorani Kurd males aged 18 years old and above. All work in this manuscript was reviewed and approved following institutional ethical guidelines. The Ethics Committee of the Kurdistan Institution for Strategic Studies and Scientific Research/Department of Biology granted ethical approval on February 15, 2020, with a unique

reference number (KI-ET 20/February 2020). Informed consent forms were completed by all participants and authors had no access to information that could identify individual participants during or after data collection. The DNA was extracted using a PrimePrep Genomic DNA Extraction Kit from Blood (GeNet Bio-Korea) according to the manufacturer's protocol [12]. The purity and concentration of the DNA were determined by using an Eppendorf Bio-photometer Plus (Eppendorf-Germany) [13]. The data were then collected in September 2020 and made accessible for research purposes in January 2021.

X-STRs typing

A set of X-STRs genetic markers were amplified using Investigator Argus X-12 QS kit (Qiagen) [14]. Simultaneous amplification was conducted following the manufacturer's instructions. The markers are distributed along the X chromosome and located in four LGs: LG 1—DXS10148, DXS10135, and DXS8378; LG 2—DXS7132, DXS10079, and DXS10074; LG 3—DXS10103, HPRTB, and DXS10101; LG 4—DXS10146, DXS10134, and DXS7423. The amplified fragments were separated on an ABI 3500 Genetic Analyzer, and allele calling was performed with GeneMapper[®] V.5 software (Thermo Fisher Scientific).

Statistical analysis

Forensic and population genetic parameters. The haplotype frequencies were calculated using GenAIEx 6.5 [15, 16]. The forensics statistics including gene diversity (GD), polymorphism information content (PIC) and match probability (PM) were calculated using the STRAF online tool [17]. Power of discrimination (PD) assessment was conducted for both male and female individuals, alongside the determination of the mean exclusion chance (MEC). The calculation of MEC involved the utilization of various formulas, tailored to specific scenarios. For cases involving paternal deficiency, Krüger's formula was employed, allowing for the examination of the paternal grandmother as a substitute for the putative father [18]. Kishida's formula, on the other hand, was applied to trios with a daughter [19]. In instances involving duos comprising a daughter and a putative father, or a son and mother, as well as trios with daughters, Desmarais' formulas were employed [20]. The results from the aforementioned formulas were yielded using StatsX v1.1 [21]. A pairwise exact test of linkage disequilibrium (LD) was performed for all pairs of loci using STRAF online tool [22]. The population genetic structure in our data was evaluated by the analysis of molecular variance (AMOVA). Molecular data were obtained for the Iraqi Sorani Kurds population using X-STRs based on the Investigator Argus X-12 QS kit (Qiagen) and compared with the worldwide available population data. Arlequin 3.5.2.2 software [23, 24] was used to calculate the average pairwise differences between (PiXY) and within populations (PiX), in addition to the corrected average pairwise difference between populations $(PiXY - (PiX + PiY)/2)$. More specifically, genetic distances between groups of males were quantified by F_{ST} calculations based on X-STRs data and multi-dimensional scaling (MDS) plots. MDS analysis is also used to investigate genetic similarities between populations [25] and to visualize the variances of the genetic differences in X-STRs and between populations. The genetic matrices and the MDS plot were generated using R statistical software version 4.0. The phylogenetic tree was constructed from allele frequency data using POPTREE2 online tool [26] and FigTree software [27].

Structure statistical analysis. Population structure was investigated using the program STRUCTURE version 2.3.4 [28, 29] with an admixture model. STRUCTURE is a model-based inference of population structure program in which individuals are grouped into a series of statistical clusters based on their multilocus genotypes. Each person had a membership

coefficient linked to each cluster, which represented the proportion of ancestry that is attributable to the respective cluster.

This study looked into the structure of five geographical regions: 11 Middle Eastern countries and populations [30–35], three African regions [36–38], seven South Asian countries and populations [39–44], 17 European regions and populations [32, 34, 45–52] and eight South American regions and populations [53, 54]. STRUCTURE output was processed with STRUCTURE HARVESTER to analyse and display likelihood values across numerous values of K, as well as to determine the number of genetic groupings that best suit the data [55, 56]. Then, using CLUMPP program [57, 58], The multiple replicate analyses of each data set were aligned, and the resulting files were used to create a graph of population Q-matrices for the X-STRs data for 12 markers across 46 subpopulations using the Distruct program [59, 60].

This study also looked at the levels of variation in membership coefficients among individuals who belonged to two or more designated clusters. The FSTruct program was used to investigate membership variability differences between admixed and non-admixed populations [61, 62].

To better understand genetic diversity and population relationships, allele distributions were studied across five metapopulations: Middle East, Africa, Far East Asia, Europe and South America. We evaluated the number of distinct alleles in the population and the number of unique (private) alleles to the population, that are not present in other populations, as two fundamental characteristics for a population at a given locus. Both private and distinct alleles are especially useful when looking for highly variable multiallelic markers like microsatellites in populations. The number of distinct alleles and private alleles was counted using the Allelic Diversity Analyzer (ADZE) Version 1.0 software [63, 64].

Results

The Sorani Kurds X-STRs database

Haplotype frequencies for each LG were calculated. There were 105, 81, 79, and 107 haplotypes detected in LG1, LG2, LG3, and LG4, respectively. LG4 was the most informative, with a frequency of 0.00855 for the most common haplotype, whereas LG3 was the least informative in the Iraqi Sorani Kurd males and presented in S1 Table in [S1 File](#). A total number of 159 different alleles for all loci were observed, varying from 29 for DXS10135 to 5 for DXS8378 and DXS7423. Allele frequencies of the 12 X-STRs loci shown in S2 Table ([S1 File](#)) and S1 Fig ([S2 File](#)). Variant alleles were observed in different samples. An off-ladder allele was observed at locus DXS8378 (two samples), duplicated alleles were found at locus DXS7423 in one individual and a null allele was also found in one individual at locus DXS10079 (S2 Fig in [S2 File](#)).

The PIC and the GD showed that the most polymorphic marker in the Iraqi Sorani Kurds was DXS10135 (PIC = 0.94565, GD = 0.95623). This marker had a total of 29 alleles, of which allele 24 was the most frequent (0.11111). While the least polymorphic marker was DXS8378 (PIC = 0.61026, GD = 0.68170), which had a total of 5 alleles, of which allele 11 was the most frequent (0.37607). The PD of males and females, along with different MEC computations, followed the same pattern as the GD and PIC, with the highest values observed in DXS10135 and the lowest values observed in DXS8378. The alleles frequencies and forensic parameters of Iraqi central Sorani Kurd males are shown in S2 and S3 Tables in [S1 File](#). The MEC estimates for the 12 loci in the Iraqi Sorani Kurds ethnic group were all shown in S3 Table in [S1 File](#), including MEC_Kruger, MEC_Kishida, MEC_Desmarais, and MEC_Desmarais_duo.

Linkage disequilibrium analysis

The linkage disequilibrium results for the 66 pairwise comparisons of the Iraqi Sorani Kurdish population are provided in S4 Table in [S1 File](#). After applying Bonferroni's correction, the data

revealed just one significant LD ($p < 0.05/117$), which was between the DXS10103 and DXS10101 locus pairs from LG 3 (S3 Fig in S2 File). There was no statistically significant LD between pairings of loci from different LGs.

Single locus analysis in different populations

A low level of genetic diversity was observed in our study at all 12 X-STRs of the Investigator Argus X-12 QS kit (Qiagen) panel (S4 Fig in S2 File). The locus DXS10135 had the highest gene diversity of 0.94185, while the locus DXS8378 had the lowest gene diversity of 0.68004. Some 406 different alleles were observed in the 6,634 X chromosomes analyzed, with a median number of 32 alleles per marker and a range of 12 (DXS8378) to 61 (DXS10135) as shown in S5 Table (S1 File).

Populations structure

A pairwise matrix plot of F_{ST} distances was generated to compare the 36 populations using 12 loci (Fig 1 and S6 Table in S1 File). The UAE ($F_{ST} = 0.00085$) had the closest population to Sorani Kurds, while the Philippines ($F_{ST} = 0.01053$) had the farthest. In the 36 populations studied, the closest association was identified between Jews [Chuetas] and Jews [Majorca] ($F_{ST} =$




Fig 1. The matrix of pairwise genetic distance F_{ST} of X-STRs 36 populations based on 12 X-STRs markers. This matrix was generated using Arlequin 3.5.2.2 software.

<https://doi.org/10.1371/journal.pone.0294973.g001>

-0.0059), and the farthest relationship was identified between Turkey and Lithuania ($F_{ST} = 0.0197$).

The average pairwise differences were examined to estimate the corrected genetic differences among 36 worldwide populations. These differences were specifically analysed in three contexts: between the 36 populations as a whole, within each individual population, and between different populations using Nei's distance.

The investigation revealed notable results regarding the average pairwise values obtained by using Nei's distance. The population of Turkey and the population of Lithuania had the highest average pairwise value, 0.18653. The lowest average pairwise value, -0.05476, was discovered between the Jewish subgroups of Chuetas and Majorca. Furthermore, the Sorani Kurd population had unique average pairwise values. The highest average pairwise value, 0.09900, was found between the Sorani Kurds and the Philippines. The Sorani Kurds and the United Arab Emirates, on the other side, had the lowest average pairwise value (0.00789).

The Iranian population had the highest average pairwise difference value within populations (9.44449), while Jews [Sephardim] had the lowest (9.07782). Mongolia and Jews [Chuetas] had the highest corrected average pairwise value (9.47319), while Hungary and Jews [Sephardim] had the lowest (9.09849). The Sorani Kurds had the highest corrected average pairwise value (9.43045) with Iran and the lowest (9.22487) with Jews [Sephardim]. The results are shown in S7 Table (S1 File) and Fig 2.

The MDS was calculated based on F_{ST} genetic distances of 12 X-STRs loci. The resulting MDS plot was generated using R software, depicting the relationships among samples from 36 global populations. The result is shown in Fig 3. The MDS revealed four main clusters, with the central cluster having the most populations (17), followed by clusters of 6 in the upper right quadrant, 8 in the lower right quadrant, and 4 in the lower left quadrant. Along with the populations of Saudi Arabia, the United Arab Emirates, and Iraq, Sorani Kurds are clustered in the lower left quadrant. The sole population that did not exhibit clustering was the population of Lithuania, which was isolated within the upper left quadrant.

To measure the diversity between populations a phylogenetic tree was constructed based on allele frequency data using POPTREE2 software [14]. FigTree software was subsequently used to visualise the phylogenetic tree. Five subpopulations ($K = 5$) were the optimal clustering for the 36 global populations (Fig 4). The Iraqi Sorani Kurds were assigned to the largest cluster (cluster 1). The inferred subpopulations were as follows: cluster 1: Sweden, Germany, Belarus, Portugal, UAE, Bahrain, Switzerland, Slovenia, Croatia, Hungary, Italy [Sardinia], Ethiopia [Tigray], Iran, Turkey, Saudi Arabia, Iraq [Sorani Kurd], Argentina, cluster 2: China [Hebei Han], Korean [Yanbian], Malaysia [Malay], Mongolia, Mexico, Bangladesh, China [Han], Philippines, cluster 3: Egypt, Guinea Bissau, cluster 4: Jews [North Africa] and Jews [Majorca], and cluster 5: Iraq [Iraqi], Albania. In addition to Lithuania, four Jewish populations, Chuetas, Ashkenazim, Middle Eastern, and Sephardim, did not demonstrate any clustering.

The continental pairwise matrix showed more differentiation between the Middle East and South America ($F_{ST} = 0.00257$) compared to the differentiation between the Middle East and Africa ($F_{ST} = 0.00206$), Far East Asia ($F_{ST} = 0.00203$) and Europe ($F_{ST} = 0.00125$), as shown in Fig 5A and S9 Table (S1 File). Pairwise F_{ST} and the average pairwise differences between and within populations in different continents, in addition to the corrected average pairwise difference between populations of the five different continents were calculated. Nei's distance results showed that the highest genetic distance was between the Middle East and South America (0.02383) while the lowest genetic distance was between the Middle East and Europe (0.01162). The highest average number of pairwise differences within the population was in Europe (9.27788) and the lowest was in Far East Asia (9.24154). The highest corrected average




Fig 2. Matrix plot showing population average pairwise differences based on 12 loci and between 36 populations. The area above the diagonal (green) shows the average number of pairwise differences between populations (Pi_{XY}); the diagonal (orange) shows the average number of pairwise differences within the population (Pi_X); and below the diagonal (blue) shows the corrected average pairwise difference $(Pi_{XY} - (Pi_X + Pi_Y) / 2)$. The scale of differences is shown on the right side of the matrix. This matrix was generated using Arlequin 3.5.2.2 software.

<https://doi.org/10.1371/journal.pone.0294973.g002>

pairwise difference was between Africa and Europe (9.30024) and the lowest was between Far East Asia and the Middle East (9.26607), as shown in Fig 5B and S9 Table in S1 File.

Population admixtures and ancestry variability

The structure of 46 ethnic groups and populations was investigated using the program STRUCTURE version 2.3.4 and an admixture model. The populations Q-matrices were created using X-STRs data from 12 markers across 46 populations (6,634 individuals). Six clusters have developed. The five geographical regions studied (Africa, the Middle East, South East Asia, Europe, and South America) did not reveal any distinct subpopulation genetic structures (Fig 6).

The FSTruct programme was used to analyse the F_{ST} / F_{ST}^{max} ratio of the membership coefficient variability differences between admixed and non-admixed populations provided by the STRUCTURE analysis to investigate ancestry variability in the five geographical regions.

The results showed that the F_{ST} / F_{ST}^{max} ratio was lowest in the Middle Eastern and African populations, 0.23243 and 0.23395 respectively, while it was the highest in both the European




Fig 3. Multidimensional scaling plot of the 36 populations. Population codes are as follows: ALB: Albania; ARG: Argentina; BLR: Belarus; BGD: Bangladesh; BHR: Bahrain; CHE: Switzerland; CHN [H]: China [Han]; CHN [HH]: China [Hebie Han]; DEU: Germany; EGY: Egypt; ETH: Ethiopia; GNB: Guinea Bissau; HRV: Croatia; HUN: Hungary; IRN: Iran; IQ: Iraq; IQ [Ku]: Iraq [Sorani Kurd]; ITA: Italy; Jew [Ash]: Jews [Ashkenazim]; Jew [Chu]: Jews [Chuetas]; Jew [Maj]: Jews [Majorca]; Jew [ME]: Jews [Middle East]; Jew [NA]: Jews [North Africa]; Jew [Seph]: Jews [Sephardim]; KOR: Korea; LTU: Lithuania; MEX: Mexico; MGN: Mongolia; MYS: Malaysia; PHL: Philippines; PRT: Portugal; SUA: Saudi Arabia; SVN: Slovenia; SWE: Sweden; TUR: Turkey; UAE: United Arab Emirates. The MDS fig was generated using R statistical software version 4.0.3.

<https://doi.org/10.1371/journal.pone.0294973.g003>




Fig 4. Phylogenetic tree of genetic relationships among the 36 global populations. This phylogenetic tree was generated using POPTREE2 software. Five clusters (K = 5) were created. Iraqi Sorani Kurds fell into the biggest cluster. The cyan-coloured cluster is the largest (17 populations), followed by the purple-coloured cluster (8 populations), and the remaining clusters each have two populations.

<https://doi.org/10.1371/journal.pone.0294973.g004>




Fig 5. Plots displaying the average number of pairwise differences within and between the five continents. These matrices were generated using Arlequin 3.5.2.2 software.

<https://doi.org/10.1371/journal.pone.0294973.g005>

and South American populations, 0.27122 and 0.26324 respectively, as shown in Fig 7 and S10 Table (S1 File).

Pairs of bootstrap distributions of F_{ST}/F_{ST}^{max} were significantly different ($p < 2e-16$) for four pairwise combinations and $P = 1.40E-08$ between the Middle Eastern and the African populations, Wilcoxon rank-sum test. For all stats and ratios, and all stats and matrices, the Kruskal-Wallis chi-squared showed that there are significant differences between the groups being compared in the study (3921, $df = 4$, p -value = $2.2e-16$).

Allelic richness in different population groups

This study examined two allelic richness parameters, specifically distinct and private alleles, across five distinct geographical regions. The findings revealed that the distinct alleles demonstrated similarities between Europe and South America, while displaying the highest levels in Africa and the Middle East, and the lowest levels in Far East Asia (Fig 8A, S11A Table (S1 File)). Moreover, the analysis revealed that Africa and the Middle East exhibited the highest levels of private alleles, while Europe and Far East Asia displayed the lowest levels (Fig 8B, S11B Table (S1 File)).

Discussion

The Argus X-12 (Qiagen, Hilden, Germany) is the most often used X-STRs kit in the field of forensic genetics. However, because the 12 X-STRs markers in this kit were largely characterised for European ancestry, several of the genetic variants observed in other population groups were missed out. To address this issue, the Argus X-12 QS emerged in the forensic field as an optimised version containing the same markers [1].

There has been few genetic research on the Iraqi Kurdish population, with the majority of previous work focusing on Y-STRs, mtDNA, and autosomal STRs [9, 10, 65, 66]. By analysing




Fig 6. Population structure analysis using investigator Argus X-12 STR markers. The results show 6 clusters ($K = 6$) using 46 ethnic groups and populations from five different geographical regions (Europe, South America, the Middle East, Africa and South East Asia).

<https://doi.org/10.1371/journal.pone.0294973.g006>

X chromosome markers, this research study hopes to contribute to a better knowledge of genetic diversity among the Kurdish population. Previous research in this field was limited, with only one study analysing the genetic makeup of the Kurdish population utilising only five X-STRs markers and treating the entire Kurdish population as a single group for comparison analysis [67]. In contrast, this study takes a broader approach, evaluating twelve X-STRs markers and focused specifically on a subgroup of the Kurdish community, the Iraqi Sorani Kurds.

The DXS10135 locus in the Kurdish population exhibited the highest polymorphism information content (PIC), which is comparable to Bahrain, UAE, Jews, Iraq, Lithuania, Turk,




Fig 7. Box plot, violin plot and empirical cumulative distribution function (ECDF) plot of the bootstrap distribution of F_{ST}/F_{ST}^{max} for each Q matrix in the STRUCTURE analysis.

<https://doi.org/10.1371/journal.pone.0294973.g007>

Slovenia, Albania, Guinea-Bissau, Filipino, Mongolia, Eastern Han, Hebei Han, Croatia, and Argentina [30–32, 34, 37, 40, 43, 44, 49, 54]. The lowest PIC was found at the locus DXS8378, which is shared by Argentina, Croatia, Turkey, Slovenia, and Albania, whereas the lowest PIC was found among other groups at the locus DXS7423. The bi-allele marker was observed in one male at the locus DXS7423; a doubled allele at the same locus was previously observed in the Iraqi population [34].

A rare allele was observed in two individuals, which is not presented in the allelic ladder of the kit at locus DXS8378 (allele = 6). This off-ladder allele at the same locus, DXS8378, was reported in the earlier study of the German population [47]. In addition, a null allele was also observed in one sample at the locus DXS10079 which, to our knowledge, has not been previously reported in other populations.

When MEC values were considered, the forensic efficacy of X-STRs in complex scenarios including kinship and paternity was obvious. These MEC calculations proved crucial in the examination of complex kinship and paternity scenarios, especially when the reference DNA sample came from the child's putative grandmother, the father's mother, rather than the alleged father.




Fig 8. The mean number of (A) distinct alleles per locus and (B) private alleles per locus of the five regions: Middle East, Africa, Far East Asia, South America and Europe.

<https://doi.org/10.1371/journal.pone.0294973.g008>

The phylogenetic tree showed that the Sorani Kurds, Saudi Arabia, Turkey, Iran, Ethiopia and Argentina fell into one cluster with the European populations. This study confirms the findings of a previous mtDNA study of the Sorani Kurds, which found that they are genetically related to the European lineage [65]. Furthermore, based on X-STRs, the Bahraini and Emirati populations were more closely linked, supporting the findings of Al-Snan's previous study [30].

Even after applying Bonferroni's correction, the linkage disequilibrium tests revealed one significant result between the DXS10103 and DXS10101 locus pairs from LG 3. A similar result with statistically substantial LD between DXS10103 and DXS10101 has previously been observed in the Iranian, Filipino, Mongolian, and Chinese Han populations [35, 40, 43, 44]. Another study found significant associations between the markers DXS10103 and DXS10101, as well as DXS10103 and HPRTB in the Chuetas Jewish population [32]. However, various population studies have found substantial levels of LD for various pairs of loci within different linkage groups [30, 31, 34, 37, 49, 54]. The observed variations are likely attributable to the sample size, since detecting linkage disequilibrium typically requires a substantial number of samples [68].

In comparison to the Y chromosome, which demonstrated a high level of gene diversity across all PPY23 kits in a global analysis [69], this study demonstrated a low degree of gene diversity across all 12 X-STRs in the Investigator Argus X-12 QS kit (Qiagen) panel.

Unlike the Y chromosome, the X chromosome population Q-matrices did not reveal a distinct subpopulation genetic structure [11]. In light of the findings, it can be concluded that population admixture did not exhibit significant relevance in the context of subpopulation clusters. Consequently, the determination of the specific geographical region to which the Sorani Kurds can be attributed within the five primary regions investigated in this study proved to be unfeasible. This could be explained by the fact that the X chromosome differs from the Y chromosome in that it undergoes recombination. The entirety of the Y chromosome functions as a single locus and has a single genealogical history. On the other hand, the X chromosome gets broken by recombination every generation, giving different regions on the chromosome different histories [3].

Individuals within populations that have recently undergone admixture differ substantially in terms of their ancestry, with some individuals deriving the majority of their ancestry from one source population and others from a different one. As a result, it might be hypothesised that admixed populations produce more ancestry variability than nonadmixed populations when assessing inferred cluster memberships. This was clearly shown in this study, as the Middle Eastern and African populations had the lowest ancestral variability as compared to European and South American populations.

Based on numerous viewpoints on the history of human migrations, we can evaluate patterns of distinct and private allelic richness in relation to our expectations. The higher number of distinct and private alleles in Africa corresponds to the pattern expected for models of human evolution that began in Africa and spread to other regions through a succession of founder events. The combination of the physically contiguous regions of Africa and the Middle East contains the greatest number of alleles. Since many alleles in the founding population would have only moved along part of their migration outside of Africa. The findings of this study on allelic richness and ancestry variability are consistent with the predictions of African-origin models that include serial founder effects throughout outward migrations [70].

The main limitation encountered in this research endeavour was the limited availability of X-STRs data. As seen, this study only included three African countries and two South American countries, necessitating additional data to enhance the understanding of the X-STRs population genetics.

In conclusion, this study represents the pioneering investigation into the X-STRs data pertaining to the population of Iraqi Sorani Kurds. The establishment of a comprehensive

population database has enabled the utilization of STR markers included in the Argus X-12 Kit in routine forensic casework. The remarkable statistical parameters exhibited by these loci, in conjunction with their independent inheritance patterns, render them an invaluable marker set for both forensic casework and population genetics research. The use of X-STRs loci may prove to be a significant complement to autosomal and Y-STRs markers, particularly in complicated kinship and paternity instances.

Supporting information

S1 File. S1 Table. Haplotype frequencies for 12 X-STRs markers in four linkage groups (LGs), N = 117. LG1 = (DXS8378, DXS10135, DXS10148), LG2 = (DXS7132, DXS10074, DXS10079), LG3 = (DXS10101, DXS10103, HPRTB), LG4 = (DXS7423, DXS10134, DXS10146). S2 Table: Allele frequencies and sample size by locus of the Sorani Kurd. S3 Table: Forensic parameters, Number of alleles (A_{all}), Gene diversity (GD), polymorphism information content (PIC), match probability (PM), power of discrimination (PD) and Mean Exclusion Chance (MEC) of the Sorani Kurd males. S4 Table: Pairwise p values of linkage disequilibrium test for all pairs of loci using 117 male samples from Sorani Kurdish population. S5 Table: Forensic parameters, Gene diversity (GD), polymorphism information content (PIC), match probability (PM) and power of discrimination (PD), of the five geographical regions Middle East, Africa, Far East Asia, Europe and South America. S6 Table: Pairwise genetic distances (F_{st}) based on haplotypes of 36 global populations. S7 Table: Average number of pairwise differences using data from 36 populations. Above diagonal: Average number of pairwise differences between populations (P_{iXY}), Diagonal elements (yellow highlighted): Average number of pairwise differences within population (P_{iX}), Below diagonal: Corrected average pairwise difference (P_{iXY}-(P_{iX}+P_{iY})/2). S8 Table: Pairwise genetic distances based on F_{ST} between the five continents. S9 Table: average number of pairwise differences (F_{ST}) within and between continents. Above diagonal: Average number of pairwise differences between populations (P_{iXY}), Diagonal elements (yellow highlighted): Average number of pairwise differences within population (P_{iX}), Below diagonal: Corrected average pairwise difference (P_{iXY}-(P_{iX}+P_{iY})/2). S10 Table: (A) Q statistics for the five populations Middle East, Africa, Far East Asia and South America. (B) Pairwise comparisons using Wilcoxon rank sum test with continuity correction. S11 Table: distinctive alleles and private alleles on a continental scale: Middle East, Africa, Far East Asia, South America and Europe. (A) Distinctive alleles, (B) Private alleles. (XLSX)

S2 File. S1 Fig. Plot distribution of allele frequencies per locus using Investigator Argus X-12 QS kit. S2 Fig. Electropherogram shows variant alleles in three samples of the Sorani Kurd males. (A) An off-ladder allele was observed at locus DXS8378 (allele = 6). (B) Duplicated alleles (14, 15) were found at locus DXS7423. (C) A null allele was observed at locus DXS10079. S3 Fig. P-values for the pairwise exact test matrix for linkage disequilibrium among the 12 X-STR loci. S4 Fig. Ranking of Investigator Argus X-12 QS kit (Qiagen) markers by gene diversity (GD). Rank within continental residency groups, i.e. Middle East (n = 1224), Africa (n = 448), Far East Asia (n = 815), South America (n = 1167) and Europe (n = 2980). (DOCX)

Acknowledgments

We are grateful to the blood donors for their cooperation and to everyone who helped us with this project. We wish to thank the two anonymous reviewers for their helpful comments on the manuscript.

Author Contributions

Conceptualization: Balnd M. Albarzinji, Shams Hadi, Bahez Ismael, Ahmed Barqee, Abdullah Hadi, Hayder Lazim.

Data curation: Balnd M. Albarzinji, Shams Hadi, Bahez Ismael, Ahmed Barqee, Abdullah Hadi, Hayder Lazim.

Formal analysis: Shams Hadi, Abdullah Hadi, Hayder Lazim.

Funding acquisition: Balnd M. Albarzinji.

Investigation: Balnd M. Albarzinji, Bahez Ismael, Ahmed Barqee.

Methodology: Balnd M. Albarzinji, Bahez Ismael, Ahmed Barqee, Hayder Lazim.

Software: Shams Hadi, Abdullah Hadi.

Supervision: Hayder Lazim.

Validation: Abdullah Hadi, Hayder Lazim.

Visualization: Hayder Lazim.

Writing – original draft: Balnd M. Albarzinji, Shams Hadi, Bahez Ismael, Ahmed Barqee, Abdullah Hadi, Hayder Lazim.

Writing – review & editing: Balnd M. Albarzinji, Shams Hadi, Bahez Ismael, Ahmed Barqee, Abdullah Hadi, Hayder Lazim.

References

1. Gomes I, Pinto N, Antão-Sousa S, Gomes V, Gusmão L, Amorim A. Twenty Years Later: A Comprehensive Review of the X Chromosome Use in Forensic Genetics. *Frontiers in Genetics*. 2020; 11. <https://doi.org/10.3389/fgene.2020.00926> PMID: 33093840
2. Ross MT, Grafham DV, Coffey AJ, Scherer S, McLay K, Muzny D, et al. The DNA sequence of the human X chromosome. *Nature*. 2005; 434(7031):325–37. <https://doi.org/10.1038/nature03440> PMID: 15772651
3. Schaffner SF. The X chromosome in population genetics. *Nature Reviews Genetics*. 2004; 5(1):43–51. <https://doi.org/10.1038/nrg1247> PMID: 14708015
4. Szibor R. X-chromosomal markers: past, present and future. *Forensic Sci Int Genet*. 2007; 1(2):93–9. Epub 20070427. <https://doi.org/10.1016/j.fsigen.2007.03.003> PMID: 19083736.
5. Diegoli TM. Forensic typing of short tandem repeat markers on the X and Y chromosomes. *Forensic Science International: Genetics*. 2015; 18:140–51. <https://doi.org/10.1016/j.fsigen.2015.03.013> PMID: 25934544
6. Tillmar AO, Kling D, Butler JM, Parson W, Prinz M, Schneider PM, et al. DNA Commission of the International Society for Forensic Genetics (ISFG): Guidelines on the use of X-STRs in kinship analysis. *Forensic Science International: Genetics*. 2017; 29:269–75. <https://doi.org/10.1016/j.fsigen.2017.05.005> PMID: 28544956
7. Garcia FM, Bessa BGO, dos Santos EVW, Pereira JDP, Alves LNR, Vianna LA, et al. Forensic Applications of Markers Present on the X Chromosome. *Genes*. 2022; 13(9):1597. <https://doi.org/10.3390/genes13091597> PMID: 36140765
8. Scherer M, König M, Bussmann M, Prochnow A, Peist R. Development and validation of the new Investigator® Argus X-12 QS Kit. *Forensic Science International: Genetics Supplement Series*. 2015; 5: e256–e7. <https://doi.org/10.1016/j.fsigs.2015.09.102>
9. Albarzinji BM, Abdulkarim FM, Hussein SA, Rashid D, Lazim H. Population genetic study of 17 Y-STR Loci of the Sorani Kurds in the Province of Sulaymaniyah, Iraq. *BMC Genomics*. 2022; 23(1):763. <https://doi.org/10.1186/s12864-022-09005-6> PMID: 36414939
10. Dogan S, Gurkan C, Dogan M, Balkaya HE, Tunc R, Demirdov DK, et al. A glimpse at the intricate mosaic of ethnicities from Mesopotamia: Paternal lineages of the Northern Iraqi Arabs, Kurds, Syrians, Turkmens and Yazidis. *PLoS One*. 2017; 12(11):e0187408. Epub 20171103. <https://doi.org/10.1371/journal.pone.0187408> PMID: 29099847; PubMed Central PMCID: PMC5669434.

11. Lazim H, Almohammed EK, Hadi S, Smith J. Population genetic diversity in an Iraqi population and gene flow across the Arabian Peninsula. *Scientific Reports*. 2020;10(1). <https://doi.org/10.1038/s41598-020-72283-1> PMID: 32943725
12. PrimePrep Genomic DNA Extraction Kit from Blood. Available from: http://genetbio.com/en/sub/sub01_0402.php.
13. BioPhotometer plus operating manual. Available from: <https://www.marshallscientific.com/v/vspfiles/specs/biophotometer%20plus%20manual.pdf>.
14. Investigator Argus X-12 QS Kit Quick-Start Protocol. Available from: <https://www.qiagen.com/us/resources/resourcedetail?id=c2b95c89-4895-4cfc-8716-79e334f14446&lang=en>.
15. GenAIEx. Available from: <https://biology-assets.anu.edu.au/GenAIEx/Welcome.html>.
16. Peakall R, Smouse PE. GenAIEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research—an update. *Bioinformatics*. 2012; 28(19):2537–9. Epub 20120720. <https://doi.org/10.1093/bioinformatics/bts460> PMID: 22820204; PubMed Central PMCID: PMC3463245.
17. Gouy A, Zieger M. STRAF-A convenient online tool for STR data evaluation in forensic genetics. *Forensic Sci Int Genet*. 2017; 30:148–51. Epub 20170715. <https://doi.org/10.1016/j.fsigen.2017.07.007> PMID: 28743032.
18. Krüger J, Fuhrmann W, Lichte KH, Steffens C. [On the utilization of erythrocyte acid phosphatase polymorphism in paternity evaluation]. *Dtsch Z Gesamte Gerichtl Med*. 1968; 64(2):127–46. PMID: 4974841.
19. Kishida T, Wang W, Fukuda M, Tamaki Y. Duplex PCR of the Y-27H39 and HPRT loci with reference to Japanese population data on the HPRT locus. *Nihon Hoigaku Zasshi*. 1997; 51(2):67–9. PMID: 9184015.
20. Desmarais D, Zhong Y, Chakraborty R, Perreault C, Busque L. Development of a highly polymorphic STR marker for identity testing purposes at the human androgen receptor gene (HUMARA). *J Forensic Sci*. 1998; 43(5):1046–9. PMID: 9729823.
21. StatsX Package. Available from: https://www.researchgate.net/publication/308791536_StatsX_Package.
22. STRAF online tool. Available from: <https://straf-p7bdrhm3xq-ew.a.run.app/>.
23. Arlequin ver 3.5.2.2. Available from: <http://cmapg.unibe.ch/software/arlequin35/>.
24. Excoffier L, Lischer HE. Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol Ecol Resour*. 2010; 10(3):564–7. Epub 20100301. <https://doi.org/10.1111/j.1755-0998.2010.02847.x> PMID: 21565059.
25. Kruskal JB. Multidimensional scaling by optimizing goodness of fit to a nonmetric hypothesis. *Psychometrika*. 1964; 29(1):1–27. <https://doi.org/10.1007/BF02289565>
26. Takezaki N, Nei M, Tamura K. POPTREE2: Software for Constructing Population Trees from Allele Frequency Data and Computing Other Population Statistics with Windows Interface. *Molecular Biology and Evolution*. 2009; 27(4):747–52. <https://doi.org/10.1093/molbev/msp312> PMID: 20022889
27. FigTree. Available from: <http://tree.bio.ed.ac.uk/software/figtree/>.
28. Structure Software. Available from: <https://web.stanford.edu/group/pritchardlab/structure.html>.
29. Pritchard JK, Stephens M, Donnelly P. Inference of population structure using multilocus genotype data. *Genetics*. 2000; 155(2):945–59. <https://doi.org/10.1093/genetics/155.2.945> PMID: 10835412; PubMed Central PMCID: PMC1461096.
30. Al-Snan NR, Messaoudi SA, Bakhiet M, Mansoor LA. Population genetic analysis of 12 X-STRs in a Bahraini population sample (Allele and haplotype frequencies of 12 X-STRs in Bahraini population). *bioRxiv*. 2019:790915. <https://doi.org/10.1101/790915>
31. Almarri MA, Lootah RA. Allelic and haplotype diversity of 12 X-STRs in the United Arab Emirates. *Forensic Sci Int Genet*. 2018; 33:e4–e6. Epub 20171229. <https://doi.org/10.1016/j.fsigen.2017.12.013> PMID: 29305242.
32. Ferragut JF, Bentayebi K, Pereira R, Castro JA, Amorim A, Ramon C, et al. Genetic portrait of Jewish populations based on three sets of X-chromosome markers: Indels, Alu insertions and STRs. *Forensic Sci Int Genet*. 2017; 31:e5–e11. Epub 20170918. <https://doi.org/10.1016/j.fsigen.2017.09.008> PMID: 28951006.
33. Messaoudi SA, Babu SR, Alsaleh AB, Albujja M, Al-Snan NR, Chaudhary AR, et al. Population genetic data for 12 X-STR loci in the Central Saudi region using investigator Argus X-12 amplification kit. *Ann Hum Biol*. 2021; 48(4):321–6. Epub 20210803. <https://doi.org/10.1080/03014460.2021.1957147> PMID: 34340612.
34. Poulsen L, Tomas C, Drobnič K, Ivanova V, Mogensen HS, Kondili A, et al. NGMSElect™ and Investigator® Argus X-12 analysis in population samples from Albania, Iraq, Lithuania, Slovenia, and Turkey.

- Forensic Science International: Genetics. 2016; 22:110–2. <https://doi.org/10.1016/j.fsigen.2016.02.004> PMID: 26894835
35. Poulsen L, Farzad MS, Børsting C, Tomas C, Pereira V, Morling N. Population and forensic data for three sets of forensic genetic markers in four ethnic groups from Iran: Persians, Lurs, Kurds and Azeris. *Forensic Science International: Genetics*. 2015; 17:43–6. <https://doi.org/10.1016/j.fsigen.2015.03.010> PMID: 25814339
 36. Elakkary S, Hoffmeister-Ullerich S, Schulze C, Seif E, Sheta A, Hering S, et al. Genetic polymorphisms of twelve X-STRs of the investigator Argus X-12 kit and additional six X-STR centromere region loci in an Egyptian population sample. *Forensic Sci Int Genet*. 2014; 11:26–30. Epub 20140224. <https://doi.org/10.1016/j.fsigen.2014.02.007> PMID: 24632058.
 37. Gomes I, Pereira PJP, Harms S, Oliveira AM, Schneider PM, Brehm A. Genetic characterization of Guinea-Bissau using a 12 X-chromosomal STR system: Inferences from a multiethnic population. *Forensic Sci Int Genet*. 2017; 31:89–94. Epub 20170824. <https://doi.org/10.1016/j.fsigen.2017.08.016> PMID: 28858674.
 38. Haddish K, Chierito E, Di Vella G, Lacerenza D, Raddi S, Aneli S, et al. A reference database of forensic autosomal and gonosomal STR markers in the Tigray population of Ethiopia. *Forensic Sci Int Genet*. 2022; 56:102618. Epub 20211023. <https://doi.org/10.1016/j.fsigen.2021.102618> PMID: 34735940.
 39. Li XN, Zheng JL, Yao J, Ding M, Pang H, Wang BJ. Population data of 12 X-chromosome STR loci in Yanbian Korean samples from China. *Forensic Sci Int Genet*. 2014; 13:143–4. Epub 20140729. <https://doi.org/10.1016/j.fsigen.2014.07.012> PMID: 25117403.
 40. Salvador JM, Apaga DLT, Delfin FC, Calacal GC, Dennis SE, De Ungria MCA. Filipino DNA variation at 12 X-chromosome short tandem repeat markers. *Forensic Sci Int Genet*. 2018; 36:e8–e12. Epub 20180608. <https://doi.org/10.1016/j.fsigen.2018.06.008> PMID: 29909139.
 41. Samejima M, Nakamura Y, Nambiar P, Minaguchi K. Genetic study of 12 X-STRs in Malay population living in and around Kuala Lumpur using Investigator Argus X-12 kit. *Int J Legal Med*. 2012; 126(4):677–83. Epub 20120515. <https://doi.org/10.1007/s00414-012-0705-7> PMID: 22584910.
 42. Sufian A, Hosen MI, Fatema K, Hossain T, Hasan MM, Mazumder AK, et al. Genetic diversity study on 12 X-STR loci of investigator® Argus X STR kit in Bangladeshi population. *Int J Legal Med*. 2017; 131(4):963–5. Epub 20161208. <https://doi.org/10.1007/s00414-016-1513-2> PMID: 27933412.
 43. Tao R, Zhang J, Bian Y, Dong R, Liu X, Jin C, et al. Investigation of 12 X-STR loci in Mongolian and Eastern Han populations of China with comparison to other populations. *Scientific Reports*. 2018; 8(1):4287. <https://doi.org/10.1038/s41598-018-22665-3> PMID: 29523825
 44. Xu J, Lei L, Zhang XJ, Fu LH, Yu F, Ma CL, et al. Population data of 12 X-chromosomal STR loci in Chinese Han samples from Hebei Province. *Forensic Sci Int Genet*. 2013; 7(2):e43–4. Epub 20121221. <https://doi.org/10.1016/j.fsigen.2012.11.007> PMID: 23266349.
 45. Bottinelli M, Gouy A, Utz S, Zieger M. Population genetic analysis of 12 X-chromosomal STRs in a Swiss sample. *Int J Legal Med*. 2022; 136(2):561–3. Epub 20210822. <https://doi.org/10.1007/s00414-021-02684-y> PMID: 34420081; PubMed Central PMCID: PMC8847170.
 46. Cainé L, Costa S, Pinheiro MF. Population data of 12 X-STR loci in a North of Portugal sample. *Int J Legal Med*. 2013; 127(1):63–4. <https://doi.org/10.1007/s00414-012-0672-z> PMID: 22297426.
 47. Edelmann J, Lutz-Bonengel S, Naue J, Hering S. X-chromosomal haplotype frequencies of four linkage groups using the Investigator Argus X-12 Kit. *Forensic Sci Int Genet*. 2012; 6(1):e24–34. Epub 20110202. <https://doi.org/10.1016/j.fsigen.2011.01.001> PMID: 21292579.
 48. Horváth G, Zalán A, Kis Z, Pamjav H. A genetic study of 12 X-STR loci in the Hungarian population. *Forensic Sci Int Genet*. 2012; 6(1):e46–7. Epub 20110403. <https://doi.org/10.1016/j.fsigen.2011.03.007> PMID: 21466981.
 49. Mršić G, Ozretić P, Crnjac J, Merkaš S, Sukser V, Račić I, et al. Expanded Croatian 12 X-STR loci database with an overview of anomalous profiles. *Forensic Sci Int Genet*. 2018; 34:249–56. Epub 20180309. <https://doi.org/10.1016/j.fsigen.2018.03.004> PMID: 29573605.
 50. Rebała K, Kotova SA, Rybakova VI, Zabauskaya TV, Shyla AA, Spivak AA, et al. Variation of X-chromosomal microsatellites in Belarus within the context of their genetic diversity in Europe. *Forensic Sci Int Genet*. 2015; 16:105–11. Epub 20150103. <https://doi.org/10.1016/j.fsigen.2014.12.011> PMID: 25579687.
 51. Robino C, Lacerenza D, Aneli S, Di Gaetano C, Matullo G, Robledo R, et al. Allele and haplotype diversity of 12 X-STRs in Sardinia. *Forensic Sci Int Genet*. 2018; 33:e1–e3. Epub 20171205. <https://doi.org/10.1016/j.fsigen.2017.12.002> PMID: 29221994.
 52. Tillmar AO. Population genetic analysis of 12 X-STRs in Swedish population. *Forensic Sci Int Genet*. 2012; 6(2):e80–1. Epub 20110802. <https://doi.org/10.1016/j.fsigen.2011.07.008> PMID: 21813348.

53. Cortés-Trujillo I, Zuñiga-Chiquette F, Ramos-González B, Chávez-Briones ML, Islas-González KL, Betancourt-Guerra DA, et al. Allele and haplotype frequencies of 12 X-STRs in Mexican population. *Forensic Sci Int Genet.* 2019; 38:e11–e3. Epub 20181023. <https://doi.org/10.1016/j.fsigen.2018.10.012> PMID: 30389253.
54. García MG, Catanesi CI, Penacino GA, Gusmão L, Pinto N. X-chromosome data for 12 STRs: Towards an Argentinian database of forensic haplotype frequencies. *Forensic Sci Int Genet.* 2019; 41:e8–e13. Epub 20190510. <https://doi.org/10.1016/j.fsigen.2019.04.005> PMID: 31085140.
55. STRUCTURE HARVESTER. Available from: <https://github.com/dentearl/structureHarvester>.
56. Earl DA, vonHoldt BM. STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conservation Genetics Resources.* 2012; 4(2):359–61. <https://doi.org/10.1007/s12686-011-9548-7>
57. CLUMPP. Available from: <https://rosenberglab.stanford.edu/clumpp.html>.
58. Jakobsson M, Rosenberg NA. CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics.* 2007; 23(14):1801–6. <https://doi.org/10.1093/bioinformatics/btm233> PMID: 17485429
59. Distruct Available from: <https://rosenberglab.stanford.edu/distruct.html>.
60. Rosenberg NA. distruct: a program for the graphical display of population structure. *Molecular Ecology Notes.* 2004; 4(1):137–8. <https://doi.org/10.1046/j.1471-8286.2003.00566.x>
61. FSTruct. Available from: <https://github.com/maikemorrison/fstruc>.
62. Morrison ML, Alcalá N, Rosenberg NA. FSTruct: An FST-based tool for measuring ancestry variation in inference of population structure. *Molecular Ecology Resources.* 2022; 22(7):2614–26. <https://doi.org/10.1111/1755-0998.13647> PMID: 35596736
63. ADZE. Available from: <https://rosenberglab.stanford.edu/adze.html>.
64. Szpiech ZA, Jakobsson M, Rosenberg NA. ADZE: a rarefaction approach for counting alleles private to combinations of populations. *Bioinformatics.* 2008; 24(21):2498–504. Epub 20080908. <https://doi.org/10.1093/bioinformatics/btn478> PMID: 18779233; PubMed Central PMCID: PMC2732282.
65. Fadhl HNM, Mohammed SA, Abdulkarim FM. Mitochondrial DNA haplogroup study: residents of Sulaymaniyah city in the Iraqi Kurdistan Region may be genetically closer to European lineage. *Egyptian Journal of Forensic Sciences.* 2021; 11(1):31. <https://doi.org/10.1186/s41935-021-00246-2>
66. Shimada I, Rand S, Brinkmann B, Hohoff C. Kurdish population data for 11 STR loci (ACTBP2, CSF1PO, FGA, TH01, TPOX, vWA, D3S1358, D5S818, D7S820, D13S317 and D21S11). *Int J Legal Med.* 2002; 116(5):301–3. Epub 20020528. <https://doi.org/10.1007/s00414-002-0294-y> PMID: 12376843.
67. Fracasso T, Schürenkamp M, Brinkmann B, Hohoff C. An X-STR meiosis study in Kurds and Germans: allele frequencies and mutation rates. *International Journal of Legal Medicine.* 2008; 122(4):353–6. <https://doi.org/10.1007/s00414-008-0231-9> PMID: 18389271
68. Kling D, Dell'Amico B, Tillmar AO. FamLinkX—implementation of a general model for likelihood computations for X-chromosomal marker data. *Forensic Science International: Genetics.* 2015; 17:1–7. <https://doi.org/10.1016/j.fsigen.2015.02.007> PMID: 25771099
69. Purps J, Siegert S, Willuweit S, Nagy M, Alves C, Salazar R, et al. A global analysis of Y-chromosomal haplotype diversity for 23 STR loci. *Forensic Science International: Genetics.* 2014; 12:12–23. <https://doi.org/10.1016/j.fsigen.2014.04.008> PMID: 24854874
70. Ramachandran S, Deshpande O, Roseman CC, Rosenberg NA, Feldman MW, Cavalli-Sforza LL. Support from the relationship of genetic and geographic distance in human populations for a serial founder effect originating in Africa. *Proceedings of the National Academy of Sciences.* 2005; 102(44):15942–7. <https://doi.org/10.1073/pnas.0507611102> PMID: 16243969