

CORRECTION

Correction: Prognosis of extracolonic findings on clinical computed tomographic colonography: A single-center experience

Hideto Tomimatsu, Keita Fujimoto, Taketo Suto, Masayuki Matsuo


In the Abstract, there is an error in the sixth sentence. The correct sentence is: Kaplan–Meier analysis revealed worse prognosis for E4 (mean survival: 5.6 years) than for E1–E3 combined (10.1 years) ($p < 0.0001$).

In the Results section, there is an error in the sixth sentence of the third paragraph. The correct sentence is: None of the patients in the Past E2 or E3 categories required further examinations that necessitated changes to the treatment plan (mean OP for E2: 2,880 days; mean OP for E3: 2,976 days).

In the Results, there is an error in the third and seventh sentence of the fourth paragraph. The correct third sentence is: The hazard ratio for Past E4 (vs the combined Past E1–E3) was 8.2 (95% CI: 3.7–18.4). The correct seventh sentence is: The hazard ratio for Revised E4 (vs the combined Revised E1–E3) was 5.2 (95% CI: 2.4–11.2).

Reference

1. Tomimatsu H, Fujimoto K, Suto T, Matsuo M. Prognosis of extracolonic findings on clinical computed tomographic colonography: A single-center experience. PLoS One. 2025;20(2):e0315601. <https://doi.org/10.1371/journal.pone.0315601> PMID: [40019913](https://pubmed.ncbi.nlm.nih.gov/40019913/)



OPEN ACCESS

Citation: Tomimatsu H, Fujimoto K, Suto T, Matsuo M (2025) Correction: Prognosis of extracolonic findings on clinical computed tomographic colonography: A single-center experience. PLoS One 20(10): e0334534. <https://doi.org/10.1371/journal.pone.0334534>

Published: October 14, 2025

Copyright: © 2025 Tomimatsu et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.