

Age-Patterns of Malaria Vary with Severity, Transmission Intensity and Seasonality in Sub-Saharan Africa: A Systematic Review and Pooled Analysis

Ilona Carneiro^{1*}, Arantxa Roca-Feltrer¹, Jamie T. Griffin², Lucy Smith¹, Marcel Tanner³, Joanna Armstrong Schellenberg¹, Brian Greenwood¹, David Schellenberg¹

1 Disease Control and Vector Biology Unit, Department of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, London, United Kingdom, **2** Department of Infectious Disease Epidemiology, MRC Centre for Outbreak Analysis and Modelling, Imperial College London, London, United Kingdom, **3** Swiss Tropical Institute, Basel, Switzerland

Abstract

Background: There is evidence that the age-pattern of *Plasmodium falciparum* malaria varies with transmission intensity. A better understanding of how this varies with the severity of outcome and across a range of transmission settings could enable locally appropriate targeting of interventions to those most at risk. We have, therefore, undertaken a pooled analysis of existing data from multiple sites to enable a comprehensive overview of the age-patterns of malaria outcomes under different epidemiological conditions in sub-Saharan Africa.

Methodology/Principal Findings: A systematic review using PubMed and CAB Abstracts (1980–2005), contacts with experts and searching bibliographies identified epidemiological studies with data on the age distribution of children with *P. falciparum* clinical malaria, hospital admissions with malaria and malaria-diagnosed mortality. Studies were allocated to a 3×2 matrix of intensity and seasonality of malaria transmission. Maximum likelihood methods were used to fit five continuous probability distributions to the percentage of each outcome by age for each of the six transmission scenarios. The best-fitting distributions are presented graphically, together with the estimated median age for each outcome. Clinical malaria incidence was relatively evenly distributed across the first 10 years of life for all transmission scenarios. Hospital admissions with malaria were more concentrated in younger children, with this effect being even more pronounced for malaria-diagnosed deaths. For all outcomes, the burden of malaria shifted towards younger ages with increasing transmission intensity, although marked seasonality moderated this effect.

Conclusions: The most severe consequences of *P. falciparum* malaria were concentrated in the youngest age groups across all settings. Despite recently observed declines in malaria transmission in several countries, which will shift the burden of malaria cases towards older children, it is still appropriate to target strategies for preventing malaria mortality and severe morbidity at very young children who will continue to bear the brunt of malaria deaths in Sub-Saharan Africa.

Citation: Carneiro I, Roca-Feltrer A, Griffin JT, Smith L, Tanner M, et al. (2010) Age-Patterns of Malaria Vary with Severity, Transmission Intensity and Seasonality in Sub-Saharan Africa: A Systematic Review and Pooled Analysis. PLoS ONE 5(2): e8988. doi:10.1371/journal.pone.0008988

Editor: Abdisalan M. Noor, The Kenya Medical Research Institute, Kenya

Received: September 18, 2009; **Accepted:** January 6, 2010; **Published:** February 1, 2010

Copyright: © 2010 Carneiro et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This project was funded by The Bill and Melinda Gates Foundation (Grant ID 33679; www.gatesfoundation.org) through the Intermittent Preventive Treatment in Infants (IPTi) consortium. I.C. is also funded by the Department for International Development, United Kingdom (www.dfid.gov.uk) through the TARGETS Communicable Disease Consortium (Grant ID HD205). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: ilona.carneiro@lshtm.ac.uk

Introduction

The peak prevalence of *Plasmodium falciparum* malaria infection is known to shift to younger age groups as transmission intensity increases [1]; there is ample evidence for this from the field [2]. However, while there is a general acceptance that the same shift in peak age occurs for severe malaria [3–8], there has been limited evidence of this for uncomplicated clinical malaria [9,10]. A recent analysis of data from seven demographic surveillance sites in sub-Saharan Africa suggested that the peak age of malaria mortality is similar across sites, despite differences in transmission intensity [11].

The “peak shift” phenomenon has been described for malaria and other infections to which immunity is acquired [12,13] and

has two implications for control. First, identification of the age groups bearing the greatest burden of clinical malaria for a given transmission setting would enable interventions to be targeted to those worst affected. Secondly, it implies that reported declines in malaria transmission intensity, and future progress towards malaria elimination, will result in a shift of malaria morbidity towards older children, as has recently been observed [14–16].

Several research groups have undertaken exercises to quantify the burden of malaria regionally or globally [17–21], or to assess how the intensity of malaria of different severities varies with transmission intensity [22–24]. However, there has been no comprehensive assessment of the comparative age-patterns of different malaria outcomes across a wide range of epidemiological

settings. This study builds on previous data abstraction efforts to collate and analyse the age distribution among children of clinical malaria, hospital admissions with malaria and deaths diagnosed as malaria. Data are abstracted from epidemiological studies across sub-Saharan Africa. We have not considered data from outside sub-Saharan Africa because of the very different epidemiology of malaria in many of these areas with *P. vivax* playing an important role. We use statistical modelling of these data to describe the age-patterns of *P. falciparum* malaria for six broad epidemiological settings categorised according to distinct combinations of malaria transmission intensity and seasonality.

Methods

Data Sources and Definitions

Systematic literature reviews were undertaken using the PubMed and CAB Abstracts (BIDS) online abstracting databases, the WHO publication Library (WHOLIS: www.who.int/library/databases/en/) and the SIGLE grey literature database (www.opensigle.inist.fr), conducting key author searches and cross-checking the bibliographies of references identified and of previous literature reviews on malaria morbidity and anaemia [25]. A review of studies that measured entomological inoculation rates (EIR) between 1970–2005 was undertaken in November 2005 using combinations of the following search terms: EIR, entomologic* inoculation rate, sporozoites inoculation rate, anoph*, vector* capacity, biting rate, sporozoite rate, sporozoites index, malaria transmission, entomol*, malaria control, light trap, pyreth* spray, human bait. The EIR studies were screened for quality, including sampling methods, length of sampling period and frequency of sampling. A review of studies that measured parasite prevalence between 1980–2006 was undertaken in February 2006 using combinations of the following search terms: malaria, falciparum, parasit*, epidemiology, parasitology, transmission, prevalence, morbidity, cross-sectional survey, child, infant. A review of studies that measured malaria morbidity and mortality outcomes between 1980–2005 was undertaken in October 2005 (see Figure 1) using combinations of the following search terms: malaria, falciparum, morbidity, mortality, epidemiology. An additional review of studies on all-cause hospital admissions from 1980–2008 was undertaken in July 2008 using combinations of the following search terms: hospitalization, hospitals/statistics and numerical data, hospitals/trends, hospitals/utilization, inpatients, hospital mortality, pediatrics. Only studies undertaken from 1980 in sub-Saharan Africa in areas endemic for *Plasmodium falciparum* and which reported on malaria outcomes in children (under 15 years of age) were included. Age of clinical malaria episodes and malaria-diagnosed deaths was obtained from community-based studies of a representative sample of the population, while age of hospital admissions with malaria was identified from hospital-based studies. Only data from baseline periods or control arms of intervention studies were included, to prevent confounding of the relationships under investigation, especially by transmission reducing interventions.

All relevant references were collected, examined and key information abstracted directly into a relational database form in Microsoft Access, which was developed to link the studies using geographical co-ordinates. When additional information was needed, authors were contacted for further details. A number of authors shared individual-level datasets, or a more detailed age-breakdown of malaria outcome data than was publically available.

Data on the incidence of clinical malaria were obtained from longitudinal studies using either active or passive case detection. Case definitions varied according to (i) whether or not reported

history of fever was considered, and over what time period (24 or 48 hours), (ii) whether a concurrent axillary ($\geq 37.5^{\circ}\text{C}$) or rectal ($\geq 38.5^{\circ}\text{C}$) temperature measurement was considered, and (iii) whether a parasite density threshold was used. An age-specific fever density threshold calculated according to previously described methods to ascertain the malaria-attributable fraction of fevers in endemic areas [26] was used in some studies [27–32]. Where data were available to us [33–40], the number of malaria-attributable fevers in each age group was calculated using the following formula:

$$I_m = I_p - \frac{(I_n \times P)}{(1 - P)}$$

where I_m is the incidence of fevers truly attributable to malaria, I_p is the incidence of fevers associated with malaria parasites, I_n is the incidence of fevers not associated with malaria parasites and P is the prevalence of *P. falciparum*. In some cases, data on age-specific fever incidence or parasite prevalence was obtained from other studies from the same site and linearly interpolated if the same age-categories were not available.

It was not possible to estimate the malaria-attributable incidence of fevers adjusting for changes in a parasite density threshold with age for 11 of the 29 studies included in this analysis. Of these, 6/11 used a fixed fever density threshold, with 3/6 calculating a locally specific density threshold based on the malaria-attributable fraction. A sensitivity analysis was undertaken to assess the impact of using different threshold densities on the results.

Hospital admissions with malaria were all (or, in some cases, a proportional sample) of admissions to the paediatric ward of a given hospital who had a positive rapid diagnostic test or blood film for malaria. All-cause hospital admissions were all admissions to a paediatric ward. Malaria-diagnosed deaths were obtained from studies using recognised verbal autopsy methods, and include all acute febrile illness deaths minus those with an obvious alternative cause. Data on all-cause deaths were obtained from the malaria-diagnosed deaths literature review, and were not searched for separately.

Classification of Study Site Transmission Intensity and Seasonality

Given the imprecision and temporal variability of EIR estimates, malaria transmission intensity was arbitrarily categorised into three broad categories as low (EIR < 10 bites per person per year (pppy)), medium (EIR 10–100 bites pppy) or high (EIR > 100 bites pppy). Settings where malaria is epidemic were not included. Few studies reported good quality EIR estimates, and the previously described log-linear relationship between EIR and parasite prevalence [41,42] was independently confirmed from the studies identified through our searches. The log-linear relationship identified cut-offs of <25%, 25–60% and >60% parasite prevalence in children <5 years of age to be approximately concordant with the low, medium and high transmission intensity using EIR [25]. Using the geo-referenced relational database, each study was assigned to a transmission intensity category on the basis of EIR or parasite prevalence studies from within the same second administrative level, matched as closely as possible in time.

The seasonality of malaria transmission for a given setting was difficult to define, with contradictory descriptions sometimes given by different authors for the same site. Preliminary analysis of monthly incidence data revealed a continuum of seasonal patterns, which could be arbitrarily divided into those with $\geq 75\%$ of




Figure 1. Systematic literature review process. Flow diagram describing the systematic review of literature on the age-breakdown of uncomplicated clinical malaria, hospital admissions with malaria and malaria-diagnosed deaths.
doi:10.1371/journal.pone.0008988.g001

episodes concentrated in ≤ 6 months of the year, considered to have “marked seasonality”, and those with no marked seasonality [43]. The MARA (Mapping Malaria in Africa) maps of seasonality of climate suitability for malaria were consulted [44] if consistent definitions were not available from published references. In a few instances where no additional data on transmission intensity or seasonality were available for a given study, malaria experts with local knowledge were consulted. See Tables S1, S2, and S3 for criteria and sources used to allocate each study to one of six cells of a transmission intensity-seasonality matrix as presented in Table 1: low, medium, high transmission, with marked or no marked seasonality.

Statistical Analyses

The malaria outcomes described are incidence of clinical malaria with fever, numbers of hospital admissions with malaria parasites and numbers of malaria-diagnosed deaths in the community. Data on age of all-cause hospital admissions and all-cause mortality were also abstracted for comparative purposes. Data on each outcome were abstracted by age (in months under five-years of age, if possible, and then by year) up to 15 years. Studies based on very small sample sizes (< 50 cases or person-years at risk) were excluded because the percentage age-distributions could easily be spurious. For the hospital admissions with malaria and malaria-diagnosed deaths, 1 [45] and 2 [46,47] studies respectively which reported only

Table 1. Distribution of research study countries and sites by matrix of transmission intensity and seasonality.

Outcome	Seasonality	Transmission Intensity (Entomological Inoculation Rate [bites per person per year (pppy)])		
		<10 bites pppy*	10–100 bites pppy	>100 bites pppy
Clinical malaria	Marked seasonality	Mali (Sotuba), Senegal (Central & South Dakar)	Cameroon (Ebolakounou), Mali (Sotuba), Mozambique (Maputo), Senegal (Niakhar), The Gambia (Upper Baddibu)	Burkina Faso (Nouna), Ghana (Kassena-Nankhana), Mali (Doneguebougou)
	No marked seasonality	Ghana (Prampam), Uganda (Kampala),	Gabon (Lambarene), Mozambique (Manhica), Tanzania (Ifakara, Muheza highlands)	Benin (Atlantic Coast), Cameroon (Koundou), Cote d'Ivoire (Korhogoro), Ghana (Afigya Sekyere), Kenya (Asembo Bay), Senegal (Dielmo), Tanzania (Idete, Muheza, Tanga)
Hospital admissions with malaria parasites	Marked Seasonality	Guinea Bissau (Bissau), Senegal (Dakar), Sudan (Geradif), Tanzania (Huruma, Kibosho), The Gambia (Banjul, Sibanor)	Burkina Faso (Ougadougou), Mozambique (Maputo), Tanzania (Same), Zambia (Choma)	Burkina Faso (Sorou & Nayala)
	No marked seasonality	-	Benin (Cotonou), D. R. Congo (Brazzaville, Kinshasa), Gabon (Lambarene, Libreville), Ghana (Kumasi), Kenya (Kilifi), Malawi (Blantyre, Mangochi, Zomba), Mozambique (Manhica), Nigeria (Ilorin), Tanzania (Ifakara, Moshi)	Ghana (Tamale), Tanzania (Tanga)
Malaria-diagnosed mortality	Marked Seasonality	-	The Gambia (Farafenni, South Bank, Upper River Division), Senegal (Niakhar)	Burkina Faso (Kourweogo, Nouna, Oubritenga), Ghana (Kassena-Nankhana), Senegal (Bandafassi)
	No marked seasonality	-	Kenya (Kilifi), Mozambique (Manhica), Tanzania (Ifakara, Rufiji)	Burundi (Nyanzalac), D. R. Congo (Katana), Guinea (Mandiana), Kenya (Asembo Bay, Kisumu), Tanzania (Bagamoyo)

doi:10.1371/journal.pone.0008988.t001

on children less than 5 years of age were excluded as these would otherwise skew the age distribution as there was no denominator for analysis of these outcomes.

All data were analysed using Stata 10.1 (StataCorp. 2007. Stata Statistical Software: Release 10. College Station, TX: StataCorp LP). A user-written (JG) command (details available on request from the authors) was used to censor the data at different age intervals (interval-censored) and to use data with different maximum age values (right-truncated). This enabled the inclusion of studies that reported on different age ranges and used different age groupings. The percentage of each outcome by age (0–10 years - excluding neonates for malaria-diagnosed mortality) was calculated for each study. Studies from the same research setting were included if there was no temporal overlap. However, it was not possible to adjust the analyses for intra-cluster correlation within site due to small numbers of studies from the same site. Data from sites in the same cell of the transmission matrix were analysed together and five continuous probability distributions (gamma, Weibull, log-normal, log-logistic, exponential) were fitted to the data using a time-to-event likelihood for severe events, or Poisson likelihood to account for the person-time at risk for the incidence of clinical malaria. The distributions with the lowest AIC (Akaike Information Criterion) value were identified as the best fitting [48]. For the analysis of malaria-diagnosed deaths, the fitted distributions were re-scaled to exclude children under 1 month of age, as neonatal deaths are difficult to attribute and malaria is not considered a major problem in this age group. The median age was calculated as the 50th percentile of the best-fitting distribution for each outcome and transmission scenario.

Results

A total of 29, 36 and 21 studies were included in the analysis of clinical malaria, hospital admissions with malaria, and malaria-diagnosed mortality respectively (see Table 2). In addition, 22 studies on all-cause admissions and 28 studies on all-cause mortality were also analysed (data not shown). For malaria and all-cause hospital admissions, there were no data from areas with no marked seasonality and low transmission intensity. For all-cause admissions there were also no data for areas of high intensity and marked seasonality. However, the analyses included 3 studies from areas with sporadic or no malaria transmission. For malaria-diagnosed mortality, there were no data from areas with low transmission.

Figure 2 shows the percentage of each outcome by age for children under 10 years, such that the integral of the curve is equal to one, i.e. 100% of expected cases. Clinical malaria is relatively evenly distributed across all ages with a shift towards younger age groups as transmission intensity increases, both in areas of non-marked and marked seasonality. The median age for clinical malaria ranges from 32 months (Inter-quartile range (IQR): 15, 61) in settings of highly intense and not markedly seasonal transmission, to 72 months (IQR: 45, 97) in settings of low intensity and markedly seasonal transmission. In six studies, the estimation of malaria-attributable fevers resulted in negative values for some age-groups, and these were considered to be zero cases of malaria-attributable fevers. For one study [37], all age-groups had a higher asymptomatic prevalence of infection than prevalence of fevers associated with parasites, resulting in this study contributing

Table 2. Number of studies, (research sites) and [sources of data] included in the analyses, with best-fitting probability distributions.

Outcome	Seasonality	Transmission Intensity (Entomological Inoculation Rate [bites per person per year (pppy)])		
		<10 bites pppy	10–100 bites pppy	>100 bites pppy
Clinical malaria	Marked seasonality	3 (2) [10,39,60] Log-normal	5 (5) [10,29,32,61,62] Log-normal	4 (3) [10,37,45,63] Log-normal
	No marked seasonality	2 (2) [27,38] Log-normal	6 (5) [34,36,40,64–66] Gamma	9 (9) [28–31,33,35,36,67–69] Log-normal
Hospital admissions with malaria parasites	Marked Seasonality	9 (6) [6,70–76] Log-logistic	6 (4) [5,6,77–80] Log-logistic	1 (1) [5] Log-logistic
	No marked seasonality	0-	18 (14) [6,14,74,81–90] Log-normal	2 (2) [6,91] Log-logistic
Malaria-diagnosed mortality	Marked Seasonality	0-	6 (4) [92–97] Log-normal	6 (5) [11,96,98,99] Log-logistic
	No marked seasonality	0-	4 (4) [11,100] Log-logistic	5 (5) [11,101–104] Gamma


doi:10.1371/journal.pone.0008988.t002

an incidence of zero malaria-attributable cases across the age-groups 6–24 months.

A sensitivity analysis using no threshold density and a fixed density cut-off for those studies for which data were available suggested that the age-pattern would be slightly more evenly distributed across age groups if an age-specific cut-off was not used (data not shown). Therefore, the results presented here are likely to partially underestimate the proportion of clinical malaria that occurs in younger age groups, especially for areas of marked seasonality where 7/12 studies used a fixed density threshold or no threshold density to define clinical malaria.

Hospital admissions with malaria parasites are more concentrated in younger children than is clinical malaria in all settings, and these severe cases become more concentrated in younger ages with increasing transmission intensity and less seasonality. The median age ranges from 17 months (IQR: 10, 29) at high, not markedly seasonal transmission to 36 months (IQR: 20, 60) at low and markedly seasonal transmission.

Malaria-diagnosed mortality is more focussed in younger children than admissions with malaria in all settings for which there are comparative data. Once more there is a distinctive shift of the peak age towards infants as transmission becomes more

**Figure 2.** Age-patterns of *P. falciparum* malaria in Sub-Saharan Africa. Percentage of uncomplicated clinical malaria, hospital admissions with malaria and malaria-diagnosed deaths per month of age in children under ten years of age, by transmission intensity (TI) and seasonality of malaria transmission.

doi:10.1371/journal.pone.0008988.g002

intense. The median age for malaria-diagnosed mortality ranges from 12 months (IQR: 6, 22) at high, not markedly seasonal transmission to 28 months (IQR: 15, 51) at medium and markedly seasonal transmission.

For all three outcomes, the age-patterns are less biased towards younger ages in areas of marked seasonality than in areas without marked seasonality for a given intensity of transmission. Table 3 shows the median ages and inter-quartile ranges for each outcome for each cell of the transmission matrix.

Discussion

We have compiled and analysed available data on *P. falciparum* malaria outcomes of different severities across all transmission intensities in sub-Saharan Africa. Our analyses confirm previous findings that the peak age of uncomplicated clinical malaria declines with increasing transmission intensity. The most striking finding, however, was the extent to which hospital admissions with malaria, and especially malaria-diagnosed deaths, occur in the youngest age groups regardless of transmission intensity. This has implications for the age-targeting of malaria control strategies.

For uncomplicated clinical malaria, the distribution is relatively even across all ages under-10 years with a slight tendency for the age-pattern to shift towards younger age groups with increasing transmission intensity. This is similar, although less pronounced, to previously published comparative results from Senegal [9] and Mali [10], some of which was excluded from this pooled analysis (due to lack of denominator data for the Ndiop study [9]).

It is likely that inclusion of data without an age-specific density threshold for one third of the studies will have resulted in a dampening of the peak-shift effect reported here. The incidence of non-malarial fever with incidental parasitaemia would be expected to increase with age as parasite tolerance is acquired. The use of no fever density threshold would therefore result in an overestimate of the proportion of cases in older age groups, while the use of a fixed “average” density threshold would result in an underestimate of the proportion of cases at younger ages and an overestimate of cases at older ages. Estimation of the true malaria-attributable incidence for many studies was hampered by a lack of site-specific information on either all-cause fever or parasite prevalence for the same age-ranges covered by our data. An increase in the specificity of the case definition would shift the age-distribution of malaria-attributable fevers further towards younger ages, and this would increase with transmission intensity, as parasite tolerance would develop earlier in life. In addition, not all studies removed the time at risk after an antimalarial treatment, and this will serve to underestimate the true incidence. This bias will be more pronounced in the peak incidence age-groups

resulting in a dampening of the age-patterns, especially at higher transmission intensities.

Hospital admissions for malaria are concentrated in children under-5 years of age in all settings, with a shift towards younger ages with increasing intensity of malaria transmission. This is consistent with previous reports that have compared data across sites with different transmission intensities [3,4], as well as within sites with observed declines in transmission intensity over time [14–16]. The age-distributions of different syndromes of severe malarial disease have previously been shown to vary with transmission intensity [23,49,50]. A comprehensive analysis of specific outcomes across a range of transmission settings has been undertaken and will be presented elsewhere (Roca-Feltrer *et al.*, manuscript in preparation).

Our analysis shows that, across all ranges of transmission intensity, malaria-diagnosed mortality is focussed in the very young. This is in keeping with recently published data on age-specific malaria mortality rates from a number of sites in Africa, which showed the peak age of malaria deaths to be in children under 1 year of age [11].

Our findings have implications for malaria control programmes that aim to tackle the worst effects of malaria. Based on the age-pattern of clinical episodes and severe disease, the burden of malaria deaths in the very young would likely be underestimated. The situation is compounded by the fact that, in sub-Saharan Africa, the majority of malaria deaths occur outside hospital and are therefore less visible than cases of severe disease. Age-targeted strategies such as intermittent preventive treatment of infants (IPTi) [51] and children (IPTc) [52], and vaccination against malaria [53] are currently being investigated and are likely to have a role to play in settings with moderate and high transmission [54]. The intensity of malaria is expected to decline in many areas as efforts are made towards elimination [55], which will require a move towards universal coverage with preventive interventions [56]. However, it is likely that many of the countries in Africa will not reach this situation for many years to come.

The age-patterns of all-cause mortality and all-cause hospital admissions did not vary by malaria transmission setting (data not shown). This may be due, in part, to the strong influence of neonatal disease and death on these age-patterns, which we were unable to deconstruct, as most data sets did not sufficiently disaggregate the neonatal data. One exception was for all-cause admissions at medium transmission intensity and marked seasonality of malaria, for which the data were only available for a smaller age-range (no neonates for 2 studies and <8 years for all 3 studies).

The effect of seasonality of malaria transmission on the three outcomes was to dampen the impact of transmission intensity on

Table 3. Median ages in months (inter-quartile range) for each outcome for each transmission matrix cell.

Outcome	Seasonality	Transmission Intensity (Entomological Inoculation Rate [bites per person per year (pppy)])		
		<10 bites pppy	10–100 bites pppy	>100 bites pppy
Clinical malaria	Marked seasonality	72 (45, 97)	65 (35, 92)	37 (17, 67)
	No marked seasonality	62 (37, 89)	37 (20, 60)	33 (16, 62)
Hospital admissions with malaria parasites	Marked Seasonality	36 (20, 60)	34 (20, 55)	22 (14, 34)
	No marked seasonality	-	22 (12, 41)	17 (10, 29)
Malaria-diagnosed mortality	Marked Seasonality	-	28 (15, 51)	15 (8, 29)
	No marked seasonality	-	17 (7, 36)	12 (6, 22)

doi:10.1371/journal.pone.0008988.t003

the age-patterns. Within each intensity category, the peak age is lower for settings with no marked seasonality than for those with marked seasonality. This has not previously been reported, but could have been predicted. The “peak-shift” phenomenon is consistent with gradually acquired protective immunity [12], and the effect of seasonal transmission would be to reduce the cumulative exposure of individuals to malaria and hence their acquisition of immunity.

The fact that the peak age declines with increasing severity of the outcome, implies that protection against the more severe forms of malaria is acquired more rapidly than protection against clinical malaria, as previously suggested [57]. This phenomenon may be compounded by a tendency for those who are unable to develop a protective immune response to succumb to the disease and die in their early years. The strong bias for severe malaria outcomes to be concentrated in infants across different transmission intensities lends weight to the suggestion that protection against severe falciparum malaria may be acquired with age, independently of, or in synergy with prior exposure through mechanisms that are not yet fully understood.

As with any major data abstraction exercise our analyses had some limitations. The data abstracted showed a bias towards established research settings, with gaps in the availability of data from countries in central Africa. The lack of data from settings with no marked seasonality and low intensity probably reflects a lack of research facilities in such settings; most malaria research sites are in areas of medium to high transmission intensity. Nevertheless, data from research settings are likely to be susceptible to the Hawthorne effect (where the presence of a dedicated research team alters the disease epidemiology) as a result of increased implementation of interventions potentially leading to a decline in transmission [14–16]. With further progress in malaria control, there is likely to be an increase in data for low transmission intensity settings. These geographic and epidemiological ‘black holes’ in malaria research need to be filled urgently.

In addition to the lack of data from some epidemiological categories, it was generally difficult to obtain age-specific data on malaria morbidity and mortality, which are usually not published in sufficient detail to be included in a thorough statistical analysis. This was ameliorated to some extent by contacting authors. Allocating sites to transmission intensity bands may also have been subject to some misclassification, both in terms of a lack of local data on intensity parameters (EIR or parasite prevalence), and also in relation to temporal changes. It is very rare to have geographically and temporally matched data on EIR or parasite prevalence and morbidity/mortality outcomes, and local or expert knowledge was sought where possible to support the classification. The categorisation of settings as having marked seasonality or no marked seasonality may also be subject to some disagreement, as seasonality is a continuum, with most settings showing some seasonal peaks and some perennial settings having a distinct clustering of cases. However, sensitivity analyses of the definition used suggested that sites showed consistent patterns of seasonality for different malaria outcomes and across several years [43]. Given the limitations in defining the epidemiological context for a particular setting, the analyses and results were restricted to six broad categories defined by the transmission matrix.

The allocation of some sites may change over time, e.g. the measured EIR for Sotuba, Mali was 12 in 1999 and 4 in 2000 [10]. We temporally matched the outcome data to the epidemiological data, where possible, so that sites may appear in more than one cell of the matrix depending on the timing of the studies. The relationships presented here are between the burden of disease, and the intensity and seasonality of malaria transmission,

and as such, are independent of the changing epidemiology of malaria.

We attempted to estimate the incidence of truly malaria-attributable fevers, rather than fevers incidentally associated with parasites, by using the age-specific all-cause fever incidence and parasite prevalence, where this was available. In addition to a lack of suitable data to enable this adjustment to be made for many studies, it is likely that the use of non-contemporary data reduced the utility of this approach. For example, using cross-sectional parasite prevalence from peak transmission seasons to adjust the incidence over a longer period of time could result in an underestimate of malaria-attributable cases.

It is also important to note the limitations in diagnosing malaria deaths through the verbal autopsy method, which suffers from a lack of a distinct symptom complex for malaria [11,58]. In addition, malaria is also likely to be an underlying cause in many deaths, but missed as the final diagnosis if a more distinct terminal event occurs. While it is difficult to estimate the incidence of true malaria-attributable mortality, it is likely that the mis-classification of malaria deaths will be internally consistent within a study, and thus analysis of percentage age-distributions will suffer less from such errors than would cross-site comparisons.

No attempt was made here to assess the relationship between the *incidence* (as opposed to the percentage distribution) of disease and transmission intensity using these data. Such exercises have previously been undertaken [4,22–24], but there are several limitations to such an endeavour, limiting the number of studies that can be included. While most recent estimates of mortality come from community-based demographic surveillance systems, it was difficult to obtain age-specific denominator population data from many of the older studies. Data on severe malaria outcomes can only be obtained at the hospital level, as continued community surveillance will reduce the progression to severe disease. Unfortunately, it is difficult to estimate hospital catchment populations, as individuals may not access the nearest facility, and local access issues (geography, socio-economics, treatment-seeking behaviour) make it difficult to develop a standardised definition. While mathematical modelling may go some way to alleviating this issue [59] it is reliant on additional data from the site, which is not always available. Estimates of incidence are thus hampered by the lack of a denominator population for many studies of severe outcomes. In addition, given that most data come from established research settings, the presence of a dedicated malaria research team, with improved health infrastructure and/or surveillance, is itself likely to reduce the burden of malaria that would otherwise occur in that setting. In addition, local practices and access to interventions and health care may confound the relationship between transmission intensity and the age-specific burden of malaria. Nevertheless, this relationship should be the focus of any future extensions to such an exercise, as this will be most informative for public health planning.

The data presented here show that the most severe consequences of malaria are concentrated in the youngest age groups, and that the burden of malaria shifts towards younger age groups with increasing transmission intensity. Recently observed declines in malaria transmission intensity are likely to result in the burden of malaria shifting towards older children. Nevertheless, while the aim of malaria control remains the prevention of the most severe forms of malaria disease and death [56], targeting malaria interventions to very young children will produce important benefits. Integrated malaria strategies need to be tailored to the epidemiological context, and to be able to respond to changes in malaria transmission over time. As malaria control improves, the median age of severe disease manifestations will increase.

Nevertheless, a disproportionate number of malaria deaths will continue to occur in the very young and it is still worth targeting specific control strategies at this age group.

Supporting Information

Table S1 Sources used to allocate clinical malaria studies to a matrix of intensity and seasonality of malaria.

Found at: doi:10.1371/journal.pone.0008988.s001 (0.45 MB DOC)

Table S2 Sources used to allocate studies on hospital admissions with malaria to a matrix of intensity and seasonality of malaria.

Found at: doi:10.1371/journal.pone.0008988.s002 (0.51 MB DOC)

Table S3 Sources used to allocate studies of malaria-diagnosed deaths to a matrix of intensity and seasonality of malaria.

Found at: doi:10.1371/journal.pone.0008988.s003 (0.27 MB DOC)

Acknowledgments

All authors that took the time to respond to queries, to provide more detailed data than is publically available, or to publish their data with sufficiently detailed information to enable these analyses. Specifically Pedro Alonso, Kevin Baird, Quique Bassat, Fred Binka, Sarah Bonnet, Daniel

Chandramohan, Badara Cisse, Tamara Clark, Chris Curtis, Alassane Dicko, Grant Dorsey, Juliana Gabor, Hayder Giha, Martin Grobusch, Robert Guigumde, Caterina Guinovart, Lars Henning, Marie Claire Henry, Michael van Hensbroek, Patrick Imbert, Muminatou Jallow, Lawrence Kazembe, Andrew Kitua, Peter Kremsner, Peter McElroy, Caroline Maxwell, Frank Mockenhaupt, Saul Morris, Paul Mulumba, Dr Olanrewaju, Hugh Keyburn, Eleanor Riley, Christophe Rogier, Alan Schapira, Erich Schmutzhard, Phil Thuma, Jim Todd, Jean-Francois Trape, Janneke van Dijk, and Luis Varandas; the Scientific Co-ordination Committee of the MRC Laboratories in The Gambia and the Co-ordinating Committee of the Joint Malaria Programme in Tanzania. We are extremely grateful to Terrie Taylor on behalf of the Severe Malaria in African Children (SMAC) Network who provided individual-level hospital admissions data from six sites, the INDEPTH Network who provided age-breakdown data on mortality from seven demographic surveillance sites and Konstantina Boutsika on behalf of the Mapping Malaria Risk in Africa (MARA) collaboration for additional data on parasite prevalence age-breakdown. See www.iptwebtool.org for further acknowledgements relating to these and other data. Tom Smith for data and helpful insights, and Amanda Ross, Rob Newman and Andrea Egan for comments on the draft manuscript.

Author Contributions

Conceived and designed the experiments: IC ARF MT JAS BG DS. Analyzed the data: IC ARF. Contributed reagents/materials/analysis tools: JTG. Wrote the paper: IC ARF JTG LS MT JAS BG DS. Conducted associated systematic reviews and abstracted data: LS.

References

- Boyd MF (1949) Epidemiology of malaria: factors related to the intermediate host. In: Boyd MF, ed. *Malaria: A comprehensive survey of all aspects of this group of diseases from a global standpoint*. W.B. Saunders Company.
- Molineaux L, Gramiccia G (1980) The Garki Project: Research on the epidemiology and control of malaria in the Sudan Savanna of West Africa. Geneva: World Health Organization.
- Snow RW, Bastos de Azevedo I, Lowe BS, Kabiru EW, Nevill CG, et al. (1994) Severe childhood malaria in two areas of markedly different falciparum transmission in east Africa. *Acta-Trop* 57: 289–300.
- Snow RW, Omumbo JA, Lowe B, Molyneux CS, Obiero JO, et al. (1997) Relation between severe malaria morbidity in children and level of Plasmodium falciparum transmission in Africa [see comments]. *Lancet* 349: 1650–1654.
- Modiano D, Sirima BS, Sawadogo A, Sanou I, Pare J, et al. (1998) Severe malaria in Burkina Faso: influence of age and transmission level on clinical presentation. *American Journal of Tropical Medicine and Hygiene* 59: 539–542.
- Reyburn H, Mbatia R, Drakeley C, Bruce J, Carneiro I, et al. (2005) Association of transmission intensity and age with clinical manifestations and case fatality of severe Plasmodium falciparum malaria. *JAMA* 293: 1461–1470.
- Idro R, Aloyo J, Mayende L, Bitarwakwe E, John CC, et al. (2006) Severe malaria in children in areas with low, moderate and high transmission intensity in Uganda. *Trop Med Int Health* 11: 115–124.
- Okiro EA, Al-Taiar A, Reyburn H, Idro R, Berkley JA, et al. (2009) Age patterns of severe paediatric malaria and their relationship to Plasmodium falciparum transmission intensity. *Malar J* 8: 4.
- Trape J-F, Rogier C (1996) Combating malaria morbidity and mortality by reducing transmission. *Parasitology Today* 12: 236–240.
- Dicko A, Sagara I, Diemert D, Sogoba M, Niambele MB, et al. (2007) Year-to-year variation in the age-specific incidence of clinical malaria in two potential vaccine testing sites in Mali with different levels of malaria transmission intensity. *Am J Trop Med Hyg* 77: 1028–1033.
- Abdullah S, Adazu K, Masanja H, Diallo D, Hodgson A, et al. (2007) Patterns of age-specific mortality in children in endemic areas of sub-Saharan Africa. *Am J Trop Med Hyg* 77: 99–105.
- Anderson RM, May RM (1992) *Indirectly transmitted microparasites. Infectious diseases of humans: Dynamics and control*. Oxford: Oxford University Press. pp 379–383.
- Woolhouse ME (1998) Patterns in parasite epidemiology: the peak shift. *Parasitol Today* 14: 428–434.
- Schellenberg D, Menendez C, Aponte J, Guinovart C, Mshinda H, et al. (2004) The Changing Epidemiology of Malaria in Ifakara Town, southern Tanzania. *Tropical Medicine and International Health* 9: 68–76.
- Ceesay SJ, Casals-Pascual C, Erskine J, Anya SE, Duah NO, et al. (2008) Changes in malaria indices between 1999 and 2007 in The Gambia: a retrospective analysis. *Lancet* 372: 1545–1554.
- O'Meara WP, Bejon P, Mwangi TW, Okiro EA, Peshu N, et al. (2008) Effect of a fall in malaria transmission on morbidity and mortality in Kilifi, Kenya. *Lancet* 372: 1555–1562.
- Murray CJL, Lopez A (1996) *Global Health Statistics*. Cambridge, Harvard University Press (Global Burden of disease and Injury Series) 2.
- Snow RW, Craig M, Deichmann U, Marsh K (1999) Estimating mortality, morbidity and disability due to malaria among Africa's non-pregnant population [see comments]. *Bull World Health Organ* 77: 624–640.
- Snow RW, Guerra CA, Noor AM, Myint HY, Hay SI (2005) The global distribution of clinical episodes of Plasmodium falciparum malaria. *Nature* 434: 214–217.
- Murphy SC, Breman JG (2001) Gaps in the childhood malaria burden in Africa: cerebral malaria, neurological sequelae, anemia, respiratory distress, hypoglycemia, and complications of pregnancy. *Am J Trop Med Hyg* 64: 57–67.
- Roca-Feltrer A, Carneiro I, Armstrong Schellenberg JR (2008) Estimates of the burden of malaria morbidity in Africa in children under the age of 5 years. *Trop Med Int Health* 13: 771–783.
- Smith TA, Leuenberger R, Lengeler C (2001) Child mortality and malaria transmission intensity in Africa. *Trends Parasitol* 17: 145–149.
- Snow RW, Marsh K (2002) The consequences of reducing transmission of Plasmodium falciparum in Africa. *Adv Parasitol* 52: 235–264.
- Smith T, Killeen G, Lengeler C, Tanner M (2004) Relationships between the outcome of Plasmodium falciparum infection and the intensity of transmission in Africa. *Am J Trop Med Hyg* 71: 80–86.
- Roca-Feltrer A (2008) *Estimating The Burden And The Age Pattern Of Malaria Morbidity In Sub-Saharan Africa In Under-Fives*. London: University of London. pp 423.
- Smith T, Schellenberg JA, Hayes R (1994) Attributable fraction estimates and case definitions for malaria in endemic areas. *Stat-Med* 13: 2345–2358.
- McGuinness D, Koram K, Bennett S, Wagner G, Nkrumah F, et al. (1998) Clinical case definitions for malaria: clinical malaria associated with very low parasite densities in African infants. *Trans R Soc Trop Med Hyg* 92: 527–531.
- Bloland PB, Boriga DA, Ruebush TK, McCormick JB, Roberts JM, et al. (1999) Longitudinal cohort study of the epidemiology of malaria infections in an area of intense malaria transmission II. Descriptive epidemiology of malaria infection and disease among children. *Am J Trop Med Hyg* 60: 641–648.
- Bonnet S, Paul RE, Gouagna C, Safeukui I, Meunier JY, et al. (2002) Level and dynamics of malaria transmission and morbidity in an equatorial area of South Cameroon. *Trop Med Int Health* 7: 249–256.
- Henry MC, Rogier C, Nzeyimana I, Assi SB, Dossou-Yovo J, et al. (2003) Inland valley rice production systems and malaria infection and disease in the savannah of Cote d'Ivoire. *Trop Med Int Health* 8: 449–458.
- Rogier C, Trape JF (1993) Malaria attacks in children exposed to high transmission: who is protected? *Trans R Soc Trop Med Hyg* 87: 245–246.
- Thompson R, Begtrup K, Cuamba N, Dgedge M, Mendis C, et al. (1997) The Matola malaria project: a temporal and spatial study of malaria transmission

- and disease in a suburban area of Maputo, Mozambique. *Am J Trop Med Hyg* 57: 550–559.
33. Alonso PL, Smith T, Armstrong Schellenberg JRM, Masanja H, Mwankusye S, et al. (1994) Randomised trial of efficacy of SPf66 vaccine against *Plasmodium falciparum* malaria in children in southern Tanzania. *Lancet (British edition)* 344: 1175–1181.
 34. Henning L, Schellenberg D, Smith T, Henning D, Alonso P, et al. (2004) A prospective study of *Plasmodium falciparum* multiplicity of infection and morbidity in Tanzanian children. *Trans R Soc Trop Med Hyg* 98: 687–694.
 35. Kitua AY, Smith T, Alonso PL, Masanja H, Urassa H, et al. (1996) *Plasmodium falciparum* malaria in the first year of life in an area of intense and perennial transmission. *Tropical Medicine and International Health* 1: 475–484.
 36. Maxwell CA, Chambo W, Mwaimu M, Magogo F, Carneiro IA, et al. (2003) Variation of malaria transmission and morbidity with altitude in Tanzania and with introduction of alphacypermethrin treated nets. *Malar J* 2: 28.
 37. Baird JK, Owusu AS, Utz GC, Koram K, Barcus MJ, et al. (2002) Seasonal malaria attack rates in infants and young children in northern Ghana. *Am J Trop Med Hyg* 66: 280–286.
 38. Davis JC, Clark TD, Kemble SK, Talemwa N, Njama-Meya D, et al. (2006) Longitudinal study of urban malaria in a cohort of Ugandan children: description of study site, census and recruitment. *Malar J* 5: 18.
 39. Diallo S, Ndir O, Faye O, Diop BM, Dieng Y, et al. (1998) [Malaria in the southern sanitary district of Dakar (Senegal). 1. Parasitemia and malarial attacks]. *Bull Soc Pathol Exot* 91: 208–213.
 40. Guinovart C, Bassat Q, Sigauque B, Aide P, Sacarlal J, et al. (2008) Malaria in rural Mozambique. Part I: children attending the outpatient clinic. *Malar J* 7: 36.
 41. Beier JC, Killeen GF, Githure JJ (1999) Short report: entomologic inoculation rates and *Plasmodium falciparum* malaria prevalence in Africa. *Am J Trop Med Hyg* 61: 109–113.
 42. Hay SI, Rogers DJ, Toomer JF, Snow RW (2000) Annual *Plasmodium falciparum* entomological inoculation rates (EIR) across Africa: literature survey, Internet access and review. *Trans R Soc Trop Med Hyg* 94: 113–127.
 43. Roca-Feltrer A, Armstrong Schellenberg JR, Smith L, Carneiro I (2009) A simple method for defining malaria seasonality. *Malar J* 8: 276.
 44. Mapping Malaria Risk in Africa (2008) Duration of Malaria Transmission Season.
 45. Chandramohan D, Owusu-Agyei S, Carneiro I, Awine T, Amponsa-Achiano K, et al. (2005) Cluster randomised trial of intermittent preventive treatment for malaria in infants in area of high, seasonal transmission in Ghana. *Bmj* 331: 727–733.
 46. Slutsker L, Bloland P, Steketee RW, Wirima JJ, Heymann DL, et al. (1996) Infant and second-year mortality in rural Malawi: causes and descriptive epidemiology. *Am J Trop Med Hyg* 55: 77–81.
 47. Kuate Defo B (1995) Epidemiology and control of infant and early childhood malaria: a competing risks analysis. *Int J Epidemiol* 24: 204–217.
 48. Anderson DR, Burnham KP (2002) Model selection and multimodel inference: a practical information-theoretic approach. Springer.
 49. Marsh K (1992) Malaria - a neglected disease? *Parasitology* 104: S53–S69.
 50. Marsh K, Snow RW (1997) Host-parasite interaction and morbidity in malaria endemic areas. *Philos Trans R Soc Lond B Biol Sci* 352: 1385–1394.
 51. Grobusch MP, Egan A, Gosling RD, Newman RD (2007) Intermittent preventive therapy for malaria: progress and future directions. *Curr Opin Infect Dis* 20: 613–620.
 52. Greenwood B (2006) Review: Intermittent preventive treatment—a new approach to the prevention of malaria in children in areas with seasonal malaria transmission. *Trop Med Int Health* 11: 983–991.
 53. Bejon P, Lusingu J, Olotu A, Leach A, Lievens M, et al. (2008) Efficacy of RTS,S/AS01E vaccine against malaria in children 5 to 17 months of age. *N Engl J Med* 359: 2521–2532.
 54. IOM IoM (2008) Assessment of the Role of Intermittent Preventive Treatment for Malaria in Infants: Letter Report. Washington DC: IOM. 80 p.
 55. Tanner M, de Savigny D (2008) Malaria eradication back on the table. *Bull World Health Organ* 86: 82.
 56. Roll Back Malaria Partnership (2008) Global Malaria Action Plan for a Malaria-free World.
 57. Snow RW, Gilles HM (2002) The epidemiology of malaria. In: Warrell DA, Gilles HM, eds. *Essential malariaology*. 4th ed. London: Arnold. pp 97–100.
 58. Rowe AK, Rowe SY, Snow RW, Korenromp EL, Schellenberg JR, et al. (2006) The burden of malaria mortality among African children in the year 2000. *Int J Epidemiol* 35: 691–704.
 59. Ross A, Maire N, Molineaux L, Smith T (2006) An epidemiologic model of severe morbidity and mortality caused by *Plasmodium falciparum*. *Am J Trop Med Hyg* 75: 63–73.
 60. Diallo S, Konate L, Ndir O, Dieng T, Dieng Y, et al. (2000) [Malaria in the central health district of Dakar (Senegal). Entomological, parasitological and clinical data]. *Sante* 10: 221–229.
 61. Greenwood BM, Bradley AK, Greenwood AM (1987) Mortality and morbidity from malaria among children in a rural area of the Gambia, West Africa. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 81: 478–486.
 62. Cisse B, Sokhna C, Boulanger D, Milet J, Ba el H, et al. (2006) Seasonal intermittent preventive treatment with artesunate and sulfadoxine-pyrimethamine for prevention of malaria in Senegalese children: a randomised, placebo-controlled, double-blind trial. *Lancet* 367: 659–667.
 63. Ye Y, Kyobutungi C, Louis VR, Sauerborn R (2007) Micro-epidemiology of *Plasmodium falciparum* malaria: Is there any difference in transmission risk between neighbouring villages? *Malar J* 6: 46.
 64. Schellenberg D, Aponte J, Kahigwa E, Mshinda H, Tanner M, et al. (2003) The incidence in children of clinical malaria detected by active case detection in Ifakara, southern Tanzania. *Trans R Soc Trop Med Hyg* 97: TR/2002/020410.
 65. Saute F, Aponte J, Ahmeda J, Ascaso C, Vaz N, et al. (2003) Malaria in southern Mozambique: incidence of clinical malaria in children living in a rural community in Manhica district. *Trans R Soc Trop Med Hyg* 97: 655–660.
 66. Grobusch MP, Lell B, Schwarz NG, Gabor J, Dornemann J, et al. (2007) Intermittent preventive treatment against malaria in infants in Gabon—a randomized, double-blind, placebo-controlled trial. *J Infect Dis* 196: 1595–1602.
 67. Velema JP, Alihonou EM, Chippaux JP, Van Boxel Y, Gbedji E, et al. (1991) Malaria morbidity and mortality in children under three years of age on the coast of Benin, West Africa. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 85: 430–435.
 68. Lusingu JP, Vestergaard LS, Mmbando BP, Drakeley CJ, Jones C, et al. (2004) Malaria morbidity and immunity among residents of villages with different *Plasmodium falciparum* transmission intensity in North-Eastern Tanzania. *Malar J* 3: 26.
 69. Kobbe R, Kreuzberg C, Adjei S, Thompson B, Langefeld I, et al. (2007) A randomized controlled trial of extended intermittent preventive antimalarial treatment in infants. *Clin Infect Dis* 45: 16–25.
 70. Imbert P, Sartelet I, Rogier C, Ka S, Baujat G, et al. (1997) Severe malaria among children in a low seasonal transmission area, Dakar, Senegal: influence of age on clinical presentation. *Trans R Soc Trop Med Hyg* 91: 22–24.
 71. Imbert P, Gerardin P, Rogier C, Ka AS, Jouvenel P, et al. (2002) Severe falciparum malaria in children: a comparative study of 1990 and 2000 WHO criteria for clinical presentation, prognosis and intensive care in Dakar, Senegal. *Trans R Soc Trop Med Hyg* 96: 278–281.
 72. Camara B, Diouf S, Diagne I, Fall L, Ba A, et al. (2003) Severe malaria in children in a Senegal hospital setting. *Médecine et Maladies Infectieuses* 33: 45–48.
 73. Giha HA, El-Ghazali G, A-Elgadir TME, A-Elbasit IE, Elthahir EM, et al. (2005) Clinical pattern of severe *Plasmodium falciparum* malaria in Sudan in an area characterized by seasonal and unstable malaria transmission. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 99: 243–251.
 74. Taylor T, Olola C, Valim C, Agbenyega T, Kremsner P, et al. (2006) Standardized data collection for multi-center clinical studies of severe malaria in African children: establishing the SMAC network. *Trans R Soc Trop Med Hyg* 100: 615–622.
 75. Jallow M MalariaGEN project.
 76. Rodrigues A, Schellenberg JA, Kofoed PE, Aaby P, Greenwood B (2008) Changing pattern of malaria in Bissau, Guinea Bissau. *Trop Med Int Health* 13: 410–417.
 77. Guiguemde TR, Dabire E, Coulibaly CO, Pare J, Kam LK (1991) Etude de la mortalité palustre dans le service de pédiatrie de l'hôpital de Ouagadougou (Burkina Faso). *Bull Soc Pathol Exot* 84: 338–339.
 78. Schapira A, Solomon T, Julien M, Macome A, Parmar N, et al. (1993) Comparison of intramuscular and intravenous quinine for the treatment of severe and complicated malaria in children. *Trans R Soc Trop Med Hyg* 87: 299–302.
 79. Varandas L, Julien M, Gomes A, Rodrigues P, Van Lerberghe W, et al. (2001) A randomised, double-blind, placebo-controlled clinical trial of vitamin A in severe malaria in hospitalised Mozambican children. *Ann Trop Paediatr* 21: 211–222.
 80. Thuma PE Malaria admissions to Macha Mission Hospital Zambia 2003–2004.
 81. Boulard JC, Chippaux JP, Ayivi B, Akogbeto M, Massougbdji A, et al. (1990) Study of malarial morbidity in a hospital paediatric service in Benin (West Africa) in 1988 and 1989. *Medicine Tropicale* 50: 315–320.
 82. Snow RW, Armstrong Schellenberg JRM, Peshu N, Forster D, Newton CRJC, et al. (1993) Periodicity and space-time clustering of severe childhood malaria on the coast of Kenya. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 87: 386–390.
 83. Slutsker L, Taylor TE, Wirima JJ, Steketee RW (1994) In-hospital morbidity and mortality due to malaria-associated severe anaemia in two areas of Malawi with different patterns of malaria infection. *Trans R Soc Trop Med Hyg* 88: 548–551.
 84. Koko J, Duffillot D, Zima-Ebeyard AM, Duong TH, Gahouma D, et al. (1999) Clinical and epidemiological aspects of malaria in children in Libreville, Gabon. *Medicine d'Afrique Noire* 46: 10–14.
 85. Olanrewaju WI, Johnson AWBR (2001) Malaria in children in Ilorin, Nigeria. *East African Medical Journal* 78: 131–134.
 86. Kazembe LN, Kleinschmidt I, Sharp BL (2006) Patterns of malaria-related hospital admissions and mortality among Malawian children: an example of spatial modelling of hospital register data. *Malar J* 5: 93.
 87. Bassat Q, Guinovart C, Sigauque B, Aide P, Sacarlal J, et al. (2008) Malaria in rural Mozambique. Part II: children admitted to hospital. *Malar J* 7: 37.

88. Maitland K, Levin M, English M, Mithwani S, Peshu N, et al. (2003) Severe *P. falciparum* malaria in Kenyan children: evidence for hypovolaemia. *Qjm* 96: 427–434.
89. Mulumba MP, Ilunga I, Bankoto A, Kamba VD (Submitted) Radioscopic du polymorphisme clinique de l'access de pernicious palustre de l'enfant a Kinshasa. *Congo Medical Journal*.
90. Allen SJ, O'Donnell A, Alexander NDE, Clegg JB (1996) Severe malaria in children in Papua New Guinea. *Qjm* 89: 779–788.
91. Mockenhaupt FP, Ehrhardt S, Burkhardt J, Bosomtwe SY, Laryea S, et al. (2004) Manifestation and outcome of severe malaria in children in northern Ghana. *Am J Trop Med Hyg* 71: 167–172.
92. Alonso PL, Lindsay SW, Armstrong JR, Conteh M, Hill AG, et al. (1991) The effect of insecticide-treated bed nets on mortality of Gambian children. *Lancet* 337: 1499–1502.
93. Greenwood BM, David PH, Otoo-Forbes LN, Allen SJ, Alonso PL, et al. (1995) Mortality and morbidity from malaria after stopping malaria chemoprophylaxis. *Trans R Soc Trop Med Hyg* 89: 629–633.
94. D'Alessandro U, Olalaye BO, McGuire W, Langerock P, Bennett S, et al. (1995) Mortality and morbidity from malaria in Gambian children after introduction of an impregnated bednet programme [see comments]. *Lancet* 345: 479–483.
95. Jaffar S, Leach A, Greenwood AM, Jepson A, Muller O, et al. (1997) Changes in the pattern of infant and childhood mortality in upper river division, The Gambia, from 1989 to 1993. *Trop Med Int Health* 2: 28–37.
96. Trape JF, Pison G, Preziosi MP, Enel C, Desgrees du Lou A, et al. (1998) Impact of chloroquine resistance on malaria mortality. *C R Acad Sci III* 321: 689–697.
97. Etard J-F, Le Hesran J-Y, Diallo A, Diallo J-P, Ndiaye J-L, et al. (2004) Childhood mortality and probable causes of death using verbal autopsy in Niakhar, Senegal, 1989–2000. *International Journal of Epidemiology* 33: 1286–1292(1287).
98. Ghana VAST Study Team (1993) Vitamin A supplementation in northern Ghana: effects on clinic attendances, hospital admissions, and child mortality. *Lancet* 342: 7–12.
99. Becher H, Kynast-Wolf G, Sie A, Ndugwa R, Ramroth H, et al. (2008) Patterns of malaria: cause-specific and all-cause mortality in a malaria-endemic area of west Africa. *Am J Trop Med Hyg* 78: 106–113.
100. Snow RW, Mung'ala VO, Foster D, Marsh K (1994) The role of the district hospital in child survival at the Kenyan Coast. *Afr J Health Sci* 1: 71–75.
101. Delacollette C, Barutwanayo M (1993) [Mortality and morbidity at young ages in a stable hyperendemic malaria region, community Nyanza-Lac, Imbo South, Burundi]. *Bull Soc Pathol Exot* 86: 373–379.
102. Premji Z, Ndayanga P, Shiff C, Minjas J, Lubega P, et al. (1997) Community based studies on childhood mortality in a malaria holoendemic area on the Tanzanian coast. *Acta Trop* 63: 101–109.
103. Schumacher R, Swedberg E, Diallo M (2002) Mortality study in Guinea: investigating the causes of death in children under 5. Published by Save the Children Federation, Inc. and the Basic Support for Institutionalizing Child Survival (BASICS II) Project.
104. Phillips-Howard PA, Nahlen BL, Kolczak MS, Hightower AW, ter Kuile FO, et al. (2003) Efficacy of permethrin-treated bed nets in the prevention of mortality in young children in an area of high perennial malaria transmission in western Kenya. *Am J Trop Med Hyg* 68: 23–29.