Discrimination of Juvenile Yellowfin (*Thunnus albacares*) and Bigeye (*T. obesus*) Tunas using Mitochondrial DNA Control Region and Liver Morphology

Ivane R. Pedrosa-Gerasmio^{1*}✉, Ricardo P. Babaran¹, Mudjekeewis D. Santos²

¹ College of Fisheries and Ocean Sciences, University of the Philippines Visayas, Iloilo, Philippines, ² National Fisheries Research and Development Institute, Quezon City, Philippines

Abstract

Yellowfin tuna, *Thunnus albacares* (Bonnaterre, 1788) and bigeye tuna, *Thunnus obesus* (Lowe, 1839) are two of the most economically important tuna species in the world. However, identification of their juveniles, especially at sizes less than 40 cm, is very difficult, often leading to misidentification and miscalculation of their catch estimates. Here, we applied the mitochondrial DNA control region D-loop, a recently validated genetic marker used for identifying tuna species (Genus *Thunnus*), to discriminate juvenile tunas caught by purse seine and ringnet sets around fish aggregating devices (FADs) off the Southern Iloilo Peninsula in Central Philippines. We checked individual identifications using the Neighbor-Joining Method and compared results with morphometric analyses and the liver phenotype. We tested 48 specimens ranging from 13 to 31 cm fork length. Morpho-meristic analyses suggested that 12 specimens (25%) were bigeye tuna and 36 specimens (75%) were yellowfin tuna. In contrast, the genetic and liver analyses both showed that 5 specimens (10%) were bigeye tuna and 43 (90%) yellowfin tuna. This suggests that misidentification can occur even with highly stringent morpho-meristic characters and that the mtDNA control region and liver phenotype are excellent markers to discriminate juveniles of yellowfin and bigeye tunas.

Citation: Pedrosa-Gerasmio IR, Babaran RP, Santos MD (2012) Discrimination of Juvenile Yellowfin (*Thunnus albacares*) and Bigeye (*T. obesus*) Tunas using Mitochondrial DNA Control Region and Liver Morphology. PLoS ONE 7(4): e35604. doi:10.1371/journal.pone.0035604

Editor: Dirk Steinke, Biodiversity Institute of Ontario - University of Guelph, Canada

Received: December 6, 2011; **Accepted:** March 20, 2012; **Published:** April 19, 2012

Copyright: © 2012 Pedrosa-Gerasmio et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by the Accelerated Science and Technology Human Resource Development (ASTHRD) and Grants-in-Aid (GIA) Programs of the Department of Science and Technology-Philippine Council for Aquatic and Marine Research and Development (DOST-PCAMRD) and the National Fisheries Research and Development Institute (NFRDI), Philippines. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: ivanepedrosa@yahoo.com.ph

✉ Current address: National Fisheries Research and Development Institute, Quezon, Philippines

Introduction

Yellowfin tuna (*Thunnus albacares*) and bigeye tuna (*T. obesus*) are the second and third most important large tuna commodity in the Philippines by catch weight, after skipjack tuna [1,2,3]. Following the introduction of fish aggregating devices (FADs), locally known as payao, their catches increased significantly in the mid 1970s [4] especially for smaller-sized individuals [1]. Reliable estimates of the numbers of these two species are very important in fisheries management to illustrate annual production, demonstrate utilization rates, monitor catch quotas, estimate fishing mortality and to calculate catch per unit effort especially in light of declining population due to overfishing in recent years [2]. However, differentiation of these tuna species in commercial landings poses a problem since the two species are morphologically very similar, especially at sizes less than 40 cm Fork Length (FL). It has been suggested that misidentification by fishery-data collectors can be as high as 30% (Chow and Inoue, 1993). The difficulty in distinguishing these two species, particularly to non-expert field staff has long been problematic in Philippine fisheries statistics as yellowfin and bigeye tuna data were collectively grouped as yellowfin [5].

Mis-identification is not uncommon for yellowfin and bigeye tuna species [6], with the frequency of misidentification as high as 30% [7]. Grewe and Hampton (1998) reported a 0–10.4% frequency of yellowfin among collected bigeye tunas at sizes 40 to 60 cm FL indicating the need for individual genetic identification. Interestingly, in the genetic component of the same study up to 30% of juvenile fish identified in the field as yellowfin tuna were genetically confirmed to be bigeye tuna [8].

Recently, the mitochondrial DNA control region (CR) has been validated as a molecular marker for differentiating *Thunnus* species, and is a more robust marker, than the standard mitochondrial DNA barcode marker, CO1, for differentiating all tuna species including those belonging to the subgenus *Neothunnus* (*Thunnus albacares*, *T. atlanticus*, *T. tonggol*) that are very closely related [9]. In tunas, the mtDNA CR displacement-loop (D-loop) region is highly polymorphic [10]. This does not encode proteins and typically have a high mutation rate presumably due to reduced functional constraints and relaxed selection pressure [11], increasing their likelihood of discriminating between species. In the study of Niwa *et al.* (2003), the genetic variation of the yellowfin tuna mtDNA CR D-loop was shown to be extremely high and a suitable region for investigations of population structure [10].

Figure 1. Neighbor-Joining Analyses of a 298 base pair fragment of the mtDNA CR D-loop. Numbers above the branches indicate bootstrap values inferred from 1000 replicates. Branches corresponding to partitions reproduced in less than 50% bootstrap replicates were collapsed. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. doi:10.1371/journal.pone.0035604.g001

Here, we applied the mtDNA CR D-loop as a marker to differentiate juvenile yellowfin and bigeye tunas, and compared molecular results against identifications based on traditional characters. In addition, the liver phenotypes of the two species [12] were validated for identification of juvenile yellowfin and bigeye tuna. The right, medial and left lobes of the liver measured were analyzed using Principal Component Analysis (PCA).

Results

Species identification based on morpho-meristic characters

External characters such as body coloration, marks and bandings, eye diameter, and body depth have been used to identify and distinguish tunas. Body color is ideal when the specimens are fresh, but colors fade quickly after death. Similarly, bandings and lines can become washed or rubbed out. In this study, the specimens were taken from the landed catch and were stored on ice to maintain colors, markings and bands until examination. Eye [12] diameter and body depth have been used to distinguish the two species but are unreliable in juvenile specimens, since the eye of yellowfin tuna juveniles may appear quite large and indistinguishable from that of bigeye tuna. Body depths are also very similar in juveniles of both species. Takeyama *et al.* (2001) have claimed that there are no external morphological characters for species identification of small juvenile tunas.

In this study, the number of gill rakers in the lower gill arch varied between juveniles (13–31 cm FL) with 18–22 in yellowfin

tuna and 17–21 in bigeye tuna and this character did not provide identification since the ranges overlap.

Using the combined traditional morpho-meristic characters above, 36 juvenile specimens were tentatively identified as yellowfin tuna and 12 specimens as bigeye tuna. These initial identifications were recorded to test the commonality of misidentification of juvenile yellowfin and bigeye tuna.

Species identification based on mtDNA CR D-Loop

Reference control region sequences (397 base pairs) were extracted from the study of Martinez *et al.* (2006, [13]) for *T. albacares* (GenBank Accession Number DQ126342 and DQ126343) and *T. obesus* (GenBank Accession Number DQ126501 and DQ126502). Percent homology or percent identity between *T. albacares* and *T. obesus* was 90% [14].

Of the 48 DNA samples of juvenile tuna examined, Neighbor-Joining analysis identified 43 as yellowfin tuna (90%) and five as bigeye tuna (10%) (Fig. 1). Additional tree building methods (UPGMA, Minimum Evolution, and Maximum Parsimony) generated similar trees.

Species identification based on liver morphology

Whole livers obtained from the 48 specimens showed two distinct phenotypes that corresponded with yellowfin tuna and bigeye tuna (Fig. 2). In yellowfin tuna, the right lobe of the liver is longer than the round medial and left lobes, and the lobes are smooth and clear, with no striations. In bigeye tuna, the three lobes are rounded and about equal in size, with a striated ventral surface [12]. The relative size of the right lobe provided an

Figure 2. Graph Ordination from Principal Component Analysis. Principal Components I and II separated juveniles of yellowfin (Upper; $n=43$) and bigeye (Lower; $n=5$) tunas and comparison of their liver morphologies; yellowfin (A. Itano, 2005 and B. Pedrosa-Gerasmio *et al.*, 2011) and bigeye tuna (C. Itano, 2005 and D. Pedrosa-Gerasmio *et al.*, 2011).

doi:10.1371/journal.pone.0035604.g002

unambiguous separation between the juvenile specimens, but the striations were not obvious.

Principal Component Analysis (PCA) generated a graph ordination (Fig. 2) clearly separating the two species of juvenile tunas based on liver measurement data. The first two eigenvalues are approximately 95% of the total meaningful variance. In general, once eigenvectors are found from the covariance matrix, the next step is to sort them by eigenvalue, in decreasing order which gives the components (Tables 1 and 2) in order of significance [15]. In this case, the first and second components (PCI and PCII) were retained for the analysis, which yielded two

Table 1. Variance extracted from the 3 axes using liver measurement data.

Axis	Eigenvalue	% of Variance	Cum.% of Var.
1	2.42	80.654	80.654
2	0.443	14.772	95.426
3	0.137	4.574	100

doi:10.1371/journal.pone.0035604.t001

axes. Genetic-based identification and liver morphology identification were in complete agreement for all individuals (Table 3).

Discussion

Juveniles of yellowfin tuna and bigeye tuna, especially at sizes less than 40 cm FL, are difficult to distinguish using external morphology while DNA-based methods and liver morphology are more reliable for obtaining species identifications (e.g. [8,16]). With an increasing catch of tuna juveniles, accurate species-level catch data are necessary to determine reproductive activity and to clarify species distributions for fisheries conservation and management [16].

Tuna species can be identified using several genetic markers developed in population-based studies. However, misidentification can occur if the genetic marker is not appropriate for species discrimination [9]. For instance, certain nuclear genetic markers cannot distinguish between Atlantic and Pacific bluefin tuna [17]. Further, genetic markers with low genetic variability, such as mtDNA CO1, infer low genetic distance among *T. albacares*, *T. atlanticus* and *T. tonggol* and prove limited use in differentiation between these species [18]. There is therefore a need to consider several premises before attempting the identification of tuna species using mitochondrial genetic markers. Validation of the genetic marker is even more critical due to the observed introgression in some *Thunnus* species [17]. Recently, the mitochondrial DNA control region has been demonstrated to accurately discriminate all species in the genus *Thunnus* [9].

Here, the use of traditional morphological and meristic characters resulted in misidentification of juvenile tuna about 27% of the time. Alternatively, mtDNA CR D-loop data was highly accurate at discriminating juveniles of yellowfin and bigeye tuna with an unambiguous separation between species of 100%. Furthermore, differentiation of the two juveniles liver morphology using the right-lobe liver criterion [12] was confirmed by genetic data showing 1:1 correspondence; 5 samples (10%) bigeye tunas and 43 samples (90%) yellowfin tunas. This result suggests that liver phenotypes can be a powerful identification tool for fisheries managers on board ships, in the marketplace or in

Table 2. Loadings from a Principal Component Analysis of the log transformed right, middle and left lobe measurements of *T. albacares* and *T. obesus*. Variables with the highest values on principal components I and II (in asterisks) are shown.

Characters	EIGENVECTORS		
	1	2	3
RIGHT	−0.531	0.8455*	−0.054
MIDDLE	−0.6026*	−0.333	0.7254

doi:10.1371/journal.pone.0035604.t002

Table 3. Identification of juvenile tunas caught in Southern Iloilo, Philippines based on body morphology, liver morphology and mtDNA Control Region sequence Data (n = 48).

SAMPLE	FORK	BODY MORPHOLOGY	LIVER	CR D-LOOP	GENBANK
NAME	LENGTH (cm)	INITIAL ID	MORPHOLOGY	ID	ACCESSION
			ID		NUMBER
1. Oc6	12.8	Bigeye	Bigeye	Bigeye	JN988649
2. Oct11	21.4	Yellowfin	Yellowfin	Yellowfin	JN988644
3. Oct9	21.1	Yellowfin	Yellowfin	Yellowfin	JN988643
4. Oc4	16.4	Yellowfin	Yellowfin	Yellowfin	JN988642
5. BET3	31.6	Bigeye*	Yellowfin	Yellowfin	JN988641
6. Oc13	14.8	Yellowfin	Yellowfin	Yellowfin	JN988640
7. Oct6	22.4	Yellowfin	Yellowfin	Yellowfin	JN988639
8. BET2	31.3	Bigeye*	Yellowfin	Yellowfin	JN988638
9. BET4	28.8	Bigeye*	Yellowfin	Yellowfin	JN988637
10. B3	30	Bigeye*	Yellowfin	Yellowfin	JN988636
11. YFT2	30.4	Yellowfin	Yellowfin	Yellowfin	JN988635
12. Oc18	13.7	Yellowfin	Yellowfin	Yellowfin	JN988634
13. Oct18	21.3	Bigeye*	Yellowfin	Yellowfin	JN988633
14. Oct2	22.4	Bigeye*	Yellowfin	Yellowfin	JN988632
15. Oc282	25.5	Yellowfin	Yellowfin	Yellowfin	JN988631
16. Oct19	20.8	Yellowfin	Yellowfin	Yellowfin	JN988630
17. Oc5	17	Yellowfin	Yellowfin	Yellowfin	JN988629
18. Oct16	20.9	Bigeye*	Yellowfin	Yellowfin	JN988628
19. YFT3	29.6	Yellowfin	Yellowfin	Yellowfin	JN988627
20. BET 1	28.8	Bigeye	Bigeye	Bigeye	JN988648
21. Oct12	21.5	Yellowfin	Yellowfin	Yellowfin	JN988626
22. Oct15	22	Yellowfin	Yellowfin	Yellowfin	JN988625
23. N13	20	Yellowfin	Yellowfin	Yellowfin	JN988624
24. N3	17.5	Yellowfin	Yellowfin	Yellowfin	JN988623
25. N17	18.6	Yellowfin	Yellowfin	Yellowfin	JN988622
26. N9	19.5	Yellowfin	Yellowfin	Yellowfin	JN988621
27. Oc19	14.8	Yellowfin	Yellowfin	Yellowfin	JN988620
28. Oc20	18.4	Yellowfin	Yellowfin	Yellowfin	JN988619
29. Oct1	23.2	Yellowfin	Yellowfin	Yellowfin	JN988618
30. N2	19.2	Yellowfin	Yellowfin	Yellowfin	JN988617
31. Oc2650	23.2	Yellowfin*	Bigeye	Bigeye	JN988647
32. N10	18.8	Yellowfin	Yellowfin	Yellowfin	JN988616
33. N1	20.2	Yellowfin	Yellowfin	Yellowfin	JN988615
34. N14	22	Bigeye*	Yellowfin	Yellowfin	JN988614
35. Oct20	22.9	Yellowfin	Yellowfin	Yellowfin	JN988613
36. Oc2611	23	Yellowfin*	Bigeye	Bigeye	JN988646
37. Na23	26	Yellowfin*	Bigeye	Bigeye	JN988645
38. B2	30	Bigeye*	Yellowfin	Yellowfin	JN988612
39. Na24	24.3	Yellowfin	Yellowfin	Yellowfin	JN988611
40. Na26	26.2	Yellowfin	Yellowfin	Yellowfin	JN988610
41. Oc3	15.5	Yellowfin	Yellowfin	Yellowfin	JN988609
42. Oc1	16.6	Yellowfin	Yellowfin	Yellowfin	JN988608
43. Oc8	15.3	Bigeye*	Yellowfin	Yellowfin	JN988607
44. N7	18.7	Yellowfin	Yellowfin	Yellowfin	JN988606
45. Oc11	21.4	Yellowfin	Yellowfin	Yellowfin	JN988605
46. Oc12	13.8	Yellowfin	Yellowfin	Yellowfin	JN988604

doi:10.1371/journal.pone.0035604.t003

Figure 3. Sampling Site. Map of Southeast Asia showing the fish sampling site (off Southern Iloilo, Philippines) with juveniles of yellowfin and bigeye tunas shown.
doi:10.1371/journal.pone.0035604.g003

(NAOAc) and 900 µL of 95% ethanol were then added and mixed with the supernate in new tubes. Tubes were shaken by hand for 3 min and placed overnight in -20°C freezer to allow the DNA to precipitate out. After precipitation, the samples were centrifuged at 12,000 rpm for 10 min. The aqueous phase was carefully pipetted out and the DNA pellet was left at the bottom of each tube. The pellets were then rinsed twice with 500 µL 70% ethanol and centrifuged at 12,000 rpm for 5 min. The tubes were then opened and allowed to air dry for 30 min then rehydrated in 300 µL of $1\times\text{TE}$ buffer. The DNA extracts were then stored at -20°C until molecular analysis.

The mtDNA CR D-loop was amplified from the genomic DNA using the polymerase chain reaction (PCR) technique and two primers (CB3R420 5' CCCCCCTAACTCCCAAAGCTAGG-3' and 12Sar430 5' GCCTGCGGGGCTTTCTAGGGCC3') primarily designed for tuna under the genus *Euthynnus* but also suggested for use in fish closely related to these genus [6]. PCR was carried out in a final volume of 25 µL, in a reaction mixture containing the following reagents: 11.3 µL ddH₂O, 2.5 µL $10\times$ PCR Buffer with 1.5 mM MgCl₂, 5.0 µL 2 mM dNTPs, 2.5 µL 10 µM Primer 1, 2.5 µL 10 µM Primer 2, 0.2 µL *Taq* DNA Polymerase and 1 µL of DNA template. Individual tubes were subjected to the following cycling parameters in a PCR machine (Labnet International, Inc.): initial denaturation phase of 5 min at 94°C , followed by 35 amplification cycles, each consisting of 1 min of denaturation at 94°C , 1 min of annealing at 50°C , and 1 min of extension at 72°C . Final extension was set at 72°C for 5 minutes.

Agarose gel electrophoresis was used to confirm the successful DNA amplification before sending samples for DNA sequencing. A 1% agarose gel was made by suspending dry agarose in a buffer solution (1 g of agarose to 100 mL of $1\times\text{TAE}$ buffer), boiled for approximately 5 minutes or until agarose was completely dissolved, and then poured into a casting tray and allowed it to cool. During electrophoresis, the gel was submerged in a chamber containing a $1\times\text{TE}$ buffer solution. The 2 µL of each PCR product was loaded into individual wells with a 3 µL loading dye. The DNA for analysis was forced through the pores of the gel by the electrical current. Electrode wires were connected to the power supply. Positive (red) and negative (black) were made sure to be properly connected. Under an electrical field, DNA moves to the positive electrode and away from the negative electrode [20]. Electrophoresis was allowed to run for 15 min at 100 volts and was visualized under UV light in a gel documentation system. Unclean DNA samples were sent to Macrogen in Korea for purification and DNA Sequencing using 3730/3730xl DNA Analyzer.

References

- Dickson J, Natividad A (1997) Tuna fishing and a review of payaos in the Philippines. In: Le-Gall JY, Cayre P, Taquet M, eds. Mechanisms and effects of the aggregation of tuna by Fish Aggregating Devices (FADs) Elsevier, Paris, France. pp 141–158.
- Langley A, Williams P, Hampton J (2006) The Western and Central Pacific tuna fishery: 2005 overview and status of stocks. Tuna Fisheries Assessment Report 7. Noumea, New Caledonia: Secretariat of the Pacific Community.
- Barut N, Garvilles E (2010) Annual Report to the Commission Part 1: Information on Fisheries, Research, and Statistics. Scientific Committee Regular Session, Nuku'alofa, Tonga, 6th. Western and Central Pacific Fisheries Commission, Pohnpei, Federated States of Micronesia. 11 p.
- Babaran R (2006) Payao fishing and its impacts to tuna stocks. College of Fisheries and Ocean Sciences, University of the Philippines Visayas. WCPFC-SC2-2006/FT WP-7 1–12 Available: http://www.wcpfc.int/DigitalLibrary/Doc/FAME/Meetings/WCPFC/SC2/SC2_FT_WP7.pdf.
- Langley A, Hampton J (2007) Spatio-temporal patterns of purse seine catches of skipjack and juvenile bigeye and yellowfin tuna caught in association with floating objects. WCPFC-SC3-SA SWG/WP-04. pp 1–10.
- Santos M, Lopez G, Barut N (2010) A pilot study on the genetic variation of Eastern little tuna (*Euthynnus affinis*) in Southeast Asia. Phil J of Science 139(1): 43–50.
- Chow S, Inoue S (1993) Intra and Interspecific restriction fragment length polymorphism in mitochondrial genes of *Thunnus* tuna species. Nat Res Inst Far Seas Fish Bull 30: 207–225.
- Grewe P, Hampton J (1998) An Assessment of Bigeye (*Thunnus obesus*) population structure in the Pacific Ocean, based on mitochondrial DNA and DNA microsatellite analysis. SOEST Publication 98-05, JIMAR Contribution. 98-320, 23.
- Viñas J, Tudela S (2009) A Validated Methodology for Genetic Identification of Tuna Species (Genus *Thunnus*). PLoS ONE 4(10): e7606. 10.1371/journal.pone.0007606.
- Niwa Y, Nakazawa A, Margulies D, Scholey V, Wexler J, et al. (2003) Genetic monitoring for spawning ecology of captive Yellowfin Tuna (*Thunnus albacares*) using mitochondrial DNA variation. Aquaculture 218: 387–395.
- Liu ZJ, Cordes JF (2004) DNA marker technologies and their applications in aquaculture genetics. Aquaculture 238: 1–37.
- Itano D (2005) A handbook for the identification of Yellowfin and Bigeye Tunas in Fresh Condition. Pelagic Fisheries Research Program, Honolulu, Hawaii, USA. Ver 2. 1–27. Available: BE-YF ID Fresh_ENGLISH_v2.pdf.
- Martínez P, González EG, Castilho R, Zardoya R (2006) Genetic diversity and historical demography of Atlantic bigeye tuna (*Thunnus obesus*). Mol Phylogenet Evol 39: 404–416.

Resulting DNA sequences were edited and aligned using alignment explorer MEGA 4.0 [21] and ClustalX 2.0.11 [22]. Percent homology or percent identity between the two species was obtained using Align software [14]. Evolutionary distances were computed using the Maximum Composite Likelihood method [23] and reported as the number of base substitutions per site. All positions containing gaps and missing data were eliminated (complete deletion option) from the dataset. A neighbor-joining phylogenetic tree was inferred using MEGA4 [21] with 1000 bootstrap probability replicates.

All sequences were deposited in GenBank with Accession Numbers JN988602–JN988644 for *T. albacares* and JN988645–JN988649 for *T. obesus*.

Liver Morphology Analysis

Whole livers were also investigated for each of the samples. These were photographed using a digital camera and the lobes were measured on the longest axis using the Pixel Caliper (version 1.0, UPV, Iloilo, Philippines). Measurement data of the right, left and middle lobes of juvenile yellowfin and bigeye tunas were then log transformed for PCA using PC-ORD 4.10 [24].

Quantifiable difference in the length of the right lobe and overall texture of the liver was noted for each individual. Assumptions were done using the criteria of Itano (2005) for larger individuals, i.e., the right lobe is longer than round medial and left lobes in yellowfin and three rounded lobes are about equal sizes for bigeye. The texture of the livers was also distinct for larger individuals [12], i.e., bigeye livers have striated ventral surface while yellowfins have smooth, clear lobes, and with no striations.

Acknowledgments

The authors would like to thank Roselyn D. Aguila, Minerva H. Ventolero, Sweedy Kay L. Perez, Benedict A. Maralit and Altair B. Agmata of the National Fisheries Research and Development Institute-Genetic Fingerprinting Laboratory (NFRDI-GFL) who made significant contributions for the completion of this study; to Dr. Maria Rowena Romana-Eguia, Prof. Encarnacion Emilia S. Yap, Dr. Rex Ferdinand M. Traifalgar and Dr. Demian A. Willette for their valuable comments and suggestions. We also thank NFRDI for letting us use the equipment in the laboratory as well as providing us some chemicals, laboratory space, and utilities.

Author Contributions

Performed the experiments: IRPG MDS RPB. Analyzed the data: IRPG. Contributed reagents/materials/analysis tools: RPB MDS. Wrote the paper: IRPG RPB.

14. Pearson W, Wood T, Zhang Z, Miller W (1997) Comparison of DNA sequences with protein sequences. *Genomics* 46: 24–36.
15. Smith L (2002) A tutorial on Principal Component Analysis. Available: http://www.cs.otago.ac.nz/cosc453/student_tutorials/principal_components.pdf.
16. Takeyama H, Chow S, Tsuduki H, Matsunaga T (2001) Mitochondrial DNA sequence variation within and between *Thunnus* tuna species and its application to species identification. *J of Fish Biol* 58: 1646–1657.
17. Chow S, Nakagawa T, Suzuki N, Takeyama H, Matsunaga T (2006) Phylogenetic relationships among *Thunnus* species inferred from rDNA ITS1 sequence. *J Fish Biol* 68: 24–35.
18. Ward RD, Zemlak TS, Innes BH, Last PR, Hebert PD (2005) DNA Barcoding of Australia's fish species. *Phil Trans R Soc B* 360: 1847–1857.
19. NOAA (2009) National Oceanic and Atmospheric Administration. A Guide to the Tunas of the Western Atlantic Ocean. Available: http://www.nmfs.noaa.gov/sfa/hms/brochures/2009/Tuna_Guide_2009.pdf.
20. Basit S (2009) Comparison of DNA extraction methods for muscle tissues of tuna and tuna-like species: Its implications for identification using polymerase chain reaction. *Emilio Aguinaldo Coll Res Bull* 7(1): 1–8.
21. Tamura K, Dudley J, Nei M, Kumar S (2007) MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Molecular Biology and Evolution* 24: 1596–1599.
22. Larkin M, Blackshields G, Brown N, Chenna R, McGettigan P, et al. (2007) Clustal W and Clustal X version 2.0. *Bioinformatics* 23: 2947–2948.
23. Tamura K, Nei M, Kumar S (2004) Prospects for inferring very large phylogenies by using the neighbor-joining method. *PNAS* 101: 11030–11035.
24. McCune B, Mefford M (1999) *Multivariate Analysis of Ecological Data*. Version 4.10. MjM Software, Glenden Beach, Oregon. USA.