

Age and Gender Differences in the Social Patterning of Cardiovascular Risk Factors in Switzerland: The CoLaus Study

Silvia Stringhini^{1*}, Brenda Spencer¹, Pedro Marques-Vidal¹, Gerard Waeber², Peter Vollenweider², Fred Paccaud¹, Pascal Bovet¹

1 Institute of Social and Preventive Medicine (IUMSP), Lausanne University Hospital, Lausanne, Switzerland, **2** Department of Medicine, Internal Medicine, CHUV and Faculty of Biology and Medicine, Lausanne, Switzerland

Abstract

Objectives: We examined the social distribution of a comprehensive range of cardiovascular risk factors (CVRF) in a Swiss population and assessed whether socioeconomic differences varied by age and gender.

Methods: Participants were 2960 men and 3343 women aged 35–75 years from a population-based survey conducted in Lausanne, Switzerland (CoLaus study). Educational level was the indicator of socioeconomic status used in this study. Analyses were stratified by gender and age group (35–54 years; 55–75 years).

Results: There were large educational differences in the prevalence of CVRF such as current smoking (Δ =absolute difference in prevalence between highest and lowest educational group:15.1%/12.6% in men/women aged 35–54 years), physical inactivity (Δ =25.3%/22.7% in men/women aged 35–54 years), overweight and obesity (Δ =14.6%/14.8% in men/women aged 55–75 years for obesity), hypertension (Δ =16.7%/11.4% in men/women aged 55–75 years), dyslipidemia (Δ =2.8%/6.2% in men/women aged 35–54 years for high LDL-cholesterol) and diabetes (Δ =6.0%/2.6% in men/women aged 55–75 years). Educational inequalities in the distribution of CVRF were larger in women than in men for alcohol consumption, obesity, hypertension and dyslipidemia ($p<0.05$). Relative educational inequalities in CVRF tended to be greater among the younger (35–54 years) than among the older age group (55–75 years), particularly for behavioral CVRF and abdominal obesity among men and for physiological CVRF among women ($p<0.05$).

Conclusion: Large absolute differences in the prevalence of CVRF according to education categories were observed in this Swiss population. The socioeconomic gradient in CVRF tended to be larger in women and in younger persons.

Citation: Stringhini S, Spencer B, Marques-Vidal P, Waeber G, Vollenweider P, et al. (2012) Age and Gender Differences in the Social Patterning of Cardiovascular Risk Factors in Switzerland: The CoLaus Study. PLoS ONE 7(11): e49443. doi:10.1371/journal.pone.0049443

Editor: M Maria Glymour, Harvard School of Public Health, United States of America

Received: May 7, 2012; **Accepted:** October 7, 2012; **Published:** November 13, 2012

Copyright: © 2012 Stringhini et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: SS is supported by a post-doctoral fellowship awarded by the Swiss School of Public Health (SSPH+). PV and GW received an unrestricted grant for GSK to build the CoLaus study. The CoLaus study was supported by research grants from GlaxoSmithKline, the University Hospital Center, the Faculty of Biology and Medicine of Lausanne, Switzerland and the Swiss National Science Foundation (grant no: 33CSO-122661). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: GSK provided grants for the CoLaus study. This does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials, as detailed online in the guide for authors.

* E-mail: silvia.stringhini@chuv.ch

Introduction

In high income countries, cardiovascular disease (CVD) disproportionately affects the lower socioeconomic groups [1], probably reflecting an unequal distribution of cardiovascular risk factors (CVRF) across society [2,3,4] and differential access to and/or use of treatment [5]. However, the magnitude of socioeconomic inequalities in relation to CVD mortality differs substantially between countries [6,7]. In Europe, there appears to be a North-South gradient in socioeconomic inequalities in CVD, with larger differences in Northern than in Southern European countries [7].

Between-country variations in the magnitude of socioeconomic inequalities in CVD tend to mirror cross-country differences in the social patterning of CVRF. Indeed, strong socioeconomic inequalities in CVRF have frequently been reported in Northern European regions such as in Scandinavian countries or in the United Kingdom [8,9,10], while in several Southern European countries such as Italy, Greece or Spain the association between socioeconomic indicators and CVRF seems to be weaker [11,12,13,14]. For example, Schroder et al. [14] and de Vogli et al. [13] reported a lack of educational/occupational differences in CVRF in Spain and Italy, respectively. Stringhini et al. [15] showed large occupational inequalities in the prevalence of unhealthy behaviors among British civil servants but small inequalities among French employees of the national gas and electricity company. Cavelaars et al. [2] noted a North-South pattern in the social distribution of smoking and vegetable consumption with small associations with educational level in Southern European regions.

ities in CVRF have frequently been reported in Northern European regions such as in Scandinavian countries or in the United Kingdom [8,9,10], while in several Southern European countries such as Italy, Greece or Spain the association between socioeconomic indicators and CVRF seems to be weaker [11,12,13,14]. For example, Schroder et al. [14] and de Vogli et al. [13] reported a lack of educational/occupational differences in CVRF in Spain and Italy, respectively. Stringhini et al. [15] showed large occupational inequalities in the prevalence of unhealthy behaviors among British civil servants but small inequalities among French employees of the national gas and electricity company. Cavelaars et al. [2] noted a North-South pattern in the social distribution of smoking and vegetable consumption with small associations with educational level in Southern European regions.

These North-South differences might be explained by the fact that CVRF were originally more prevalent in the higher socioeconomic groups and the direction of this association has gradually reversed over the last century [16,17]. The “social transition” of CVRF from the higher to the lower socioeconomic groups appears to have started earlier in Northern than in Southern Europe, and to have occurred in men before women [18]. In some Southern European countries certain CVRF such as smoking (among women) or low consumption of fruit and vegetables are still more prevalent in the higher socioeconomic groups [11,19,20]. For example, Huisman et al. reported large educational differences in current smoking in both Northern and Southern Europe, but in Italy, Spain, Greece and Portugal the socioeconomic gradient in women was inverted, the prevalence of smoking being higher among higher educated women [10]. However, most studies examining the social patterning of CVRF in Southern European countries, including Switzerland, are based on data from the 1990's [2,11,14,19,21,22].

In the French-speaking region of Switzerland, the most recent comprehensive assessment of social inequalities in CVRF dates back to the early 2000s [22]. It showed small but significant socioeconomic differences in the prevalence of several CVRF such as current smoking, physical inactivity, obesity and hypertension (but not hypercholesterolemia) among men. Among women, a similar pattern was observed, but current smoking was not socially patterned. More recent studies examining only one risk factor at a time reported decreasing educational inequalities in smoking [23], but increasing educational differences in overweight and obesity [24].

The overall aim of our study is to provide an updated and comprehensive assessment of social inequalities in major risk factors for lifestyle-related diseases (current smoking, heavy drinking, physical inactivity, overweight and obesity, hypertension, dyslipidemia and diabetes) in a French-speaking Swiss town. As the French-speaking region of Switzerland is generally assimilated to Southern European countries for its CVD profile [25], this study allows assessing whether it is still the case that social inequalities in major CVRF are small in Southern Europe. A key feature of this study is that it additionally examines whether socioeconomic differences in CVRF vary by age and gender.

Data and Methods

Study Population and Design

The Colaus study is a cross-sectional population-based study conducted in Lausanne, Switzerland (approximately 180'000 inhabitants). Details of the study have been previously described [26]. Briefly, a simple random sampling of 19,830 participants was drawn, corresponding to 35% of the source population, of which 6738 participants were eventually included. The following inclusion criteria applied: (a) written informed consent; (b) age 35–75 years; (c) willingness to take part in the examination and donate a blood sample; and (d) Caucasian origin. Recruitment began in June 2003 and ended in May 2006. The age and sex distribution of the 6738 participants included in the Colaus study were similar to those of the 19,830 individuals originally sampled. Participants attended the outpatient clinic at the University Hospital of Lausanne (CHUV) in the morning after an overnight fast. Data were collected by trained field interviewers during a single visit lasting about 60 minutes. Venous blood samples were drawn after an overnight fast, and assays were performed by the CHUV Clinical Laboratory on fresh plasma samples within 2 hour of blood collection in a Modular P apparatus (Roche Diagnostics, Switzerland). Information on demographic data, socioeconomic

and marital status, lifestyle factors, personal and family history of disease, CVRF and treatment was collected. The study was approved by the Institutional Ethics Committee of the University of Lausanne (Switzerland).

Measures

Socioeconomic status (SES). *Education* was the indicator of socioeconomic status used in this study. It was assessed as the highest qualification achieved and categorized as “high” (tertiary education), “middle” (upper secondary education or post-secondary non tertiary education, including vocational education) and “low” (lower secondary education or lower) [27].

Cardiovascular risk factors (CVRF). *Current smoking* was assessed using questions on current smoking status and was classified as yes/no. Former smokers were included in the non-smokers category. For current smokers, the *number of pack years* of smoking was calculated by multiplying the number of packs of cigarettes smoked per day (average number of cigarettes smoked per day divided by 20) by the number of years the person reported to have smoked. *Alcohol consumption* was assessed using questions on the number of alcoholic drinks consumed in the past week, then categorized as “abstainers” (0 unit/week), “moderate drinkers” (1–21/1–14 units/week for men/women) or “heavy drinkers” ($\geq 21/\geq 14$ units/week for men/women). We considered both abstaining from alcohol and heavy drinking as CVRF. Participants were classified as *physically active* if they reported participating in a physical activity of more than 20 minutes once a week or more, and as physically inactive otherwise. Body weight and height were measured with participants standing without shoes in light indoor clothing. Body weight was measured in kilograms to the nearest 0.1 kg using a Seca® Scale (Hamburg, Germany), which was calibrated regularly. Height was measured to the nearest 5 mm using a Seca® height gauge (Hamburg, Germany). Waist circumference was measured twice with a non-stretchable tape over the unclothed abdomen at the mid-point between the lowest rib and the iliac crest. The mean of the two measurements was used for analyses [26]. *Body Mass Index* (BMI) was calculated and categorized in three groups (normal <25; overweight 25–29; obese ≥ 30 kg/m²) based on the World Health Organization recommendations [28]. *Abdominal obesity* was considered as a waist circumference ≥ 102 cm for men and ≥ 88 cm for women. Blood pressure (BP) was measured three times on the left arm after at least 10 minutes of rest in a seated position using a clinically validated automated oscillometric device (Omron HEM-907, Matsusaka, Japan) with a cuff adapted to the arm circumference. Three readings were obtained and the average of the last two BP readings was used. *Hypertension* was defined as systolic/diastolic BP $\geq 140/90$ mmHg or use of antihypertensive medication. *Low HDL-cholesterol* was defined for values <1.0 mmol/l in men and <1.2 mmol/l in women; *high LDL-cholesterol* for a value ≥ 3.4 mmol/l; *high triglycerides* for a value ≥ 1.7 mmol/l. *Diabetes* was defined as fasting plasma glucose ≥ 7.0 mmol/L or glucose lowering treatment.

Other covariates. *Place of birth* was classified as “born in Switzerland” or “not born in Switzerland”.

Statistical Analysis

Statistical analysis was conducted using Stata v.12 (Stata corp, College Station, TX, USA). With the few exceptions mentioned below, all analyses were performed separately for men and women and in two age groups (35–54 years and 55–75 years). We used least squares regression to calculate age and place of birth-adjusted prevalence rates or mean values of CVRF for each educational group. Differences in CVRF prevalence and mean values between

the lowest and the highest educational group, with their 95% confidence intervals (CI), were also calculated. As suggested in previous studies [29,30], relative inequalities in CVRF were examined using the Relative Index of Inequality (RII) calculated by log-binomial regression [31]. The RII is a regression-based index taking into account both the size and relative position of each educational group in the educational hierarchy. To compute the RII, education was transformed into a summary measure ranging from zero (highest level of education) to one (lowest level of education). The population in each educational category was assigned a score corresponding to the midpoint of the relative position of their category in the cumulative population distribution. For example, if the highest educational category comprises 24% of the population, all participants in this category are assigned a value of 0.12 (0.24/2), and if the second category comprises 30% of the population, the corresponding value is 0.27 (0.12+ [0.3/2]), and so forth. The RII was calculated using log-binomial regression, as the RII by logistic regression has been shown to produce biased estimates of relative inequalities when the prevalence of the health outcome is relatively high (i.e.: >10%) [30]. As such, the RII can be interpreted as the prevalence ratio between the two ends of the educational hierarchy [30]. Log-binomial regressions were adjusted for age (treated as a continuous variable) and place of birth. Analyses including HDL-cholesterol were additionally adjusted for oral contraceptive intake among women. In order to test whether the associations between education and CVRF differed by gender or by age, interaction terms between education (lowest versus highest education in analysis of absolute inequalities and RII in analysis of relative inequalities) and sex or between education and age group were included in the different regression models described.

Results

From the initial 6738 participants, 435 (6% of the original sample) were excluded because of missing values on one or more covariates (N = 18 for education, N = 157 for alcohol consumption, N = 114 for physical inactivity and N < 20 for the other CVRF, categories not mutually exclusive). Hence, 6303 participants (53% women) were included in the present analyses. Excluded women were slightly older than those included in the study (p = 0.03), but there were no age differences between included and excluded men. Excluded participants were more likely to have CVRF than those included in the analysis (for example, OR = 1.54; 95%CI: 1.25; 1.90 for smoking, OR = 1.29; 95%CI: 1.00; 1.66 for obesity and OR = 2.14; 95%CI: 1.55; 2.94) and they were also more likely to be in the lowest educational group than those included in the study (OR = 1.69; 95%CI: 1.27; 2.27). However, educational inequalities in CVRF were similar in both the excluded and included samples (p for interaction between education and inclusion status > 0.05 for smoking, obesity or diabetes).

Table 1 shows the characteristics of the participants included in the study. Mean age was 52 years for both men and women. One half of participants reported “lower than secondary” education (52.8% of men and 57.8% of women). The distribution of participants across educational categories was similar in the two age groups for men, while women in the older age group tended to report a lower educational level than those in the younger group. The majority of men and women were born in Switzerland.

Absolute Inequalities in CVRF

For men, age and place of birth-adjusted prevalence and mean values of CVRF by educational level and age group are presented

in **Table 2**. Lower education was associated with higher levels of CVRF with a marked dose-response pattern (p for linear trends < 0.05 for all CVRF apart from LDL-cholesterol in the younger age group and alcohol consumption, LDL and HDL-cholesterol in the older age group). There was a 15% (95%CI: 10.0; 20.2) difference in the prevalence of smoking between the lowest and the highest educational group in the youngest age group, but there were no significant educational differences in smoking prevalence in the oldest age group [Δ = 3.1% (95%CI: -3.0; 9.2)]. In both age groups, the number of pack-years smoked increased with decreasing educational level. Physical inactivity, overweight, obesity and abdominal obesity were also far more prevalent in the lowest as compared with the highest educational group (Δ = 25.3%/19.4% in the youngest/oldest age group for physical inactivity; 14.7%/12.5% for overweight; 8.6%/14.2% for obesity and 9.3%/4.6% for abdominal obesity). Large differences were also seen for hypertension (particularly in the oldest age group (Δ = 16.7%)), but less so for dyslipidemia and diabetes. Absolute educational differences in CVRF tended to be larger in the younger than in the older age group for smoking, heavy drinking, physical inactivity and abdominal obesity, but they were larger in the older age group for obesity and hypertension.

For women, the prevalence and mean values of CVRF according to educational level and age group are presented in **Table 3**. As for men, most CVRF showed a linear association with educational level (p for linear trends < 0.05 for all CVRF apart from heavy drinking in the younger age group and smoking, diastolic blood pressure, HDL and LDL-cholesterol and diabetes in the older age group). In the younger age group, but not in the older, large absolute inequalities were observed for current smoking (Δ = 12.6%). Physical inactivity (Δ = 22.7%/21.2% in the younger/older age group), overweight (Δ = 22.9%/27.9%), obesity (Δ = 10.3%/14.8%), abdominal obesity (Δ = 15.7%/21.6%), hypertension (Δ = 8.6%/11.4%) and dyslipidemia (Δ = 9.5%/7.8% for high LDL-cholesterol) were more prevalent in the lowest educational group in both age groups.

Relative Inequalities in CVRF

Results for relative educational inequalities in CVRF are shown in **Table 4**. Participants at the bottom end of the educational hierarchy were more likely to be current smokers than those at the top, but in analysis stratified by age group the association of smoking status with education was evident only in the younger age group (p for interaction between education and age group < 0.05). In general, relative educational inequalities in CVRF were larger in the younger age group (although interaction terms reached statistical significance only for smoking, heavy drinking and abdominal obesity in men and for smoking, hypertension, and dyslipidemia in women). For example, men at the bottom of the educational hierarchy were more than four times more likely to have diabetes than those at the top in the younger age group [RII = 4.61 (95%CI: 1.62; 13.10)], but only 1.9 times more likely in the older age group [RII = 1.89 (95%CI: 1.10; 3.27)]. The corresponding figure for women was RII = 5.12 (95%CI: 0.84; 31.33) among those 35–54 years and RII = 1.87 (95%CI: 0.71; 4.88) among those 55–75 years. Among women, relative inequalities were particularly strong for obesity [RII = 4.77 (95%CI: 3.15; 7.22)] and low HDL-cholesterol [RII = 5.62 (95%CI: 2.64; 12.94)]. Educational inequalities in alcohol abstinence, hypertension and dyslipidemia in the younger age group and in abdominal obesity in the older age group were larger in women than in men (all p < 0.05).

Table 1. Socio-demographic characteristics by gender and age group.

	MEN			WOMEN		
	Overall	35–54 years	55–75 years	Overall	35–54 years	55–75 years
N (%)	2960 (47.0)	1727 (58.3)	1233 (41.7)	3343 (53.0)	1842 (55.1)	1501 (44.9)
Age (mean, SD)	52.2 (10.8)	44.3 (5.3)	63.1 (5.8)	52.9 (10.7)	44.6 (5.4)	63.0 (5.7)
Education, N (%)						
Tertiary	710 (24.0)	463 (26.8)	247 (20.0)	550 (16.5)	396 (21.5)	154 (10.3)
Post secondary/secondary	685 (23.2)	391 (22.6)	294 (23.8)	859 (25.7)	504 (27.4)	355 (23.7)
Lower than secondary	1565 (52.8)	873 (50.6)	692 (56.2)	1934 (57.8)	942 (51.1)	992 (66.0)
Born in Switzerland, N (%)						
Yes	1765 (59.6)	910 (52.7)	855 (69.3)	2035 (60.9)	1016 (55.2)	1019 (67.9)
No	1195 (40.4)	817 (47.3)	378 (30.7)	1308 (39.1)	826 (44.8)	482 (32.1)

SD: standard deviation.

doi:10.1371/journal.pone.0049443.t001

Sensitivity Analyses

About 40% of participants were not born in Switzerland. As education can have different meanings in different populations, depending on the school system and the level of economic development, we repeated the analyses for relative inequalities in CVRF stratifying for place of birth (Switzerland or not Switzerland). In general, there were no substantial differences in educational inequalities in CVRF by place of birth (**Table S1**). However, inequalities in heavy drinking and diabetes were larger among men not born in Switzerland, and inequalities in obesity were larger among women born in Switzerland (all $p < 0.05$). About 6% of participants had missing values on one or more covariates. As missingness was found to be patterned by education, we assessed whether missing data could have biased our results. Analyses for relative educational inequalities in CVRF were rerun using multiple multivariate imputation (STATA procedures “ice/micombine”) to replace missing values. Results did not differ from those reported in the main analysis. Although socioeconomic status is a complex concept, we focused on educational level in this study. However, analyses were also performed using occupational position as the indicator of SES for the 4512 participants who were currently working. Overall, results were very similar to those using education as an indicator of SES. However, in general socioeconomic differences in CVRF tended to be more pronounced for education than for occupational position, especially among women. Finally, we repeated all analyses adjusting for marital status and results were virtually unchanged. All results from sensitivity analysis not shown in Table S1 are available upon request.

Discussion

We found large absolute differences in the prevalence of CVRF according to educational level in this Swiss population. Moreover, relative inequalities by education differed by gender and tended to be greater in the younger than in the older age group, particularly for behavioral risk factors among men and for physiological risk factors among women.

Overall Prevalence of CVRF

Prevalence estimates of smoking, physical inactivity, obesity, and hypertension in our study were comparable to other population-based estimates (several of them telephone health surveys) of the Swiss general population [23,32,33,34]. On the

other hand, the prevalence of measured hypercholesterolemia and diabetes in Colaus was higher than self-reported prevalence from the Swiss health surveys [35,36], or than that measured in the neighboring region of Geneva [22,37]. The prevalence of overweight and obesity, hypertension, high triglycerides and diabetes was higher in men than in women, as reported previously in Switzerland [38].

Absolute Educational Differences in CVRF

Overall, the prevalence of CVRF was lower in higher socioeconomic groups, consistent with general findings in high income countries [2,3,4]. Absolute socioeconomic differences were particularly large for behaviors such as smoking and physical activity, and anthropometric measures such as weight (reflecting the balance between physical activity and diet). This was particularly true among women. This result is in line with findings from recent studies reporting strong educational inequalities in physical inactivity and obesity in Switzerland [32,39]. Absolute educational differences were also large for hypertension but smaller for dyslipidemia and diabetes, as observed previously [40].

Relative Educational Differences in CVRF

Relative educational inequalities differed by age and gender for several CVRF. Among both men and women, the educational gradient in current smoking was stronger in the younger than in the older age group. It has been observed that the smoking epidemic initially spread in the high socioeconomic groups, later reached the lower socioeconomic groups, and started declining first in the high socioeconomic group [18,21]. In addition, the “social transition” of smoking usually starts earlier in men than in women, and in Europe it was delayed in Southern Europe as compared with Northern Europe [18]. In a study conducted in Geneva (Switzerland) in the 1990s, smoking was still more prevalent among the higher educated women [19], but there were no educational differences in current smoking among young participants (35–44 years in 1993–95). The current study suggests that the social transition in smoking is now completed in Switzerland.

Gender and age differences were also observed for the association between education and heavy drinking. Low-educated young men were more likely to report heavy drinking than young men with high education, but the inverse was observed among older men. Among women, heavy drinking tended to be more

Table 2. Age-adjusted prevalence and mean values of selected cardiovascular risk factors by level of education and age group among men (N = 2960).

	Age group 35–54 years (N = 1727)							Age group 55–75 years (N = 1233)						
	Educational level							Educational level						
	Overall	High	Mid	Low	<i>p</i> ^a	<i>f</i> ^b	(95% CI)	High	Mid	Low	<i>p</i> ^a	<i>f</i> ^b	(95% CI)	<i>P</i> ^c
Health behaviors														
Current smoking (%)	29.4	21.6	29.1	36.7	0.000	15.1	(10.0; 20.2)	22.7	24.3	25.8	0.314	3.1	(−3.0; 9.2)	0.004
Pack-years (mean) ^d	24.2	13.7	18.8	23.0	0.000	9.2	(5.7; 12.7)	19.2	25.8	34.0	0.001	15.0	(6.6; 26.5)	0.341
No alcohol consumption (%)	18.0	16.1	19.1	22.1	0.008	6.0	(1.6; 10.4)	13.7	14.3	14.9	0.650	1.2	(−3.8; 6.1)	0.117
Heavy drinking (%)	12.2	6.1	8.9	11.8	0.001	5.7	(2.4; 9.0)	15.8	15.1	14.4	0.575	−1.4	(−6.5; 3.6)	0.009
Physical inactivity (%)	36.6	20.2	32.8	45.5	0.000	25.3	(20.2; 30.5)	21.9	31.6	41.3	0.000	19.4	(12.8; 26.1)	0.111
Body mass index (mean, kg/m ²)														
Overweight ^e (%)	62.0	45.2	52.6	59.9	0.000	14.7	(9.3; 20.2)	62.4	68.6	74.9	0.000	12.5	(6.1; 18.9)	0.546 <i>546</i>
Obesity ^e (%)	17.1	6.4	10.7	15.1	0.000	8.6	(5.1; 12.2)	12.8	19.9	27.0	0.000	14.2	(8.3; 20.1)	0.126
Waist circumference (mean, cm)														
Abdominal obesity ^e (%)	26.6	10.8	15.5	20.2	0.000	9.3	(5.2; 13.4)	34.8	37.1	39.4	0.186	4.6	(−2.2; 11.5)	0.220
Blood pressure (mmHg)														
Systolic (mean)	131.8	124.7	126.1	127.5	0.000	2.9	(1.4; 4.3)	135.3	137.8	140.2	0.000	4.9	(2.4; 7.3)	0.106
Diastolic (mean)	81.2	78.8	79.8	80.8	0.001	2.0	(0.8; 3.1)	81	82.1	83.3	0.003	2.3	(0.8; 3.9)	0.710
Hypertension ^e (%)	37.2	18.3	21.9	25.4	0.002	7.1	(2.5; 11.7)	43.5	51.8	60.2	0.000	16.7	(9.8; 23.6)	0.023
HDL-cholesterol (mean, mmol/l)														
Low HDL-cholesterol ^e (%)	4.2	3.1	4.5	5.9	0.020	2.8	(0.4; 5.2)	2.8	3.1	3.3	0.650	0.6	(−1.9; 3.1)	0.182
LDL-cholesterol (mean, mmol/l)														
High LDL-cholesterol ^e (%)	49.9	47.5	49.4	51.4	0.162	4.0	(−1.6; 9.6)	49.9	50.8	51.7	0.623	1.8	(−5.3; 8.8)	0.366
Triglycerides (mean, mmol/l)														
High triglycerides ^e (%)	32.5	26.4	29.5	32.7	0.015	6.4	(1.2; 11.5)	28.7	32.6	36.5	0.023	7.8	(1.1; 14.5)	0.994
Fasting glucose (mean, mmol/l)														
Diabetes ^e (%)	8.9	1.5	3.1	4.7	0.003	3.1	(1.1; 5.2)	11.1	14.1	17.1	0.021	6.0	(0.9; 11.0)	0.253

BP: blood pressure; CI: confidence interval; HDL: high-density lipoprotein; LDL: low-density lipoprotein. Prevalence and mean values are adjusted for age and place of birth (Switzerland or outside Switzerland).

^a*p* for linear trend across socioeconomic categories.

^bDifference in prevalence/mean between the highest and the lowest educational category.

^c*p* for interaction between educational level and age group.

^dAnalyses restricted to current smokers (N = 515 in the 35–54 years group and N = 293 in the 55–75 years group). 32 smokers with missing information on pack-years were not included.

^eOverweight: BMI ≥ 25 kg/m² and < 30 kg/m²; obesity: BMI ≥ 30 kg/m²; abdominal obesity: waist circumference $\geq 102/88$ cm in men/women; hypertension: BP $\geq 140/90$ mmHg or taking BP treatment; low HDL-cholesterol: $< 1.0/1.2$ mmol/l in men/women; high LDL-cholesterol: ≥ 3.4 mmol/l; high triglycerides: ≥ 1.7 mmol/l; diabetes: fasting glucose ≥ 7.0 mmol/l or taking diabetes treatment.

doi:10.1371/journal.pone.0049443.t002

common in the higher educational group, as observed previously in Switzerland and other European countries [39,41], but the associations were not statistically significant. Relative inequalities in physical inactivity were very large but did not differ by age and gender. Conversely, the social patterning of obesity was stronger in women, as previously reported [39,42]. This was mostly due to the very low prevalence of obesity among the highly educated women. It has been hypothesized that this might reflect a stronger social pressure for thinness on women with a high socioeconomic status than on women with a low socioeconomic status, in addition to greater health consciousness [43].

Hypertension was also strongly socially patterned, as reported previously in Switzerland [22]. Although absolute inequalities in dyslipidemia and diabetes were not large, relative inequalities were strong in Lausanne compared with other countries [44]. For example, young women with a low educational level were more than 10 times more likely to have low HDL-cholesterol and 5 times more likely to have diabetes than their more advantaged counterparts. This might be related to the observed inequalities in physical inactivity and obesity among younger women.

For most CVRF and for both genders, relative educational inequalities were stronger in the younger (35–54 years) than in the older (55–75 years) age group. This could either mirror cohort

Table 3. Age-adjusted prevalence and mean values of selected cardiovascular risk factors by level of education and age group among women (N = 3343).

	Age group 35–54 years (N = 1842)							Age group 55–75 years (N = 1501)						
	Educational level						<i>p</i> ^a	Educational level						<i>p</i> ^c
	Overall	High	Mid	Low	<i>p</i> ^b	(95% CI)		High	Mid	Low	<i>p</i> ^b	(95% CI)		
Health behaviors														
Current smoking (%)	23.6	20.4	26.7	33	0.000	12.6	(7.5; 17.8)	20.3	19.9	19.6	0.813	−0.7	(−6.9; 5.4)	0.000
Pack-years (mean) ^d	20.9	12.5	16.9	19.2	0.000	6.3	(2.9; 9.7)	16.7	23.8	28.7	0.002	11.4	(4.1; 18.7)	0.180
No alcohol consumption (%)	38.0	26.1	35.9	45.7	0.000	19.6	(14.2; 25.1)	30.2	35.3	40.4	0.007	10.3	(2.8; 17.7)	0.258
Heavy drinking (%)	4.7	4.5	4.5	4.6	0.894	0.2	(−2.2; 2.6)	6.4	5.3	4.2	0.187	−2.2	(−5.5; 1.1)	0.115
Physical inactivity (%)	34.8	20.3	31.7	43.0	0.000	22.7	(17.4; 28.0)	18.8	29.4	40	0.000	21.2	(14.0; 28.5)	0.776
Body mass index														
(mean, kg/m ²)	25.0	23.0	24.1	25.2	0.000	2.2	(1.7; 2.8)	23.6	25.1	26.6	0.000	3.0	(2.3; 3.8)	0.086
Overweight ^e (%)	41.8	21.3	32.7	44.1	0.000	22.9	(17.5; 28.2)	28.5	42.4	56.4	0.000	27.9	(20.3; 35.5)	0.305
Obesity ^e (%)	13.8	5.0	10.1	15.2	0.000	10.3	(6.6; 13.9)	5.4	12.8	20.3	0.000	14.8	(9.1; 20.6)	0.187
Waist circumference														
(mean, cm)	83.0	77.5	79.9	82.3	0.000	4.8	(3.5; 6.1)	81.1	84.6	88	0.000	7.0	(5.1; 8.9)	0.061
Abdominal obesity ^e (%)	31.7	13.4	21.3	29.1	0.000	15.7	(10.9; 20.5)	26.4	37.2	47.9	0.000	21.6	(14; 29.1)	0.139
Blood pressure (mmHg)														
Systolic (mean)	124.0	115.2	116.9	118.7	0.000	3.5	(1.9; 5.1)	128.8	131.2	133.6	0.001	4.8	(2.1; 7.6)	0.241
Diastolic (mean)	77.4	74.6	75.8	77.0	0.000	2.4	(1.2; 3.6)	78.0	78.7	79.4	0.101	1.4	(−0.3; 3.0)	0.087
Hypertension ^e (%)	25.0	8.0	12.3	16.6	0.000	8.6	(4.8; 12.5)	31.8	37.5	43.2	0.003	11.4	(4.0; 18.8)	0.369
HDL-cholesterol^f														
(mean, mmol/l)	1.81	1.89	1.81	1.73	0.000	−0.16	(−0.21; −0.11)	1.93	1.86	1.79	0.000	−0.14	(−0.21; −0.07)	0.970
Low HDL-cholesterol ^e (%)	4.3	0.6	3.7	6.8	0.000	6.2	(3.8; 8.6)	2.7	3.8	4.8	0.199	2.1	(−1.1; 5.2)	0.027
LDL-cholesterol														
(mean, mmol/l)	3.26	2.89	2.99	3.10	0.000	0.21	(0.12; 0.31)	3.50	3.55	3.59	0.172	0.09	(−0.04; 0.24)	0.036
High LDL-cholesterol ^e (%)	42.2	24.1	28.8	33.6	0.000	9.5	(4.4; 14.7)	52.3	56.2	60.1	0.044	7.8	(0.2; 15.4)	0.330
Triglycerides														
(mean, mmol/l)	1.12	1.37	1.43	1.50	0.005	0.13	(0.04; 0.22)	1.38	1.48	1.57	0.001	0.18	(0.08; 0.29)	0.647
High triglycerides ^e (%)	15.0	5.6	9.7	13.8	0.000	8.2	(4.6; 11.7)	14.9	18.7	22.4	0.018	7.5	(1.3; 13.8)	0.859
Fasting glucose														
(mean, mmol/l)	5.32	5.11	5.16	5.21	0.018	0.09	(0.02; 0.17)	5.33	5.45	5.58	0.008	0.25	(0.07; 0.43)	0.165
Diabetes ^e (%)	3.2	0.4	1.1	1.7	0.049	1.3	(0.0; 2.6)	4.1	5.4	6.7	0.168	2.6	(−1.1; 6.3)	0.414

BP: blood pressure; CI: confidence interval; HDL: high-density lipoprotein; LDL: low-density lipoprotein. Prevalence and mean values are adjusted for age and place of birth (Switzerland or outside Switzerland).

^a*p* for linear trend across socioeconomic categories.

^bDifference in prevalence/mean between the highest and the lowest educational category.

^c*p* for interaction between educational level and age group.

^dAnalyses restricted to current smokers (N = 503 in the 35–54 years group and N = 288 in the 55–75 years group). 31 smokers with missing information on pack-years were not included.

^eOverweight: BMI ≥25 kg/m² and <30 kg/m²; obesity: BMI ≥30 kg/m²; abdominal obesity: waist circumference ≥102/88cm in men/women; hypertension: BP ≥140/90 mmHg or taking BP treatment; low HDL-cholesterol: <1.0/1.2 mmol/l in men/women; high LDL-cholesterol: ≥3.4 mmol/l; high triglycerides: ≥1.7 mmol/l; diabetes: fasting glucose ≥7.0 mmol/l or taking diabetes treatment.

^fAnalyses for HDL-cholesterol are additionally adjusted for oral contraceptive use.

doi:10.1371/journal.pone.0049443.t003

differences in the social patterning of CVRF, with greater social inequalities in younger cohorts, or reflect a decrease in social inequalities in CVRF with ageing. As reported earlier, several studies conducted in Southern Europe (including one study in the French-speaking part of Switzerland) -mostly based on data from the early 1990s- found a small or null socioeconomic gradient in CVRF [2,11,13,14,19,21,22]. Our study is one of the first conducted in a Southern European country to find large socioeconomic differences in CVRF, which may hint at either a new situation in Southern Europe or at a difference between Switzerland and other Southern European countries. If the first

hypothesis is true, the fact that inequalities in CVRF tended to be stronger among younger than older participants may translate into an increase in social inequalities in adverse CVD outcomes over the next decades. Alternatively, smaller inequalities in CVRF in the older age group could be explained by the fact that relative inequalities in CVRF might decline with age as a result of increasing prevalence of adverse CVRF across socioeconomic groups or because of selection effects. However, both explanations remain speculative as the cross sectional nature of the study precludes distinguishing between age and cohort effects.

Table 4. Relative educational inequalities in selected cardiovascular risk factors by gender and age group.

	MEN			WOMEN			GENDER DIFFERENCES			
	Overall (N = 2963)	35-54 y (N = 1727)	55-75 y (N = 1233)	Overall (N = 3343)	35-54 y (N = 1842)	55-75 y (N = 1501)	Overall	35-54y	55-75y	
	RII (95%CI) ^a	RII (95%CI) ^a	RII (95%CI) ^a	RII (95%CI) ^{ab}	RII (95%CI) ^{ab}	RII (95%CI) ^{ab}	p ^c	p ^d	p ^d	
Behavioral risk factors										
Smoking	1.85 (1.48;2.31)	2.26 (1.71;2.98)	1.22 (0.83;1.79)	1.55 (1.22;1.98)	1.93 (1.44;2.58)	0.93 (0.6;1.45)	0.001	0.365	0.774	0.287
No alcohol consumption	1.44 (1.07;1.94)	1.63 (1.13;2.34)	1.12 (0.66;1.88)	2.03 (1.69;2.45)	2.29 (1.82;2.87)	1.52 (1.12;2.06)	0.171	0.027	0.120	0.160
Heavy drinking	1.46 (0.99;2.14)	2.75 (1.55;4.88)	0.88 (0.53;1.47)	0.75 (0.40;1.40)	1.09 (0.48;2.46)	0.49 (0.19;1.26)	0.087	0.066	0.086	0.333
Physical inactivity	2.98 (2.43;3.65)	3.27 (2.52;4.25)	2.53 (1.83;3.51)	2.84 (2.31;3.50)	2.93 (2.26;3.79)	2.64 (1.87;3.73)	0.639	0.707	0.539	0.902
Obesity ^e	2.95 (2.09;4.16)	3.30 (1.95;5.58)	2.82 (1.80;4.43)	4.77 (3.15;7.22)	4.74 (2.72;8.26)	4.76 (2.56;8.86)	0.991	0.177	0.347	0.215
Abdominal obesity ^e	1.48 (1.17;1.87)	2.42 (1.60;3.66)	1.18 (0.89;1.56)	2.60 (2.06;3.27)	3.11 (2.20;4.41)	2.24 (1.65;3.04)	0.153	0.012	0.380	0.005
Physiological risk factors										
Hypertension ^d	1.50 (1.26;1.79)	1.64 (1.19;2.26)	1.55 (1.26;1.91)	1.85 (1.43;2.39)	2.84 (1.75;4.61)	1.57 (1.16;2.11)	0.013	0.017	0.048	0.706
Low HDL-cholesterol ^e	2.06 (1.04;4.09)	2.45 (1.08;5.58)	1.32 (0.39;4.48)	5.62 (2.64;11.94)	12.67 (4.48;35.85)	1.79 (0.59;5.42)	0.008	0.048	0.019	0.763
High LDL-cholesterol ^e	1.11 (0.97;1.27)	1.16 (0.97;1.38)	1.06 (0.86;1.31)	1.37 (1.16;1.62)	1.75 (1.33;2.30)	1.21 (0.99;1.48)	0.004	<0.001	0.021	0.183
High triglycerides ^e	1.43 (1.17;1.76)	1.40 (1.07;1.83)	1.50 (1.10;2.05)	2.45 (1.69;3.54)	3.56 (2.03;6.23)	1.88 (1.15;3.08)	0.041	<0.001	0.001	0.369
Diabetes ^e	2.20 (1.35;3.58)	4.61 (1.62;13.1)	1.89 (1.09;3.27)	2.35 (1.00;5.53)	5.12 (0.84;31.33)	1.87 (0.71;4.88)	0.308	0.878	0.948	0.988

CI: confidence interval; HDL: high-density lipoprotein; LDL: low-density lipoprotein; RII: relative index of inequality; y:years.

^aAdjusted for age and place of birth (Switzerland or outside Switzerland).

^bAnalyses for HDL-cholesterol are additionally adjusted for oral contraceptive use.

^cp for interaction between RII and age group.

^dp for interaction between RII and sex.

^eObesity: BMI ≥ 30 kg/m²; abdominal obesity: waist circumference $\geq 102/88$ cm in men/women; hypertension: BP $\geq 140/90$ mmHg or taking BP treatment; low HDL-cholesterol: $<1.0/1.2$ mmol/l in men/women; high LDL-cholesterol: ≥ 3.4 mmol/l; high triglycerides: ≥ 1.7 mmol/l; diabetes: fasting glucose ≥ 7.0 mmol/l or taking diabetes treatment.

doi:10.1371/journal.pone.0049443.t004

Evidence for a prominent role of behavioral and biological risk factors such as those examined in this paper in explaining social inequalities in cardiovascular disease incidence and mortality is accumulating [45,46,47]. The determinants of the uneven distribution of CVRF across socioeconomic groups remain poorly understood, but likely include socioeconomic differences in several domains such as social norms, physical living and working environments, health education, health consciousness, attitude and motivation, psycho-social characteristics, and access to and utilization of health care [48,49,50]. We could not examine the role of this broader context in relation to our findings, as these factors were not assessed in our study. Further studies will be needed to elucidate the relative importance of specific factors in the social patterning of CVRF if effective policies to reduce social inequalities in health are to be implemented.

Strengths and Limitations

The main strength of this study was the availability of a large number of CVRF in a population-based survey covering a wide age range. This study also has potential limitations. The first relates to the inability of the cross-sectional design to distinguish between cohort and age effects. While we speculate that cohort-related changes might be taking place in the social patterning of CVRF, consistently with data from cohort studies or from repeated cross sectional surveys in other populations, we cannot exclude that the observed cohort differences in our study are accountable by age-related changes in behaviors. Second, measurement of socioeconomic position is challenging. Education is a valid indicator of SES as it allows for comparison of men and women and is applicable to the non-working population. However, it can have a different meaning for different birth cohorts, due to secular trends in educational attainment across generations [51]. Our sensitivity analysis using occupational position showed that our findings hold across indicators of socioeconomic status. Finally, health behaviors (smoking, alcohol consumption and physical activity) were self-reported and it has

been shown that questionnaire-based measures are not entirely reliable [52,53].

Conclusions

This study shows that large socioeconomic differences exist in the prevalence of several CVRF in a country enjoying one of the highest life expectancies at birth and one of the highest gross domestic products per capita in the world [54]. Although the overall prevalence of several CVRF was higher in men than in women, social inequalities tended to be greater in women. Socioeconomic gradients in CVRF were larger in the younger than in the older generations, suggesting that social inequalities in CVD might widen over the next decades. Further research is needed in order to elucidate the mechanisms underlying social inequalities in CVRF.

Supporting Information

Table S1 Relative educational inequalities in cardiovascular risk factors by gender and place of birth. (DOCX)

Acknowledgments

The authors also express their gratitude to the participants in the Lausanne CoLaus study and to the investigators who have contributed to the recruitment, in particular Yolande Barreau, Anne-Lise Bastian, Binasa Ramic, Martine Moranville, Martine Baumer, Marcy Sagette, Jeanne Ecoffey and Sylvie Mermoud for data collection.

Author Contributions

Conceived and designed the experiments: SS BS PB. Performed the experiments: SS. Analyzed the data: SS. Contributed reagents/materials/analysis tools: PV GW. Wrote the paper: SS. Critically revised the manuscript for important intellectual content: PMV PV GW FP. Collected the data: PV GW PMV FP. Received funding for the study: PW GW.

References

- Kaplan GA, Keil JE (1993) Socioeconomic factors and cardiovascular disease: a review of the literature. *Circulation* 88: 1973–1998.
- Cavelaars AE, Kunst A, Mackenbach JP (1997) Socio-economic differences in risk factors for morbidity and mortality in the European Community: An international comparison. *Journal of health psychology* 2: 353–372.
- Kanjilal S, Gregg EW, Cheng YJ, Zhang P, Nelson DE, et al. (2006) Socioeconomic status and trends in disparities in 4 major risk factors for cardiovascular disease among US adults, 1971–2002. *Archives of Internal Medicine* 166: 2348–2355.
- Ramsay SE, Whincup PH, Hardoon SL, Lennon LT, Morris RW, et al. (2011) Social class differences in secular trends in established coronary risk factors over 20 years: a cohort study of British men from 1978–80 to 1998–2000. *PLoS ONE* 6: e19742.
- Pell JP, Pell AC, Norrie J, Ford I, Cobbe SM (2000) Effect of socioeconomic deprivation on waiting time for cardiac surgery: retrospective cohort study. *BMJ* 320: 15–18.
- Mackenbach JP, Stirbu I, Roskam AJ, Schaap MM, Menvielle G, et al. (2008) Socioeconomic inequalities in health in 22 European countries. *New England Journal of Medicine* 358: 2468–2481.
- Mackenbach JP, Cavelaars AE, Kunst AE, Groenhouf F (2000) Socioeconomic inequalities in cardiovascular disease mortality; an international study. *European Heart Journal* 21: 1141–1151.
- Osler M (1993) Social class and health behaviour in Danish adults: a longitudinal study. *Public Health* 107: 251–260.
- Bartley M, Fitzpatrick R, Firth D, Marmot M (2000) Social distribution of cardiovascular disease risk factors: change among men in England 1984–1993. *Journal of Epidemiology and Community Health* 54: 806–814.
- Huisman M, Kunst A, Mackenbach JP (2005) Educational inequalities in smoking among men and women aged 16 years and older in 11 European countries. *Tobacco Control* 14: 106–113.
- Roos G, Johansson L, Kasmel A, Klumbiene J, Prattala R (2001) Disparities in vegetable and fruit consumption: European cases from the north to the south. *Public Health Nutrition* 4: 35–43.
- Wagner A, Simon C, Evans A, Ducimetiere P, Bongard V, et al. (2003) Physical activity patterns in 50–59 year men in France and Northern Ireland. Associations with socio-economic status and health behaviour. *European Journal of Epidemiology* 18: 321–329.
- de Vogli R, Gnesotto R, Goldstein M, Andersen R, Cornia GA (2005) The lack of social gradient of health behaviors and psychosocial factors in Northern Italy. *Sozial- und Präventivmedizin* 50: 197–205.
- Schroder H, Rohlfis I, Schmelz EM, Marrugat J (2004) Relationship of socioeconomic status with cardiovascular risk factors and lifestyle in a Mediterranean population. *European Journal of Nutrition* 43: 77–85.
- Stringhini S, Dugravot A, Shipley M, Goldberg M, Zins M, et al. (2011) Health Behaviours, Socioeconomic Status, and Mortality: Further Analyses of the British Whitehall II and the French GAZEL Prospective Cohorts. *PLoS Medicine* 8.
- Marmot MG, Adelstein AM, Robinson N, Rose GA (1978) Changing social-class distribution of heart disease. *British Medical Journal* 2: 1109–1112.
- Wilkinson RG (1994) The epidemiological transition: from material scarcity to social disadvantage? *Daedalus* 123: 61–77.
- Lopez AD, Collishaw NE, Piha T (1994) A descriptive model of the cigarette epidemic in developed countries. *Tobacco Control* 3: 242–247.
- Curtin F, Morabia A, Bernstein M (1997) Smoking behavior in a Swiss urban population: the role of gender and education. *Preventive Medicine* 26: 658–663.
- Borrell C, Dominguez-Berjon F, Pasarin MI, Ferrando J, Rohlfis I, et al. (2000) Social inequalities in health related behaviours in Barcelona. *Journal of Epidemiology and Community Health* 54: 24–30.
- Graham H (1996) Smoking prevalence among women in the European community 1950–1990. *Social Science and Medicine* 43: 243–254.
- Galobardes B, Costanza MC, Bernstein MS, Delhumeau C, Morabia A (2003) Trends in risk factors for lifestyle-related diseases by socioeconomic position in Geneva, Switzerland, 1993–2000: health inequalities persist. *American Journal of Public Health* 93: 1302–1309.

23. Marques-Vidal P, Cerveira J, Paccaud F, Cornuz J (2011) Smoking trends in Switzerland, 1992–2007: a time for optimism? *Journal of Epidemiology and Community Health* 65: 281–286.
24. Marques-Vidal P, Bovet P, Paccaud F, Chiolero A (2010) Changes of overweight and obesity in the adult Swiss population according to educational level, from 1992 to 2007. *BMC Public Health* 10: 87.
25. Muller-Nordhorn J, Binting S, Roll S, Willich SN (2008) An update on regional variation in cardiovascular mortality within Europe. *European Heart Journal* 29: 1316–1326.
26. Firmann M, Mayor V, Vidal PM, Bochud M, Pecoud A, et al. (2008) The CoLaus study: a population-based study to investigate the epidemiology and genetic determinants of cardiovascular risk factors and metabolic syndrome. *BMC Cardiovascular Disorders* 8: 6.
27. UNESCO IIS (2011) International Standard Classification of Education. UNESCO.
28. World Health Organisation (2007) The challenge of obesity in the WHO European Region and the strategies for response: WHO, 73 p.
29. Mackenbach JP, Kunst AE (1997) Measuring the magnitude of socio-economic inequalities in health: an overview of available measures illustrated with two examples from Europe. *Social Science and Medicine* 44: 757–771.
30. Khang YH, Yun SC, Lynch JW (2008) Monitoring trends in socioeconomic health inequalities: it matters how you measure. *BMC Public Health* 8: 66.
31. Sergeant JC, Firth D (2006) Relative index of inequality: definition, estimation, and inference. *Biostatistics (Oxford, England)* 7: 213–224.
32. Faeh D, Braun J, Bopp M (2011) Prevalence of obesity in Switzerland 1992–2007: the impact of education, income and occupational class. *Obesity Reviews* 12: 151–166.
33. Zimmermann-Sloutskis D, Wanner M, Zimmermann E, Martin BW (2010) Physical activity levels and determinants of change in young adults: a longitudinal panel study. *The international journal of behavioral nutrition and physical activity* 7: 2.
34. Danon-Hersch N, Marques-Vidal P, Bovet P, Chiolero A, Paccaud F, et al. (2009) Prevalence, awareness, treatment and control of high blood pressure in a Swiss city general population: the CoLaus study. *European Journal of Cardiovascular Prevention and Rehabilitation* 16: 66–72.
35. Estoppey D, Paccaud F, Vollenweider P, Marques-Vidal P (2011) Trends in self-reported prevalence and management of hypertension, hypercholesterolemia and diabetes in Swiss adults, 1997–2007. *BMC Public Health* 11: 114.
36. Marques-Vidal P, Paccaud F (2012) Regional differences in self-reported screening, prevalence and management of cardiovascular risk factors in Switzerland. *BMC Public Health* 12: 246.
37. Morabia A, Costanza MC (2005) The obesity epidemic as harbinger of a metabolic disorder epidemic: trends in overweight, hypercholesterolemia, and diabetes treatment in Geneva, Switzerland, 1993–2003. *American Journal of Public Health* 95: 632–635.
38. Nedelchev K, Arnold M, Baumgartner R, Devuyst G, Erne P, et al. (2005) Vascular risk factors in the Swiss population. *Journal of Neurology* 252: 1210–1216.
39. Faeh D, Bopp M (2010) Educational inequalities in mortality and associated risk factors: German–versus French-speaking Switzerland. *BMC Public Health* 10: 567.
40. Galobardes B, Costanza MC, Bernstein MS, Delhumeau CH, Morabia A (2003) Trends in risk factors for the major "lifestyle-related diseases" in Geneva, Switzerland, 1993–2000. *Annals of epidemiology* 13: 537–540.
41. Zins M, Gueguen A, Kivimaki M, Singh-Manoux A, Leclerc A, et al. (2011) Effect of retirement on alcohol consumption: longitudinal evidence from the French Gazel cohort study. *PLoS ONE* 6: e26531.
42. Roskam AJ, Kunst AE, Van Oyen H, Demarest S, Klumbiene J, et al. (2010) Comparative appraisal of educational inequalities in overweight and obesity among adults in 19 European countries. *International Journal of Epidemiology* 39: 392–404.
43. McLaren L (2007) Socioeconomic status and obesity. *Epidemiologic Reviews* 29: 29–48.
44. Espelt A, Borrell C, Roskam AJ, Rodriguez-Sanz M, Stirbu I, et al. (2008) Socioeconomic inequalities in diabetes mellitus across Europe at the beginning of the 21st century. *Diabetologia* 51: 1971–1979.
45. Stringhini S, Sabia S, Shipley M, Brunner E, Nabi H, et al. (2010) Association of socioeconomic position with health behaviors and mortality The Whitehall II study. *Journal of the American Medical Association*.
46. Kivimaki M, Shipley MJ, Ferrie JE, Singh-Manoux A, Batty GD, et al. (2008) Best-practice interventions to reduce socioeconomic inequalities of coronary heart disease mortality in UK: a prospective occupational cohort study. *Lancet* 372: 1648–1654.
47. Woodside JV, Yarnell JW, Patterson CC, Arveiler D, Amouyel P, et al. (2012) Do lifestyle behaviours explain socioeconomic differences in all-cause mortality, and fatal and non-fatal cardiovascular events? Evidence from middle aged men in France and Northern Ireland in the PRIME Study. *Preventive Medicine* 54: 247–253.
48. Wardle J, Steptoe A (2003) Socioeconomic differences in attitudes and beliefs about healthy lifestyles. *Journal of Epidemiology and Community Health* 57: 440–443.
49. van Lenthe EJ, de Bourdeaudhuij I, Klepp KI, Lien N, Moore L, et al. (2009) Preventing socioeconomic inequalities in health behaviour in adolescents in Europe: background, design and methods of project TEENAGE. *BMC Public Health* 9: 125.
50. Chen WQ, Wong TW, Yu IT (2008) Association of occupational stress and social support with health-related behaviors among chinese offshore oil workers. *Journal of Occupational Health* 50: 262–269.
51. Galobardes B, Shaw M, Lawlor DA, Lynch JW, Davey Smith G (2006) Indicators of socioeconomic position (part 1). *Journal of Epidemiology and Community Health* 60: 7–12.
52. Prince SA, Adamo KB, Hamel ME, Hardt J, Gorber SC, et al. (2008) A comparison of direct versus self-report measures for assessing physical activity in adults: a systematic review. *The international journal of behavioral nutrition and physical activity* 5: 56.
53. Hatzidandreu EJ, Pierce JP, Fiore MC, Grise V, Novotny TE, et al. (1989) The reliability of self-reported cigarette consumption in the United States. *American Journal of Public Health* 79: 1020–1023.
54. World Bank (2012) World Development Indicators.