Expression of the RNA Helicase DDX3 and the Hypoxia Response in Breast Cancer

Guus M. Bol^{1,3}, Venu Raman^{1,3,4}, Petra van der Groep^{1,2}, Jeroen F. Vermeulen¹, Arvind H. Patel⁵, Elsken van der Wall², Paul J. van Diest^{1,4*}

1 Departments of Pathology, University Medical Center Utrecht Cancer Center, Utrecht, The Netherlands, **2** Division of Internal Medicine and Dermatology, University Medical Center Utrecht Cancer Center, Utrecht, The Netherlands, **3** Department of Radiology and Radiological Science, Johns Hopkins University School of Medicine, Baltimore, Maryland, United States of America, **4** Department of Oncology, Johns Hopkins University School of Medicine, Baltimore, Maryland, United States of America, **5** MRC, University of Glasgow Centre for Virus Research, Glasgow, United Kingdom

Abstract

Aims: DDX3 is an RNA helicase that has antiapoptotic properties, and promotes proliferation and transformation. In addition, DDX3 was shown to be a direct downstream target of HIF-1 α (the master regulatory of the hypoxia response) in breast cancer cell lines. However, the relation between DDX3 and hypoxia has not been addressed in human tumors. In this paper, we studied the relation between DDX3 and the hypoxic responsive proteins in human breast cancer.

Methods and Results: DDX3 expression was investigated by immunohistochemistry in breast cancer in comparison with hypoxia related proteins HIF-1 α , GLUT1, CAIX, EGFR, HER2, Akt1, FOXO4, p53, ER α , COMMD1, FER kinase, PIN1, E-cadherin, p21, p27, Transferrin receptor, FOXO3A, c-Met and Notch1. DDX3 was overexpressed in 127 of 366 breast cancer patients, and was correlated with overexpression of HIF-1 α and its downstream genes CAIX and GLUT1. Moreover, DDX3 expression correlated with hypoxia-related proteins EGFR, HER2, FOXO4, ER α and c-Met in a HIF-1 α dependent fashion, and with COMMD1, FER kinase, Akt1, E-cadherin, TfR and FOXO3A independent of HIF-1 α .

Conclusions: In invasive breast cancer, expression of DDX3 was correlated with overexpression of HIF-1 α and many other hypoxia related proteins, pointing to a distinct role for DDX3 under hypoxic conditions and supporting the oncogenic role of DDX3 which could have clinical implication for current development of DDX3 inhibitors.

Citation: Bol GM, Raman V, van der Groep P, Vermeulen JF, Patel AH, et al. (2013) Expression of the RNA Helicase DDX3 and the Hypoxia Response in Breast Cancer. PLoS ONE 8(5): e63548. doi:10.1371/journal.pone.0063548

Editor: Sharon A. Glynn, National University of Ireland Galway, Ireland

Received: November 22, 2012; **Accepted:** April 3, 2013; **Published:** May 16, 2013

Copyright: © 2013 Bol et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by the Dr. Saal van Swanenberg foundation [GMB], the Dutch Cancer Society (UU 2010-4856)[GMB] and FAMRI [VR]. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: Paul van Diest is a PLOS ONE Editorial Board member. This does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials.

* E-mail: P.J.vandiest@umcutrecht.nl

Introduction

In the Western world, one in eight women will develop breast cancer during their life and breast cancer is causing about 458,000 deaths worldwide per year [1,2]. Aggressive forms of breast cancer are frequently refractory to treatment [3], even to established targeted therapy, and thus have a high risk of relapse and formation of distant metastases [4]. Identification of molecular pathways involved in aggressive forms of breast cancer is therefore important to design novel targeted therapeutic agents to counteract tumor progression and metastasis.

DDX3, also known as DDX3X because of its location on the X chromosome, is a member of the DEAD-box RNA helicase family which is involved in transcription, RNA splicing, nuclear export of mRNA and translation initiation [5,6]. Initially, DDX3 was studied because of its manipulation by viruses like hepatitis C (HCV) and human immunodeficiency virus (HIV) [7,8]. Recently DDX3 has been associated with cancer [9]. Conflicting evidence exists with regard to its tumor enhancing or repressing properties. Nevertheless, DDX3 was proven to have antiapoptotic properties [10,11], promotes proliferation and cellular transformation [9,12–

14]. Recently, novel compounds were developed which could potentially inhibit DDX3 activity [15–20].

A recent *in vitro* study [21] showed that DDX3 is a direct downstream target of HIF-1 α , the predominant factor in the mammalian hypoxia response [22]. Hypoxia is an important event in breast carcinogenesis [23–26], causing a more aggressive phenotype with increased invasiveness and proliferation, formation of metastases, resistance to therapy [27] and poorer survival [28,29].

However, no data are yet available on the relation between DDX3 and hypoxia in human breast cancer, or any other human tumors specimens. Therefore, we set out to correlate expression of DDX3 and HIF-1 α in a large set of human invasive breast cancers. Furthermore, we correlated DDX3 expression to expression of various other proteins upstream of HIF-1 α like EGFR [30], HER2 [31], Akt1 [32–34], p53 [35–39], COMMD1 [40,41], FER kinase [42], PIN1 [43] and FOXO4 [44]. Also we assessed proteins downstream of HIF-1 α such as ER α [45,46] Transferrin receptor (TfR) [47], FOXO3A [48] and Notch1 [49,50]. Finally, we included proteins that have been associated

with HIF-1 α without clear functional relationship like E-cadherin [51], p21 [52], c-Met [53,54] and p27 [55].

Materials and Methods

Patients

Representative paraffin embedded tissue blocks of 422 breast cancer patients collected between 2004 and 2007 were taken from the archive of the Department of Pathology of the University Medical Centre in Utrecht and routinely processed to four tissue microarrays (TMA) as described before [56,57].

Clinicopathological data including tumor stage, histological data (type, grade, mitotic index (MAI), estrogen receptor alpha (ER α) and human epidermal growth factor receptor 2 (HER2)) status was collected from patient files (Table 1). Protein expression data by immunohistochemistry of HIF-1 α , FOXO3A, FOXO4, PIN1, Akt1, COMMD1, p53, p21, p27, EGFR, E-cadherin, GLUT1 and CAIX was derived from previous studies [34,40,58–62].

Use of anonymous or coded left over material for scientific purposes is part of the standard treatment contract with patients in the UMCU [63].

Immunohistochemistry

Sections of 4 μ m were cut, mounted on SuperFrost slides (Menzel&Glaeser, Brunswick, Germany), deparaffinized and rehydrated. Endogenous peroxidase was then blocked for 15 min with a buffer solution containing 0.3% hydrogenperoxide. Antigens were retrieved by boiling for 20 min in 10 mM citrate buffer (pH 6.0) (for DDX3, c-Met, TrR, FER kinase and Notch1), cooled and washed in PBS. Nonspecific binding sites were blocked with a 2% normal goat serum, 1% BSA in PBS (pH 7.4) (Notch1). TMAs were subsequently incubated in a humidified chamber for 1 hour with polyclonal rabbit anti-DDX3 R648 [64] diluted 1:1000, TrR 1:300 (13–6800, Invitrogen, Breda, The Netherlands) and FER kinase 1:300 (clone 5D2, Cell Signaling Technologies, USA). Primary antibodies against c-Met 1:100 (18-2257, Zymed, Invitrogen) and Notch1 1:100 (Cell Signaling Technologies, USA) were incubated overnight at 4°C. Subsequently, sections were washed in PBS and incubated for 30 min with secondary antibodies (Brightvision, Immunologic, Duiven, The Netherlands) washed with PBS and developed with diaminobenzidine. Slides were counterstained with hematoxylin, dehydrated and cover-slipped. Appropriate positive and negative controls were used throughout.

Table 1. Patient characteristics.

	N (422)		missing
Mean age (range)	61.0	(28–88)	0
Tumor size			
$\leq$ 20 mm	212	50%	3
$\leq$ 50 mm	181	43%	
>50 mm	26	6%	
Lymph node status			
Positive*	193	48%	18
Negative**	211	52%	
Histological type			
ductal	343	82%	1
lobular	42	10%	
other	36	9%	
Grade			
I	80	20%	26
II	145	37%	
III	171	43%	
Mitotic index (range)	17.2	(0–196)	0
Estrogen receptor [#]			
Positive	335	79%	0
Negative	87	21%	
Progesterone receptor [#]			
Positive	247	59%	1
Negative	174	41%	
HER2 receptor			
Positive	44	10%	0
Negative	378	90%	

*Positive = $\geq$ N1mi.

**Negative = N0 or N0(i+) (according to TNM 7th edition, 2010).

[#]10% cut-off.

doi:10.1371/journal.pone.0063548.t001

Figure 1. Examples of DDX3 and HIF-1 α staining. Breast cancer photomicrographs are taken at 20X. A. low HIF-1 α expression (0%); B. low DDX3 expression (1), same patient as in A; C. high HIF-1 α expression (90%); D. high DDX3 expression (3), same patient as in C. doi:10.1371/journal.pone.0063548.g001

Scoring of Immunohistochemistry

Scoring was done by a single experienced pathologist (PJvD). Intensity of cytoplasmic DDX3, FER kinase and membranous E-cadherin, TIR and c-Met was scored semi-quantitatively from 0–3 and percentages of cells with nuclear DDX3 and Notch1 expression were estimated. Out of three cores from the same patient, the maximum cytoplasmic DDX3 score was used for further analysis.

DDX3 scores 1 and 2 were grouped as low DDX3 expression and evaluated against high DDX3 expression (scores 3). For E-cadherin, TIR, c-Met and FER kinase scores 0 and 1 were defined as low expression versus score 2 and 3 as high expression. For HIF-1 α , the 1% threshold was used as before [59].

Statistics

Expression levels of DDX3 and the other proteins were compared by chi-square test or t-test whenever applicable. Logistic regression or ANCOVA was used for multivariate analysis to determine dependence of these relations on HIF-1 α .

Since EGFR and HER2 are upstream regulators of HIF-1 α via PI-3K/AKT, we also assessed the relation of EGFR and HER2 with DDX3 independent of Akt1 and HIF-1 α . In lobular breast cancer there is very little or no expression of E-cadherin, so the lobular cancers were excluded in analysis with respect to E-cadherin.

Pearson correlation coefficient was determined for correlation analysis.

All statistical analyses were carried out with SPSS 17.0 for Windows. (SPSS Inc., Chicago, IL, USA), regarding two-sided p-values below 0.05 as significant.

Table 2. Expression of DDX3 in relation to oxygen sensing proteins.

		Cytoplasmic DDX3				Multivariate		
		<i>N</i> (%)	Low (%)	High (%)	OR	<i>p</i> value ^a	OR	<i>p</i> value ^b
		366	239	127				
HIF-1α	≤1%	214 (66)	155 (75)	59 (51)	2.83	<0.001	2.52	0.001
	>1%	108 (34)	52 (25)	56 (49)				
GLUT1	negative	123 (39)	94 (46)	29 (27)	2.36	0.001	1.94	0.021
	positive	190 (61)	110 (54)	80 (73)				
CAIX	negative	62 (19)	49 (24)	13 (11)	2.39	0.012	2.23	0.042
	positive	260 (81)	159 (76)	101 (89)				

^achi-square test.^blogistic regression.

doi:10.1371/journal.pone.0063548.t002

Results

DDX3 staining could be evaluated in 366 of the 422 breast cancer cases. The drop outs were caused by damaged or detached cores during cutting, mounting, or staining, or did not contain tumor. All breast cancer cases showed some expression of DDX3 of which 127 (35%) showed strong cytoplasmic DDX3 expression.(table 2).

HIF-1 α overexpression correlated with expression of CAIX, GLUT1, EGFR, HER2, Akt1, FER kinase, ER α , FOXO4, TIR, c-Met as expected (data not shown). Strong cytoplasmic DDX3 expression was associated with overexpression of the master

regulator of the hypoxia response HIF-1 α (OR = 2.83; $p < 0.001$) (figure 1) and its downstream proteins GLUT1 (OR = 2.36; $p = 0.001$) and CAIX (2.39; $p = 0.012$). In logistic regression, HIF-1 α (OR = 2.52; $p = 0.001$), GLUT1 (OR = 1.94; $p = 0.021$), CAIX (OR = 2.23; $p = 0.042$) predicted cytoplasmic DDX3 levels independently.(table 2).

The HIF-1 α transcription regulators HER2 (OR = 3.04; $p = 0.001$), EGFR (OR = 2.01; $p = 0.013$) and Akt1 (83% vs. 95%; $p = 0.001$) were correlated with DDX3 expression. Association of EGFR and HER2 with DDX3 was dependent on HIF-1 α .(table 3).

Table 3. Expression of DDX3 in relation to regulators of HIF-1 α .

		Cytoplasmic DDX3				Correction for HIF-1 α			
		<i>N</i> (%)	Low (%)	High (%)	OR	<i>p</i> value ^a	OR	<i>p</i> value ^b	
EGFR		366	239	127					
	negative	299 (83)	204 (86)	95 (76)	2.01	0.013	1.61	0.134	
HER2	positive	62 (17)	32 (14)	30 (24)					
	negative	325 (89)	222 (93)	103 (81)	3.04	0.001	1.88	0.092	
p53	positive	41 (11)	17 (7)	24 (19)					
	negative	80 (81)	45 (82)	35 (80)	1.16	0.802	1.42	0.571	
COMMD1	positive	19 (19)	10 (18)	9 (20)					
	low	23 (29)	17 (41)	6 (15)	3.90	0.013	5.45	0.006	
FER kinase	high	57 (71)	24 (59)	33 (85)					
	low	203 (57)	161 (70)	42 (34)	4.49	<0.001	4.10	<0.001	
PIN1	high	152 (43)	70 (30)	82 (66)					
	low	58 (71)	32 (78)	26 (63)	2.05	0.225	1.73	0.305	
	high	24 (29)	9 (22)	15 (37)					
		Cytoplasmic DDX3				Cytoplasmic DDX3			
		<i>N</i>	Low	High	<i>p</i> value ^c	<i>N</i>	Low	High	<i>p</i> value ^d
Akt1		80	83%	95%	0.001	75	86%	95%	0.026
FOXO4		75	30%	16%	0.035	61	21%	17%	0.600

^achi-square test.^blogistic regression.^cstudent's t-test.^dANCOVA.

doi:10.1371/journal.pone.0063548.t003

Proteins known to regulate HIF-1 α in a PI-3K/AKT independent fashion were associated with DDX3 as well; COMMD1 (OR = 3.90; $p = 0.013$), FER kinase (OR = 4.49; $p < 0.001$), FOXO4 (30% vs. 16%; $p = 0.035$), but not p53 (OR = 1.16; $p = 0.802$) and PIN1 (OR = 2.05; $p = 0.225$), (table 3). Logistic regression indicated a HIF-1 α independent relation between cytoplasmic DDX3 on the one hand and COMMD1 (OR = 5.45; $p = 0.006$) and FER kinase (OR = 4.10; $p < 0.001$) on the other, (table 3).

DDX3 was further associated with ER α (OR = 0.48; $p = 0.005$), E-cadherin (OR = 2.84; $p = 0.005$), TfR (OR = 2.77; $p < 0.001$), c-Met (OR = 1.72; $p = 0.042$) and FOXO3A (83% vs. 94%; $p = 0.021$). (table 4) After correction for HIF-1 α expression, E-cadherin (OR = 2.91; $p = 0.009$), TfR (OR = 2.01; $p = 0.007$) and FOXO3A (78% vs. 95%; $p = 0.007$) were still associated with DDX3, (table 4). Table 5 shows the Pearson correlation analysis results.

Discussion

The aim of this study was to investigate the relation between DDX3 and the hypoxic response in human breast cancer in the light of *in vitro* results pointing to regulation of DDX3 by HIF-1 α . We indeed show a positive correlation between HIF-1 α and DDX3 overexpression in a large series of human breast cancer cases, as well as an association between DDX3 overexpression and various other hypoxia related proteins.

However, we have established a correlation between DDX3 overexpression and nuclear HIF-1 α overexpression which supports the direct regulation of DDX3 by HIF-1 α found *in vitro* [21], but this is obviously no more than an association at this point no

Table 5. DDX3 correlations with the most important hypoxia related proteins.

	N	r ^a	p value
HIF-1 α	322	0.276	<0.001
GLUT1	313	0.186	0.001
CAIX	322	0.136	0.015
HER2	366	0.185	<0.001
ER α	366	-0.132	0.011

^aPearson correlation coefficient.

doi:10.1371/journal.pone.0063548.t005

proof for a causal relationship. Immunohistochemistry has some limitations like being inherently a more qualitative than quantitative method, and semiquantitative scoring and dichotomization with non-optimal reproducibility. To compensate for these issues we standardized the IHC procedure, used control tissue throughout, scored three samples per patient, studied a large cohort of breast cancer patients and results obtained from dichotomized parameters were confirmed by correlation analysis for the most important parameters with the DAKO score of DDX3 (table 5). Patient features in this study corresponded with known clinicopathological characteristics in breast cancer (table 1) [65].

Furthermore, DDX3 correlated with EGFR, HER2, FOXO4, ER α and c-Met in a HIF-1 α dependent way. Also, we found a positive correlation with COMMD1, FER kinase, Akt1, E-cadherin, TfR and FOXO3A independent of HIF-1 α . COMMD1 down regulates HIF-1 α by competition with HSP90 β [41], or

Table 4. Expression of DDX3 in relation to various other hypoxia induced proteins.

		<i>N</i> (%)	Low (%)	High (%)	OR	<i>p</i> value ^a	OR	<i>p</i> value ^b	
		366	239	127					
ERα	negative	81 (22)	42 (18)	39 (31)	0.48	0.005	0.67	0.166	
	positive	285 (78)	197 (82)	88 (69)					
E-cadherin*	low	48 (17)	38 (23)	10 (9)	2.84	0.005	2.91	0.009	
	high	227 (83)	130 (77)	97 (91)					
p21	low	46 (46)	28 (51)	18 (41)	1.50	0.418	1.26	0.625	
	high	53 (54)	27 (49)	26 (59)					
TfR	low	221 (63)	161 (72)	60 (48)	2.77	<0.001	2.01	0.007	
	high	128 (37)	63 (28)	65 (52)					
c-Met	negative	264 (77)	181 (81)	83 (71)	1.72	0.042	1.62	0.096	
	positive	77 (23)	43 (19)	34 (29)					
Cytoplasmic DDX3					Cytoplasmic DDX3				
		<i>N</i>	Low	High	<i>p</i> value ^c	<i>N</i>	Low	High	<i>p</i> value ^d
p27		99	42%	44%	0.683	77	40%	45%	0.507
FOXO3A		86	83%	94%	0.021	71	78%	95%	0.007
Notch1		305	63%	55%	0.064	282	61%	56%	0.299

^achi-square test.

^blogistic regression.

^cstudent's t-test.

^dANCOVA.

*in ductal breast cancer.

TfR = Transferrin receptor.

doi:10.1371/journal.pone.0063548.t004

down regulates the transcriptional activity of HIF-1 α [40]. However, we could not detect an association with COMMD1 and HIF-1 α expression or its downstream targets: E-cadherin, TIR, p21, p27 or c-Met. Nonetheless, COMMD1 correlates with DDX3 independent of HIF-1 α .

FER kinases help cells to withstand stress, including hypoxia, via up regulation of HIF-1 α [42]. We found a strong relation with FER kinase with both HIF-1 α and DDX3. After correction for the effect FER kinase has on HIF-1 α , a strong relation between FER kinase and DDX3 remained, implying a HIF-1 α dependent and independent relation.

DDX3 was shown to down regulate E-cadherin [12], but in the present study we show a positive correlation, for which we have no obvious explanation. TIR is also under transcriptional control of HIF-1 α [47]. TIR is overexpressed in many cancers, which could be attributed to the increased need for iron as a cofactor of the ribonucleotide reductase enzyme involved in DNA synthesis of rapidly dividing cells. Thus, the HIF-1 α independent relation between DDX3 and TIR corroborates previous reports on the oncogenic properties of DDX3. Nuclear expression of FOXO3A in breast cancer is associated with anti-apoptotic signaling via Akt1, an aggressive phenotype and poor survival [66]. In response to hypoxia, FOXO3A accumulates in a HIF-1 α dependent way to inhibit HIF-1 α induced apoptosis [48]. Although we did not find a relation between HIF-1 α and FOXO3A we did find a relation between FOXO3A and DDX3, independent of HIF-1 α and Akt1. Perhaps DDX3 and FOXO3A function in a concerted survival response after stress stimuli.

References

1. Ferlay J, Shin HR, Bray F, Forman D, Mathers C, et al. (2010) Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer* 127: 2893–2917.
2. Altekruse S, Kosary C, Krapcho M, Neyman N, Aminou R, et al. (2010) SEER Cancer Statistics Review, 1975–2007, National Cancer Institute. Bethesda, MD, based on November 2009 SEER data submission, posted to the SEER web site. http://seercancer.gov/csr/1975_2007/Accessed 2012 Nov.
3. Liedtke C, Mazouni C, Hess KR, Andre F, Tordai A, et al. (2008) Response to neoadjuvant therapy and long-term survival in patients with triple-negative breast cancer. *J Clin Oncol* 26: 1275–1281.
4. Dent R, Trudeau M, Pritchard KI, Hanna WM, Kahn HK, et al. (2007) Triple-negative breast cancer: clinical features and patterns of recurrence. *Clin Cancer Res* 13: 4429–4434.
5. Rocak S, Linder P (2004) DEAD-box proteins: the driving forces behind RNA metabolism. *Nat Rev Mol Cell Biol* 5: 232–241.
6. Lorsch JR (2002) RNA chaperones exist and DEAD box proteins get a life. *Cell* 109: 797–800.
7. Yedavalli VS, Neuvet C, Chi YH, Kleiman L, Jeang KT (2004) Requirement of DDX3 DEAD box RNA helicase for HIV-1 Rev-RRE export function. *Cell* 119: 381–392.
8. Owsianka AM, Patel AH (1999) Hepatitis C virus core protein interacts with a human DEAD box protein DDX3. *Virology* 257: 330–340.
9. Huang JS, Chao CC, Su TL, Yeh SH, Chen DS, et al. (2004) Diverse cellular transformation capability of overexpressed genes in human hepatocellular carcinoma. *Biochem Biophys Res Commun* 315: 950–958.
10. Sun M, Song L, Li Y, Zhou T, Jope RS (2008) Identification of an antiapoptotic protein complex at death receptors. *Cell Death Differ* 15: 1887–1900.
11. Li Y, Wang H, Wang Z, Makhija S, Buchsbaum D, et al. (2006) Inducible resistance of tumor cells to tumor necrosis factor-related apoptosis-inducing ligand receptor 2-mediated apoptosis by generation of a blockade at the death domain function. *Cancer Res* 66: 8520–8528.
12. Botlagunta M, Vesuna F, Mironchik Y, Raman A, Lisok A, et al. (2008) Oncogenic role of DDX3 in breast cancer biogenesis. *Oncogene* 27: 3912–3922.
13. Shih JW, Tsai TY, Chao CH, Wu Lee YH (2007) Candidate tumor suppressor DDX3 RNA helicase specifically represses cap-dependent translation by acting as an eIF4E inhibitory protein. *Oncogene* 27: 700–714.
14. Lee CS, Dias AP, Jedrychowski M, Patel AH, Hsu JL, et al. (2008) Human DDX3 functions in translation and interacts with the translation initiation factor eIF3. *Nucleic Acids Res* 36: 4708–4718.
15. Kondaskar A, Kondaskar S, Kumar R, Fishbein JC, Muvarak N, et al. (2010) Novel, Broad Spectrum Anti-Cancer Agents Containing the Tricyclic 5:7:5-Fused Diimidazodiazepine Ring System. *ACS Med Chem Lett* 2: 252–256.
16. Maga G, Falchi F, Garbelli A, Belfiore A, Witvrouw M, et al. (2008) Pharmacophore modeling and molecular docking led to the discovery of inhibitors of human immunodeficiency virus-1 replication targeting the human cellular aspartic acid-glutamic acid-alanine-aspartic acid box polypeptide 3. *J Med Chem* 51: 6635–6638.
17. Kumar R, Ujjinamata RK, Hosmane RS (2008) The first synthesis of a novel 5:7:5-fused diimidazodiazepine ring system and some of its chemical properties. *Org Lett* 10: 4681–4684.
18. Radi M, Falchi F, Garbelli A, Samuele A, Bernardo V, et al. (2012) Discovery of the first small molecule inhibitor of human DDX3 specifically designed to target the RNA binding site: towards the next generation HIV-1 inhibitors. *Bioorg Med Chem Lett* 22: 2094–2098.
19. Belon CA, High YD, Lin TI, Pauwels F, Frick DN (2010) Mechanism and specificity of a symmetrical benzimidazolephenylcarboxamide helicase inhibitor. *Biochemistry* 49: 1822–1832.
20. Yedavalli VS, Zhang N, Cai H, Zhang P, Starost MF, et al. (2008) Ring expanded nucleoside analogues inhibit RNA helicase and intracellular human immunodeficiency virus type 1 replication. *J Med Chem* 51: 5043–5051.
21. Botlagunta M, Krishnamachary B, Vesuna F, Winnard PT, Jr., Bol GM, et al. (2011) Expression of DDX3 is directly modulated by hypoxia inducible factor-1 alpha in breast epithelial cells. *PLoS ONE* 6: e17563.
22. Majumdar AJ, Wong WJ, Simon MC (2010) Hypoxia-inducible factors and the response to hypoxic stress. *Molecular cell* 40: 294–309.
23. Bos R, Zhong H, Hanrahan CF, Mommers EC, Semenza GL, et al. (2001) Levels of hypoxia-inducible factor-1 alpha during breast carcinogenesis. *J Natl Cancer Inst* 93: 309–314.
24. Bos R, Diest PJ, Groep Pvd, Greijer AE, Hermens MAJA, et al. (2003) Protein expression of B-cell lymphoma gene 6 (BCL-6) in invasive breast cancer is associated with cyclin D1 and hypoxia-inducible factor-1[alpha] (HIF-1[alpha]). *Oncogene* 22: 8948–8951.
25. Bos R, van Diest PJ, van der Groep P, Shvarts A, Greijer AE, et al. (2004) Expression of hypoxia-inducible factor-1-alpha and cell cycle proteins in invasive breast cancer are estrogen receptor related. *Breast Cancer Res* 6: R450–459.
26. Bos R, van Diest PJ, de Jong JS, van der Groep P, van der Valk P, et al. (2005) Hypoxia-inducible factor-1-alpha is associated with angiogenesis, and expression of bFGF, PDGF-BB, and EGFR in invasive breast cancer. *Histopathology* 46: 31–36.
27. Greijer AE, de Jong JS, Scheffer GL, Shvarts A, van Diest PJ, et al. (2005) Hypoxia-induced acidification causes mitoxantrone resistance not mediated by drug transporters in human breast cancer cells. *Cellular Oncology* 27: 43–49.
28. Semenza GL (2000) Hypoxia, clonal selection, and the role of HIF-1 in tumor progression. *Crit Rev Biochem Mol Biol* 35: 71–103.
29. Vaupel P (2004) The role of hypoxia-induced factors in tumor progression. *Oncologist* 9 Suppl 5: 10–17.
30. Peng XH, Karna P, Cao Z, Jiang BH, Zhou M, et al. (2006) Cross-talk between epidermal growth factor receptor and hypoxia-inducible factor-1-alpha signal

- pathways increases resistance to apoptosis by up-regulating survivin gene expression. *J Biol Chem* 281: 25903–25914.
31. Laughner E, Taghavi P, Chiles K, Mahon PC, Semenza GL (2001) HER2 (neu) signaling increases the rate of hypoxia-inducible factor 1alpha (HIF-1alpha) synthesis: novel mechanism for HIF-1-mediated vascular endothelial growth factor expression. *Mol Cell Biol* 21: 3995–4004.
  32. Mottet D, Dumont V, Deccache Y, Demazy C, Ninane N, et al. (2003) Regulation of hypoxia-inducible factor-1alpha protein level during hypoxic conditions by the phosphatidylinositol 3-kinase/Akt/glycogen synthase kinase 3beta pathway in HepG2 cells. *J Biol Chem* 278: 31277–31285.
  33. Brugarolas J, Kaelin WG (2004) Dysregulation of HIF and VEGF is a unifying feature of the familial hamartoma syndromes. *Cancer Cell* 6: 7–10.
  34. Gort EH, Groot AJ, van de Ven TLPD, van der Groep P, Verlaan I, et al. (2006) Hypoxia-inducible factor-1 alpha expression requires PI 3-kinase activity and correlates with Akt1 phosphorylation in invasive breast carcinomas. *Oncogene* 25: 6123–6127.
  35. Blagosklonny MV, An WG, Romanova LY, Trepel J, Fojo T, et al. (1998) p53 inhibits hypoxia-inducible factor-stimulated transcription. *J Biol Chem* 273: 11995–11998.
  36. Yamakuchi M, Lotterman CD, Bao C, Hruban RH, Karim B, et al. (2010) P53-induced microRNA-107 inhibits HIF-1 and tumor angiogenesis. *Proc Natl Acad Sci U S A* 107: 6334–6339.
  37. Ravi R, Mookerjee B, Bhujwala ZM, Sutter CH, Artemov D, et al. (2000) Regulation of tumor angiogenesis by p53-induced degradation of hypoxia-inducible factor 1alpha. *Genes Dev* 14: 34–44.
  38. Schmid T, Zhou J, Kohl R, Brune B (2004) p300 relieves p53-evoked transcriptional repression of hypoxia-inducible factor-1 (HIF-1). *Biochem J* 380: 289–295.
  39. Sano M, Minamino T, Toko H, Miyauchi H, Orimo M, et al. (2007) p53-induced inhibition of Hif-1 causes cardiac dysfunction during pressure overload. *Nature* 446: 444–448.
  40. van de Sluis B, Mao X, Zhai Y, Groot AJ, Vermeulen JF, et al. (2010) COMMD1 disrupts HIF-1alpha/beta dimerization and inhibits human tumor cell invasion. *J Clin Invest* 120: 2119–2130.
  41. van de Sluis B, Groot AJ, Vermeulen J, van der Wall E, van Diest PJ, et al. (2009) COMMD1 Promotes pVHL and O2-Independent Proteolysis of HIF-1alpha via HSP90/70. *PLoS ONE* 4: e3332.
  42. Salem Y, Shpungin S, Pasder O, Pomp O, Taler M, et al. (2005) Fer kinase sustains the activation level of ERK1/2 and increases the production of VEGF in hypoxic cells. *Cell Signal* 17: 341–353.
  43. Yuan WC, Lee YR, Huang SF, Lin YM, Chen TY, et al. (2011) A Cullin3-KLHL20 Ubiquitin ligase-dependent pathway targets PML to potentiate HIF-1 signaling and prostate cancer progression. *Cancer Cell* 20: 214–228.
  44. Tang TT, Lasky LA (2003) The forkhead transcription factor FOXO4 induces the down-regulation of hypoxia-inducible factor 1 alpha by a von Hippel-Lindau protein-independent mechanism. *J Biol Chem* 278: 30125–30135.
  45. Stoner M, Saville B, Wormke M, Dean D, Burghardt R, et al. (2002) Hypoxia induces proteasome-dependent degradation of estrogen receptor alpha in ZR-75 breast cancer cells. *Mol Endocrinol* 16: 2231–2242.
  46. Cho J, Kim D, Lee S, Lee Y (2005) Cobalt chloride-induced estrogen receptor alpha down-regulation involves hypoxia-inducible factor-1alpha in MCF-7 human breast cancer cells. *Mol Endocrinol* 19: 1191–1199.
  47. Tacchini L, Bianchi L, Bernelli-Zazzera A, Cairo G (1999) Transferrin receptor induction by hypoxia. HIF-1-mediated transcriptional activation and cell-specific post-transcriptional regulation. *J Biol Chem* 274: 24142–24146.
  48. Bakker WJ, Harris IS, Mak TW (2007) FOXO3a is activated in response to hypoxic stress and inhibits HIF1-induced apoptosis via regulation of CITED2. *Molecular cell* 28: 941–953.
  49. Qiang L, Wu T, Zhang HW, Lu N, Hu R, et al. (2012) HIF-1 alpha is critical for hypoxia-mediated maintenance of glioblastoma stem cells by activating Notch signaling pathway. *Cell Death and Differentiation* 19: 284–294.
  50. Gustafsson MV, Zheng X, Pereira T, Gradin K, Jin S, et al. (2005) Hypoxia requires notch signaling to maintain the undifferentiated cell state. *Dev Cell* 9: 617–628.
  51. Krishnamachary B, Zagzag D, Nagasawa H, Rainey K, Okuyama H, et al. (2006) Hypoxia-inducible factor-1-dependent repression of E-cadherin in von Hippel-Lindau tumor suppressor-null renal cell carcinoma mediated by TCF3, ZFX1A, and ZFX1B. *Cancer Res* 66: 2725–2731.
  52. Koshiji M, Kageyama Y, Pete EA, Horikawa I, Barrett JC, et al. (2004) HIF-1 alpha induces cell cycle arrest by functionally counteracting Myc. *Embo Journal* 23: 1949–1956.
  53. Pennacchietti S, Michieli P, Galluzzo M, Mazzone M, Giordano S, et al. (2003) Hypoxia promotes invasive growth by transcriptional activation of the met protooncogene. *Cancer Cell* 3: 347–361.
  54. Hayashi M, Sakata M, Takeda T, Tahara M, Yamamoto T, et al. (2005) Up-regulation of c-met protooncogene product expression through hypoxia-inducible factor-1alpha is involved in trophoblast invasion under low-oxygen tension. *Endocrinology* 146: 4682–4689.
  55. Horree N, Gort EH, van der Groep P, Heintz AP, Vooijs M, et al. (2008) Hypoxia-inducible factor 1 alpha is essential for hypoxic p27 induction in endometrioid endometrial carcinoma. *J Pathol* 214: 38–45.
  56. Moelans CB, de Weger RA, van Blokland MT, Ezendam C, Elshof S, et al. (2009) HER-2/neu amplification testing in breast cancer by multiplex ligation-dependent probe amplification in comparison with immunohistochemistry and in situ hybridization. *Cellular oncology : the official journal of the International Society for Cellular Oncology* 31: 1–10.
  57. van der Groep P, Bouter A, van der Zanden R, Menko FH, Buerger H, et al. (2004) Re: Germline BRCA1 mutations and a basal epithelial phenotype in breast cancer. *J Natl Cancer Inst* 96: 712–713; author reply 714.
  58. Vleugel MM, Greijer AE, Shvarts A, van der Groep P, van Berkel M, et al. (2005) Differential prognostic impact of hypoxia induced and diffuse HIF-1alpha expression in invasive breast cancer. *J Clin Pathol* 58: 172–177.
  59. van der Groep P, Bouter A, Menko FH, van der Wall E, van Diest PJ (2008) High frequency of HIF-1alpha overexpression in BRCA1 related breast cancer. *Breast Cancer Res Treat* 111: 475–480.
  60. Brenkman AB, de Keizer PL, van den Broek NJ, van der Groep P, van Diest PJ, et al. (2008) The peptidyl-isomerase Pin1 regulates p27kip1 expression through inhibition of Forkhead box O tumor suppressors. *Cancer Res* 68: 7597–7605.
  61. Gort EH, Suijkerbuijk KP, Roothaan SM, Raman V, Vooijs M, et al. (2008) Methylation of the TWIST1 promoter, TWIST1 mRNA levels, and immunohistochemical expression of TWIST1 in breast cancer. *Cancer Epidemiol Biomarkers Prev* 17: 3325–3330.
  62. Vermeulen JF, van de Ven RAH, Ercan C, van der Groep P, van der Wall E, et al. (2012) Nuclear Kaiso Expression Is Associated with High Grade and Triple-Negative Invasive Breast Cancer. *PLoS ONE* 7: e37864.
  63. van Diest PJ, Savulescu J (2002) For and against: No consent should be needed for using leftover body material for scientific purposes. *BMJ* 325: 648–651.
  64. Angus AG, Dalrymple D, Boulant S, McGivern DR, Clayton RF, et al. (2010) Requirement of cellular DDX3 for hepatitis C virus replication is unrelated to its interaction with the viral core protein. *J Gen Virol* 91: 122–132.
  65. Li CI, Uribe DJ, Daling JR (2005) Clinical characteristics of different histologic types of breast cancer. *Br J Cancer* 93: 1046–1052.
  66. Chen J, Gomes AR, Monteiro LJ, Wong SY, Wu LH, et al. (2010) Constitutively nuclear FOXO3a localization predicts poor survival and promotes Akt phosphorylation in breast cancer. *PLoS ONE* 5: e12293.