

Table S7. Metamorphic traits of *Rana temporaria* in the years 2009 and 2010. Given is the mean and standard deviation (SD) or coefficient of variance (CV), as well as the minimum and maximum size and weight of the measured individuals. Survival rate was based on 1117 eggs/clutch.

pond	year	clutches	metamorphs	survival	size				weight				development time				
			(total/meas)	[%]	mean±SD	min	max	CV	mean±SD	min	max	CV	mean±SD	min	max	CV	
													109.6 ±				
AW08	2009	60	2310 / 711	3.45	10.3 ± 0.6	8.7	11.8	0.06	0.09 ± 0.01	0.06	0.15	0.15	21.5	74	159	0.20	
AW08	2010	120	212 / 147	0.16	10.9 ± 0.9	8.9	14.2	0.08	0.11 ± 0.03	0.07	0.26	0.27	113.7 ± 9.7	99	144	0.08	
FS06	2009	33	812 / 290	2.20	14.4 ± 1.0	11.7	17.3	0.07	0.27 ± 0.06	0.15	0.46	0.21	99.7 ± 11.2	83	140	0.11	
													106.7 ±				
FS06	2010	86	2044 / 865	2.13	13.1 ± 0.9	8.9	16.6	0.07	0.20 ± 0.04	0.06	0.44	0.19	11.1	88	151	0.10	
WB04	2009	139	208 / 120	0.13	15.6 ± 0.7	14.1	17.2	0.04	0.38 ± 0.05	0.30	0.52	0.12	97.5 ± 4.0	91	106	0.04	
WB04	2010	382	101 / 64	0.02	14.4 ± 1.0	10.7	16.8	0.07	0.30 ± 0.06	0.11	0.44	0.19	100.8 ± 4.5	95	112	0.04	
WG02	2009	14	44 / 28	0.28	16.4 ± 0.9	14.5	18.6	0.05	0.43 ± 0.06	0.34	0.56	0.13	96.9 ± 4.5	90	107	0.05	
WG02	2010	11	6 / 6	0.05	15.2 ± 0.8	13.9	16.4	0.05	0.36 ± 0.03	0.32	0.40	0.08	98.3 ± 0.8	98	100	0.01	
WG07	2009	17	118 / 109	0.62	15.6 ± 1.0	12.6	17.5	0.06	0.39 ± 0.07	0.18	0.55	0.18	101.2 ± 9.2	88	128	0.09	
WG07	2010	25	54 / 37	0.19	16.9 ± 1.1	14.0	19.6	0.07	0.52 ± 0.11	0.22	0.78	0.21	109.3 ± 5.5	98	122	0.05	

metamorphs (total/meas) = number of metamorphs (emigrated/measured)