

Correction

# Correction: Secondary Household Transmission of 2009 Pandemic Influenza A (H1N1) Virus among an Urban and Rural Population in Kenya, 2009–2010

The PLOS ONE Staff

In Table 4, the second row in the column titled "Lwak" should be 22.5% (11/49), instead of 4.3% (21/486). Additionally, the third row of the column titled "Kibera" should be 4.3% (21/486), instead of 22.5% (11/49). Please see the correct Table 4 here:

**Table 4. Percentage (number) of Secondary Influenza-Like Illness Among Household Contacts of Confirmed pH1N1 patients, by Household Contact Age and by Patient Age - Kibera and Lwak, Kenya, 2009–2010.**

Age	Location	
	Kibera	Lwak
Household contact		
<5	6.1% (6/98)	22.5% (11/49)
≥5	4.3% (21/486)	5.9% (18/303)
Relative Risk (95% CI <sup>a</sup> )	1.4 (0.6–3.4)	3.8 (1.9–7.5)
Patient		
<5	5.0% (11/219)	3.2% (2/62)
5–15	3.3% (8/245)	11.2% (23/206)
≥15	6.7% (8/120)	4.8% (4/84)
P-value <sup>b</sup>	.33	.06

<sup>a</sup>CI = confidence interval.

<sup>b</sup>We calculated the p-values using chi-square tests.

doi:10.1371/journal.pone.0038166.t004

## Reference

- Kim CY, Breiman RF, Cosmas L, Audi A, Aura B, et al. (2012) Secondary Household Transmission of 2009 Pandemic Influenza A (H1N1) Virus among an Urban and Rural Population in Kenya, 2009–2010. PLoS ONE 7(6): e38166. doi:10.1371/journal.pone.0038166

**Citation:** The PLOS ONE Staff (2014) Correction: Secondary Household Transmission of 2009 Pandemic Influenza A (H1N1) Virus among an Urban and Rural Population in Kenya, 2009–2010. PLoS ONE 9(3): e91443. doi:10.1371/journal.pone.0091443

**Published:** March 26, 2014

**Copyright:** © 2014 The PLOS ONE Staff. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.