Correction

Correction: Statin Treatment in Hypercholesterolemic Men Does Not Attenuate Angiotensin II-Induced Ven constriction

The PLOS ONE Staff

Figure 5 is a duplicate of Figure 4. The authors have provided the correct Figure 5 here.

Figure 5. Time course of dorsal hand vein constriction with constant angiotensin II infusion. Infusion rate: 50 ng/min over 24 minutes before (pre) and after (post) treatment with irbesartan. Differences between pre- and post-treatment were analyzed with a two-way ANOVA with Bonferroni's post hoc tests to test for differences at single time points. Data are expressed as mean $\pm$ SEM. doi:10.1371/journal.pone.0103909.g005

Reference

1. Schindler C, Guenther K, Hermann C, Ferrario CM, Schroeder C, et al. (2014) Statin Treatment in Hypercholesterolemic Men Does Not Attenuate Angiotensin II-Induced Ven constriction. PLoS ONE 9(9): e103909. doi:10.1371/journal.pone.0103909

Citation: The PLOS ONE Staff (2014) Correction: Statin Treatment in Hypercholesterolemic Men Does Not Attenuate Angiotensin II-Induced Ven constriction. PLoS ONE 9(10): e112205. doi:10.1371/journal.pone.0112205

Published: October 23, 2014

Copyright: © 2014 The PLOS ONE Staff. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.