CORRECTION

Correction: Development of β -Carotene Rich Maize Hybrids through Marker-Assisted Introgression of *β -carotene hydroxylase* Allele

The *PLOS ONE* Staff

There is an error in [Fig. 1](#) in the article PDF. The authors have provided a corrected version here.

OPEN ACCESS

Citation: The *PLOS ONE* Staff (2015) Correction: Development of β -Carotene Rich Maize Hybrids through Marker-Assisted Introgression of *β -carotene hydroxylase* Allele. *PLoS ONE* 10(3): e0122130. doi:10.1371/journal.pone.0122130

Published: March 17, 2015

Copyright: © 2015 The PLOS ONE Staff. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Fig 1. Structure of the alleles of *crtRB1* gene causing variation in β -carotene concentration of maize. E: Exon; I: Intron; TE: Transposable element. A: No TE insertion at 6th Exon causing favourable *allele 1* (543 bp amplicon); B: 325 bp TE insertion at 6th Exon causing unfavourable *allele 2* (296+875 bp amplicon); C: 1250 bp TE insertion at 6th Exon causing unfavourable *allele 3* (296+1221+1880 bp amplicon).

doi:10.1371/journal.pone.0122130.g001

Reference

1. Muthusamy V, Hossain F, Thirunavukkarasu N, Choudhary M, Saha S, Bhat JS, et al. (2014) Development of β -Carotene Rich Maize Hybrids through Marker-Assisted Introgression of *β -carotene hydroxylase* Allele. PLoS ONE 9(12): e113583. doi: [10.1371/journal.pone.0113583](https://doi.org/10.1371/journal.pone.0113583) PMID: [25486271](https://pubmed.ncbi.nlm.nih.gov/25486271/)