CORRECTION

Correction: Detection of MPLW515L/K Mutations and Determination of Allele Frequencies with a Single-Tube PCR Assay

The *PLOS ONE* Staff

There are errors in the first sentence of the “Targeted deep sequencing” section of the Materials and Methods. The correct sentence is: A 169-bp fragment containing the human MPL sequence (43349278 to 43349446 of NC_000001.11) was PCR-amplified from genomic or standard plasmid DNA using forward (5'-TGACCGCTCTGCATCTAGTGC-3') and reverse (5'-GGTCACAGAGCGAACCAAGA-3') primers (see above).

There is an error in [Fig. 1](#). The first primer sequence in [Fig. 1A](#) is incorrect. Please view [Fig. 1](#) here.

OPEN ACCESS

Citation: The *PLOS ONE* Staff (2015) Correction: Detection of MPLW515L/K Mutations and Determination of Allele Frequencies with a Single-Tube PCR Assay. *PLoS ONE* 10(4): e0124208. doi:10.1371/journal.pone.0124208

Published: April 1, 2015

Copyright: © 2015 The PLOS ONE Staff. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

A

Name	Description	Sequence (5' → 3')
F_outer	Common forward primer	tgaccgctctgcatctagtc
F_inner	W515L specific forward primer	ggcctgctgctgctgaAgtt
R_inner	W515K specific reverse primer	FAM - acctgtagtgtgcaggaaactTc tt
R_outer	Common reverse primer	FAM - ggtcacagagcgaaccaaga

B

C

Fig 1. Detection of *MPLW515L/K* mutations using DARMS-PCR. (A) The primers used in the DARMS-PCR assay. The two inner primers harbored sequences (underlined) that matched *MPLW515L* or *W515K*, but not the wild-type allele. Other mismatches (capital letters) were introduced into the inner primers to reduce the annealing of the mutant-specific primers to the wild-type sequence. The reverse primers were labeled with FAM (5-carboxyfluorescein hydrate) at the 5' terminus. (B) A schematic representation of DARMS-PCR products. The two outer primers were designed to generate a 169-base-pair (bp) PCR product from all *MPL* alleles. The F_inner and R_inner primers annealed specifically to the *MPLW515L* and *W515K* alleles, respectively; in combination with the outer primers, they generated 84- and 128-bp PCR products, respectively. From a mutant allele, both 169-bp and 84- or 128-bp fragments were amplified, while, only the 169-bp fragment was generated from the wild-type allele. (C) Demonstration of DARMS-PCR. A capillary electropherogram of DARMS-PCR products showing three peaks derived from wild-type *MPL*, *W515L*, and *W515K*. This result was obtained when PCR was performed with a standard DNA mixture containing equal ratios of *MPL* wild-type, *W515L*, and *W515K* alleles with a total copy number of 10^5 . The horizontal axis represents the fragment length, and the vertical axis represents the fluorescence intensity.

doi:10.1371/journal.pone.0124208.g001

Reference

1. Takei H, Morishita S, Araki M, Edahiro Y, Sunami Y, Hironaka Y, et al. (2014) Detection of *MPLW515L/K* Mutations and Determination of Allele Frequencies with a Single-Tube PCR Assay. PLoS ONE 9(8): e104958. doi: [10.1371/journal.pone.0104958](https://doi.org/10.1371/journal.pone.0104958) PMID: [25144224](https://pubmed.ncbi.nlm.nih.gov/25144224/)