

CORRECTION

Correction: Heart Failure in a Cohort of Patients with Chronic Kidney Disease: The GCKD Study

Hanna Beck, Stephanie I. Titze, Silvia Hübner, Martin Busch, Georg Schlieper, Ulla T. Schultheiss, Christoph Wanner, Florian Kronenberg, Vera Krane, Kai-Uwe Eckardt, Anna Kötgen, GCKD Investigators

There is an error in [Table 5](#) in the third column's second row. The correct p-value should be 0.495. Please see the corrected [Table 5](#) here.



OPEN ACCESS

Citation: Beck H, Titze SI, Hübner S, Busch M, Schlieper G, Schultheiss UT, et al. (2015) Correction: Heart Failure in a Cohort of Patients with Chronic Kidney Disease: The GCKD Study. PLoS ONE 10(6): e0131034. doi:10.1371/journal.pone.0131034

Published: June 15, 2015

Copyright: © 2015 Beck et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Table 5. Multivariable adjusted analyses of factors associated with Gothenburg HF and self-reported HF (n = 4,604)

	Gothenburg HF			Self-reported HF		
	OR	95% CI	P	OR	95% CI	P
eGFR (ml/min/1.73 m²)						
≥90	0.87	0.59–1.29	0.495	0.56	0.27–1.14	0.111
60–89	1.07	0.88–1.31	0.485	1.16	0.90–1.49	0.264
45–59 (reference)	ref	ref	ref	ref	ref	ref
30–44	1.18	1.01–1.39	0.036	1.12	0.92–1.36	0.253
<30	1.71	1.32–2.20	<0.001	1.29	0.96–1.74	0.088
UACR (mg/g) (reference: <30)						
30–299	0.81	0.69–0.95	0.010	0.87	0.71–1.05	0.147
≥300	0.87	0.72–1.04	0.126	0.70	0.55–0.88	0.002
Age (5 year intervals)						
Male gender	1.13	1.09–1.17	<0.001	1.15	1.10–1.22	<0.001
Diabetes mellitus	0.65	0.57–0.76	<0.001	1.01	0.84–1.21	0.930
Hypertension	1.64	1.42–1.90	<0.001	1.61	1.35–1.92	<0.001
Valvular heart disease	1.81	1.27–2.59	0.001	1.90	1.11–3.26	0.020
BMI (kg/m ²)	2.52	2.01–3.15	<0.001	3.98	3.19–4.97	<0.001
Sleep apnea	1.10	1.09–1.11	<0.001	1.04	1.03–1.06	<0.001
Anemia	2.22	1.76–2.81	<0.001	2.15	1.70–2.73	<0.001
Education (reference: ≤9 years)	1.39	1.18–1.63	<0.001	1.06	0.88–1.29	0.534
10 years	ref	ref	ref	ref	ref	ref
>10 years	0.84	0.72–0.99	0.037	0.94	0.77–1.15	0.534
10 years	0.69	0.56–0.84	<0.001	1.00	0.78–1.29	0.977
Serum albumin (g/l)	0.95	0.94–0.97	<0.001	0.99	0.98–1.02	0.993
Heart rate (bmp)	0.99	0.99–1.00	0.584	0.99	0.98–0.99	0.011
Current smoker	0.97	0.80–1.17	0.752	0.98	0.76–1.26	0.864
Alcohol intake (≥ 3 times per week)	0.98	0.82–1.17	0.827	0.97	0.79–1.21	0.802

Of 5015 observations, values were missing in BMI (57), valvular heart disease (42), anemia (145), serum albumin (1), education (101), heart rate (49), current smoker (12), alcohol intake (28).

doi:10.1371/journal.pone.0131034.t001

Reference

- Beck H, Titze SI, Hübner S, Busch M, Schlieper G, Schultheiss UT, et al. (2015) Heart Failure in a Cohort of Patients with Chronic Kidney Disease: The GCKD Study. PLoS ONE 10(4): e0122552. doi: [10.1371/journal.pone.0122552](https://doi.org/10.1371/journal.pone.0122552) PMID: 25874373