CORRECTION

Correction: Combined Use of Hyperbaric and Hypobaric Ropivacaine Significantly Improves Hemodynamic Characteristics in Spinal Anesthesia for Caesarean Section: A Prospective, Double-Blind, Randomized, Controlled Study

The *PLOS ONE* Staff

The affiliation for the first two authors is incorrect. ZheFeng Quan and Ming Tian are not affiliated with #1 but with #2 Department of Anesthesia, Beijing Friendship Hospital, Capital Medical University, Beijing, 100050, China. The publisher apologizes for the error.

Reference

1. Quan Z, Tian M, Chi P, Li X, He H, Luo C (2015) Combined Use of Hyperbaric and Hypobaric Ropivacaine Significantly Improves Hemodynamic Characteristics in Spinal Anesthesia for Caesarean Section: A Prospective, Double-Blind, Randomized, Controlled Study. *PLoS ONE* 10(5): e0125014. doi: [10.1371/journal.pone.0125014](https://doi.org/10.1371/journal.pone.0125014) PMID: [25970485](https://pubmed.ncbi.nlm.nih.gov/25970485/)

OPEN ACCESS

Citation: The *PLOS ONE* Staff (2015) Correction: Combined Use of Hyperbaric and Hypobaric Ropivacaine Significantly Improves Hemodynamic Characteristics in Spinal Anesthesia for Caesarean Section: A Prospective, Double-Blind, Randomized, Controlled Study. *PLoS ONE* 10(6): e0132082. doi:10.1371/journal.pone.0132082

Published: June 25, 2015

Copyright: © 2015 The PLOS ONE Staff. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.