CORRECTION

Correction: Hepatocyte-Specific Depletion of UBXD8 Induces Periportal Steatosis in Mice Fed a High-Fat Diet

Norihiro Imai, Michitaka Suzuki, Kazuhiko Hayashi, Masatoshi Ishigami, Yoshiki Hirooka, Takaya Abe, Go Shioi, Hidemi Goto, Toyoshi Fujimoto

There is an error in the second sentence in the PCR section of Materials and Methods. The correct sentence is: Primers were as follows: for loxP, forward primer: 5'-CCAACCTCTGTGGG TCCTC-3'; reverse primer: 5'-GCGATTGGGGATCTGAGTTA-3'; for albumin-Cre, forward primer: 5'-TTTGCCTGCATTACCGGTCGATGCAAC-3'; reverse primer: 5'-TGCCCCTGTT TCACTATCCAGGTTACGGA-3'.

Reference

1. Imai N, Suzuki M, Hayashi K, Ishigami M, Hirooka Y, Abe T, et al. (2015) Hepatocyte-Specific Depletion of UBXD8 Induces Periportal Steatosis in Mice Fed a High-Fat Diet. PLoS ONE 10(5): e0127114. doi: [10.1371/journal.pone.0127114](https://doi.org/10.1371/journal.pone.0127114) PMID: [25970332](https://pubmed.ncbi.nlm.nih.gov/25970332/)

OPEN ACCESS

Citation: Imai N, Suzuki M, Hayashi K, Ishigami M, Hirooka Y, Abe T, et al. (2016) Correction: Hepatocyte-Specific Depletion of UBXD8 Induces Periportal Steatosis in Mice Fed a High-Fat Diet. PLoS ONE 11(1): e0148004. doi:10.1371/journal.pone.0148004

Published: January 25, 2016

Copyright: © 2016 Imai et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.