

RESEARCH ARTICLE

Naringenin Inhibits Superoxide Anion-Induced Inflammatory Pain: Role of Oxidative Stress, Cytokines, Nrf-2 and the NO-cGMP-PKG-K_{ATP} Channel Signaling Pathway

Marília F. Manchope¹, Cássia Calixto-Campos¹, Letícia Coelho-Silva¹, Ana C. Zarpelon¹, Felipe A. Pinho-Ribeiro¹, Sandra R. Georgetti², Marcela M. Baracat², Rúbia Casagrande², Waldiceu A. Verri, Jr^{1*}

1 Departamento de Ciências Patológicas, Centro de Ciências Biológicas, Universidade Estadual de Londrina, Londrina, Brazil, **2** Departamento de Ciências Farmacêuticas, Centro de Ciências de Saúde, Universidade Estadual de Londrina, Londrina, Brazil

* waverri@uel.br



OPEN ACCESS

Citation: Manchope MF, Calixto-Campos C, Coelho-Silva L, Zarpelon AC, Pinho-Ribeiro FA, Georgetti SR, et al. (2016) Naringenin Inhibits Superoxide Anion-Induced Inflammatory Pain: Role of Oxidative Stress, Cytokines, Nrf-2 and the NO-cGMP-PKG-K_{ATP} Channel Signaling Pathway. PLoS ONE 11(4): e0153015. doi:10.1371/journal.pone.0153015

Editor: Bernhard Ryffel, French National Centre for Scientific Research, FRANCE

Received: February 9, 2016

Accepted: March 22, 2016

Published: April 5, 2016

Copyright: © 2016 Manchope et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: Regarding the authors' Funding Statement/ Financial Disclosure, the authors inform that "Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Ministério da Ciência, Tecnologia e Inovação (MCTI), Secretaria da Ciência, Tecnologia e Ensino Superior (SETI), Fundação Araucária and Parana State Government grants supported this

Abstract

In the present study, the effect and mechanism of action of the flavonoid naringenin were evaluated in superoxide anion donor (KO₂)-induced inflammatory pain in mice. Naringenin reduced KO₂-induced overt-pain like behavior, mechanical hyperalgesia, and thermal hyperalgesia. The analgesic effect of naringenin depended on the activation of the NO-cGMP-PKG-ATP-sensitive potassium channel (K_{ATP}) signaling pathway. Naringenin also reduced KO₂-induced neutrophil recruitment (myeloperoxidase activity), tissue oxidative stress, and cytokine production. Furthermore, naringenin downregulated KO₂-induced mRNA expression of gp91phox, cyclooxygenase (COX)-2, and preproendothelin-1. Besides, naringenin upregulated KO₂-reduced nuclear factor (erythroid-derived 2)-like 2 (Nrf2) mRNA expression coupled with enhanced heme oxygenase (HO-1) mRNA expression. In conclusion, the present study demonstrates that the use of naringenin represents a potential therapeutic approach reducing superoxide anion-driven inflammatory pain. The antinociceptive, anti-inflammatory and antioxidant effects are mediated via activation of the NO-cGMP-PKG-K_{ATP} channel signaling involving the induction of Nrf2/HO-1 pathway.

Introduction

Pain is an unpleasant sensory and emotional experience, generally in association with tissue injury. During inflammation, pro-inflammatory mediators activate resident cells, recruited cells and nociceptors, thereby driving pain signaling. Nociceptive neurons do not express receptors for all inflammatory molecules, suggesting both direct and indirect activation and

study (Brazil). A.C.Z. received a postdoctoral fellowship from Fundação Araucária/CAPES (Brazil). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The specific roles of these authors are articulated in the 'author contributions' section.

Competing Interests: The authors have declared that no competing interests exist.

sensitization of nociceptors [1]. Increased levels of oxidative stress during the inflammatory response also contribute to nociception. For instance, reactive oxygen species (ROS) and reactive nitrogen species (RNS) can act directly and indirectly to induce nociceptor sensitization and activation [2–5]. The superoxide anion (O_2^-) is a common form of ROS that can drive nociception [5,6]. O_2^- reacts with nitric oxide (NO) generating peroxynitrite, which also contributes to nociception [3]. Superoxide dismutase (SOD), an enzymatic antioxidant, transforms superoxide anion in hydrogen peroxide, which may also induce nociception [2]. Therefore, O_2^- is a crucial ROS to the biological underpinnings driving nociception. O_2^- increases other pro-inflammatory effects, including increasing vascular permeability [7], inducing cytokine release [8,9] and increasing neutrophil recruitment [9,10], as well as provoking overt pain-like behavior and hyperalgesia [2–5]. In a physiological state, O_2^- levels remain under control by the action of the endogenous antioxidant systems, including SOD, and the endogenous antioxidant reduced glutathione (GSH) [9]. However, the imbalance between oxidants and antioxidants during inflammation leads to oxidative stress. This is important, as inhibiting the production of pro-inflammatory cytokines and ROS limit the development of inflammatory pain [6,11–13].

Naringenin (4',5,7-trihydroxy-flavonone) is a flavonoid which belongs to flavonones class found in citrus fruits, including lemon, orange, tangerine and grapefruit [14]. Naringenin inhibits the nociceptive responses in models of formalin-, acetic acid- and capsaicin-induced overt pain, as well as neuropathic pain [15–17]. Naringenin also inhibits inflammation by targeting cyclooxygenase (COX)-2 in ethanol-induced liver injury [18] and *in vitro* [19]. Moreover, naringenin inhibits phosphorylation of nuclear factor kappaB (NFκB) subunit p65 and mitogen-activated protein kinases (MAPK) in daunorubicin-induced nephrotoxicity [20] as well as inhibiting the EGFR-PI3K-Akt/ERK MAPK signaling pathway in human airway epithelial cells [21]. Naringenin also inhibits a number of aspects of oxidative stress, including lipid peroxidation and O_2^- production, as well as restoring GSH levels in UVB-induced oxidative stress in the skin of Hairless mice [22]. Furthermore, naringenin increases SOD in an experimental stroke model, highlighting its wide-acting induction of endogenous antioxidants [23]. In agreement with such antioxidant effects, naringenin also induces nuclear factor (erythroid-derived 2)-like 2 (Nrf2)/ heme oxygenase (HO)-1 in CCl_4 -induced hepatic inflammation [24]. Some flavonoids can induce antinociception by activating the NO–cGMP–PKG–ATP-sensitive potassium channel (K_{ATP}) signaling pathway [25–28]. Activating this signaling pathway is an important mechanism of action of a number of clinical analgesics, such as opioids [25], and some non-steroidal anti-inflammatory drugs including dipyrrone [26], diclofenac [27], and indomethacin [28].

Given the above, the current study addresses the analgesic effects of naringenin in a model of O_2^- -triggered inflammatory pain. It was also investigated as to whether naringenin's mechanism of action involves the NO–cGMP–PKG– K_{ATP} channel signaling pathway, the regulation of inflammatory mediators/enzymes and oxidative stress as well as the transcription factor Nrf2, and its downstream target, HO-1.

Materials and Methods

Animals

Male Swiss mice (25 ± 5 g) from Londrina State University were housed in standard plastic cages with free access to food and water, with a light/dark cycle of 12:12 h, at 21°C. All behavioral testing was performed between 9 a.m. and 5 p.m. in a temperature-controlled room. At the end of experiments, mice were anesthetized with isoflurane 3% to minimize suffering (Abbott Park, IL, USA) and killed by cervical dislocation followed by decapitation. The animal

condition was monitored daily and at indicated time points during the experiments. No unexpected animal deaths occurred during this study. Animals' care and handling procedures were in accordance with the International Association for Study of Pain (IASP) guidelines and with the approval of the Ethics Committee of the Londrina State University (process 133666.2013.71).

Drugs and reagents

Potassium superoxide (KO_2) (Alfa Aesar, 96,5%, Ward Hill, MA, USA). Naringenin (Santa Cruz Biotechnology, Inc., 98%, Dalla, TX, USA). Saline (NaCl 0,9%; Fresenius Kabi Brazil Ltda. Aquiraz, CE, Brazil). L-NAME (Research Biochemicals, Natick, MA, USA), KT5823 (Calbiochem, San Diego, CA, USA), ODQ [1H-(1,2,4)-oxadiazolol-(4,3-a)-quinoxalin-1-one, Tocris Cookson, Baldwin, MO, USA]. Glybenclamide, HTAB (Bromide, hexadecyl trimethyl ammonium), dihydrochloride O-dianisidine, GSH (reduced glutathione), EDTA sodium salt, ferric chloride hexahydrate, TPTZ (2,4,6-tripiridil-s-triazine) and Trolox (6-hydroxy-2,5,7,8-tetramethylchroman-2-11 carboxylic) were purchased from Sigma Chemical Co. (St. Louis, MO, USA). DMSO 2% and Tween 80 (Química Moderna, Barueri, SP, Brazil).

Experimental procedures

Mice were pretreated per orally (po) with 16.7, 50, or 150 mg/kg of naringenin or with vehicle (saline) 1 h before intraplantar (ipl) or intraperitoneal (ip) injection of 30 μg or 1 mg of KO_2 , respectively [6]. The writhing response was evaluated between 0–20 min after KO_2 ip injection and the paw flinching and licking nociceptive responses were quantified during 30 min after ipl injection of KO_2 . Mechanical and thermal hyperalgesia were evaluated 0.5–7 h after KO_2 . Inflammatory stimulation with KO_2 induced mechanical and thermal hyperalgesia only in the ipsilateral paw, in the side where the stimulus was injected. Myeloperoxidase (MPO) activity was evaluated 7 h after KO_2 administration. Mice were pretreated with inhibitors of the NO synthase (L-NAME; 90 mg/kg, ip, 1 h pre-treatment), guanylate cyclase (ODQ, 0,3 mg/kg, ip, 30 min pre-treatment), PKG (KT5823, 0,5 $\mu\text{g}/\text{animal}$, ip, 5 min pre-treatment), K_{ATP} channels (glibenclamide, 0,3 mg/kg, po, 45 min pre-treatment) before naringenin treatment (50 mg/kg, po). After 1 h mice received ipl injection of 30 μg of KO_2 , and mechanical and thermal hyperalgesia were assessed 0.5–7 h thereafter. Evaluations of oxidative stress (GSH, FRAP, TBARS, NBT) and cytokine production (tumor necrosis factor [TNF] α and interleukin [IL]-10) were carried out, as well as RT-qPCR measures of gp91^{phox}, COX-2, prepro endothelin (ET)-1, Nrf2, HO-1, and IL-33, 3 h after KO_2 injection in samples of paw skin.

Overt pain-like behavioral tests

Abdominal writhing was induced by ip injection of 1 mg of KO_2 [6]. Immediately after stimulus injection, each mouse was placed individually in a large glass cylinder, and the intensity of nociceptive behavior was quantified by counting the total number of writhings occurring between 0 and 20 min after stimulus injection. The writhing response consists of a contraction of the abdominal muscle together with a stretching of hind limbs, and the intensity was expressed as the cumulative number of abdominal contortions over 20 min. The number of paw flinches and the time spent licking the paw were determined between 0–30 min after ipl injection of 30 μg of KO_2 . Each mouse was placed in a large glass cylinder immediately after stimulus injection. The intensity of nociceptive behavior was quantified by counting the total number of paw flinches and the time (seconds) spent licking the ipsilateral paw [6].

Mechanical hyperalgesia test

Mechanical hyperalgesia was evaluated by the electronic version of von Frey's test, as reported previously [29]. In a quiet, temperature controlled room, mice were placed in acrylic cages (12 x 10 x 17 cm) with wire grid floors, 15–30 min before the start of testing. The test consisted of evoking a hind paw flexion reflex with a handheld force transducer (electronic anesthesiometer, IITC Life Science, Woodland Hills, CA) adapted with a 0.5 mm² polypropylene tip. The investigator was trained to apply the tip perpendicularly to the central area of the plantar hind paw with a gradual increase in pressure. The gradual increase in pressure was manually performed in blinded experiments. The upper limit pressure was 15 g. The end-point was characterized by the removal of the paw followed by clear flinching movements. After paw withdrawal, the intensity of the pressure was automatically recorded, and the final value for the response was obtained by averaging three measurements. The animals were tested before and after treatments. The results are expressed by delta (Δ) withdrawal threshold (in grams) calculated by subtracting the mean measurements 0.5, 1, 3, 5 and 7 h after stimulus from the zero-time mean measurements [6].

Thermal hyperalgesia test

Heat thermal hyperalgesia was performed using a hot plate at $55 \pm 1^\circ\text{C}$ [6]. The end-point was characterized by the removal of the paw followed by clear paw flinching or licking movements. The upper time was 15 s to avoid possible injury. The results are expressed by total withdrawal latency (in seconds) of measurements obtained 0.5, 1, 3, 5 and 7 h after stimulus [30].

MPO assay

Neutrophil migration to the hind paw skin tissue was evaluated using an MPO kinetic-colorimetric assay, as described previously [6]. Samples of paw skin tissue were collected 7 h after stimulus in ice-cold 50 mM K₂HPO₄ buffer (pH 6.0) containing 0.5% hexadecyltrimethylammonium bromide (HTAB) and kept at -80°C until use. Samples were homogenized, centrifuged (16,100g x 2 min, 4°C), and the resulting supernatant was assayed for MPO activity spectrophotometrically at 450 nm (Multiskan GO Microplate Spectrophotometer, Thermo Scientific, Vantaa, Finland), with three readings in 1 min. The MPO activity of samples was compared to a standard curve of neutrophils. Briefly, 15 μL of sample was mixed with 200 μL of 50 mM phosphate buffer, pH 6.0, containing 0.167 mg/mL o-dianisidine dihydrochloride and 0.015% hydrogen peroxide. The results are presented as MPO activity (number of neutrophils $\times 10^{10}$ per g of tissue).

GSH measurement

Paw skin sample was collected and maintained at -80°C for at least 48 h. Sample was homogenized with 200 μL of 0.02 M EDTA. The homogenate was mixed with 25 μL of 50% trichloroacetic acid and was homogenized three times during 15 min. The mixture was centrifuged (15 min x 1500 g x 4°C). The supernatant was added to 200 μL of 0.2 M TRIS buffer, pH 8.2, and 10 μL of 0.01M DTNB. After 5 min, the absorbance was measured at 412 nm against a reagent blank with no supernatant. A standard curve was performed with standard GSH. The results are expressed as GSH nmol per mg paw [31].

FRAP assay

Paw skin sample was collected and immediately homogenized with 500 μL of 1.15% KCl, and was centrifuged (10 min x 200 g x 4°C). The ability of the sample to resist oxidative damage

was determined using FRAP assays [32]. For the FRAP assay, 50 μL of supernatant was mixed with 150 μL of deionized water and 1.5 mL of freshly prepared FRAP reagent. The reaction mixture was incubated at 37°C for 30 min and the absorbance was measured at 595 nm. The results were equated against a Trolox standard curve (1.5–30 $\mu\text{mol/L}$, final concentrations). The results are expressed as Trolox equivalents per mg of paw.

Superoxide anion production

The quantitation of O_2^- production in tissue homogenates was performed using the NBT assay, as described previously [33]. Skin samples were collected 3h after the stimulus. Briefly, 50 μL of the homogenate was incubated with 100 μL of NBT (1 mg/mL) in 96-well plates at 37°C for 1h. The supernatant was carefully removed and the reduced formazan solubilized by adding 120 μL of 2M KOH and 140 μL of DMSO. The NBT reduction was measured at 600 nm using a microplate spectrophotometer reader (Multiskan GO, Thermo Scientific). The tissue weight was used for data normalization; thus the results are expressed as NBT reduction (OD/mg of paw).

Lipid Peroxidation

Tissue lipid peroxidation was assessed by the levels of thiobarbituric acid reactive substances (TBARS) [34]. For this assay, TCA 10% was added to the homogenate and the mixture was centrifuged (1000 g, 3 min, 4°C) to precipitate proteins. The protein-free supernatant was then separated and mixed with TBA (0.67%). The mixture was kept in water bath (15 min, 100°C). Malondialdehyde (MDA), an intermediate product of lipid peroxidation, was determined by difference between absorbances at 535 and 572 nm using a microplate spectrophotometer reader. The results were presented as nmol of MDA per mg of paw [34].

Cytokine measurement

Paw skin samples were collected 3 h after the injection of KO_2 , homogenized in 500 μL of ice-cold buffer containing protease inhibitors, centrifuged (3000 rpm x 10 min x 4°C) and the supernatants used to measure, TNF α and IL-10 levels, by an enzyme-linked immunosorbent assay (ELISA) using eBioscience kits. As a control, the concentrations of these cytokines were determined in animals injected with saline. The results are expressed as picograms (pg) of cytokine per mg of paw.

Reverse transcription and quantitative polymerase chain reaction (RT-qPCR)

RT-qPCR was performed as previously described [33]. Paw skin samples were collected 3 h after stimulus and homogenized in trizol reagent, with the total RNA being isolated according to manufacturer's directions. The purity of total RNA was measured with a spectrophotometer with the wavelength absorption ratio (260/280 nm) being between 1.8 and 2.0 for all preparations. Reverse transcription of total RNA to cDNA, and qPCR were carried out using GoTaq[®] 2-Step RT-qPCR System (Promega) following the manufacturer's instructions. The relative gene expression was measured using the comparative $2^{-\Delta\Delta\text{C}_q}$ method. The primers used were gp^{91phox}, sense: 5' -AGCTATGAGGTGGTGATGTTAGTGG-3', antisense: 5' -CACAAATATTTGTAC CAGACAGACTTGAG-3'; IL-33, sense: 5' -TCCTTGCTTGGCAGTATCCA-3', antisense: 5' -TGCTCAATGTGTCAACAGACG-3'; COX-2, sense: 5' -GTGGAAAAACCTCGTC CAGA-3', antisense: 5' -GCTCGGCTTC CAGTATTGAG-3'; preproET-1, sense: 5' -TGTGTCTACTTCTGCCACCT-3', antisense: 5' -CACCAGCTGCTGATAGATAC-3'; Nrf2, sense:

5' -TCACACGAGATGAGCTTAGGGCAA-3' , antisense: 5' -TACAGTTCTGGG CGGCGACTT TAT-3' ; HO-1, sense: 5' -CCCAAACCTGGCCTGTAAAA-3' , antisense: 5' -CGTGGTCAGT CAACATGGAT-3' ; β -actin, sense: 5' -AGCTGCGTTT TACACCCTTT-3' , antisense: 5' -AA GCCATGCCAATGTTGTCT-3' . The expression of β -actin mRNA was used as a reference gene to normalize data.

Statistical analysis

Results are presented as means \pm SEM of measurements made on six mice in each group per experiment, and are representative of two independent experiments. Two-way analysis of variance (ANOVA) was used to compare the groups and doses at all times (curves), when the hyperalgesic responses were measured at different times after the administration or enforcement of the stimuli. The factors analyzed were treatment, time, and time versus treatment interaction. When there was a significant time versus treatment interaction, one-way ANOVA followed by Tukey's post hoc was performed on each occasion. On the other hand, when the hyperalgesic responses were measured once after the administration or enforcement of the stimuli, the difference between responses were evaluated by one-way ANOVA followed by Tukey's post hoc. Statistical differences were considered to be significant at $p < 0.05$.

Results

Naringenin inhibits KO_2 -induced overt pain-like behavior

Mice received naringenin (16.7, 50, 150 mg/kg, po) treatment 1 h before injection of 1 mg of KO_2 ip for the assessment of the total number of writhings, or received 30 μ g of KO_2 ipl for the evaluation of the total number of paw flinches and time spent licking the paw. The naringenin doses of 50 and 150 mg/kg inhibited KO_2 -induced writhing response at a similar magnitude of effect, with the 3 mg/kg dose showing no statistically significant effect (Fig 1a). The dose of 50 mg/kg of naringenin was therefore selected for the next experiments. Naringenin also inhibited KO_2 -induced paw flinches (Fig 1b) and time spent licking the paw (Fig 1c).

Naringenin inhibits KO_2 -induced mechanical hyperalgesia, thermal hyperalgesia, and MPO activity

Mice received naringenin (50 mg/kg, po) treatment 1 h before KO_2 injection (30 μ g, ipl). Mechanical and thermal hyperalgesia were assessed 0.5, 1, 3, 5 and 7 h after KO_2 injection. Naringenin inhibited KO_2 -induced mechanical and thermal hyperalgesia at all time points (Fig 2a and 2b). Naringenin also reduced KO_2 -induced increased of MPO activity at 7 h (Fig 2c).

Naringenin inhibits KO_2 -induced mechanical and thermal hyperalgesia by activating the NO-cGMP-PKG- K_{ATP} channel signaling pathway

Mice were treated with inhibitors of a) NO synthase (L-NAME; 90 mg/kg, ip, 1 h pre-treatment), b) guanylate cyclase (ODQ, 0.3 mg/kg, ip, 30 min pre-treatment), c) PKG (KT5823, 0.5 μ g/animal, ip, 5 min pre-treatment), and d) the K_{ATP} channel (glibenclamide, 0.3 mg/kg, po, 45 min pre-treatment) before naringenin (50 mg/kg, po, 1 h before KO_2 injection) administration. After 1 h naringenin treatment, mice received a KO_2 ipl injection. L-NAME (Fig 3a and 3b), ODQ (Fig 3c and 3d), KT5823 (Fig 3e and 3f), and glibenclamide (Fig 3g and 3h) inhibited the analgesic effect of naringenin in KO_2 -induced mechanical and thermal hyperalgesia. Therefore, the anti-hyperalgesic mechanism of naringenin depends, at least in part, on activating the NO-cGMP-PKG- K_{ATP} channel signaling pathway.




Fig 1. Naringenin inhibits KO_2 -induced overt pain-like behavior. (a-c) Mice received naringenin (16.7, 50 and 150 mg/kg, po) treatment 1 h before ip injection of 1 mg of KO_2 or ipl injection of 30 μ g of KO_2 . The total number of writhings was evaluated 0–20 min after ip injection of KO_2 . (b) The number of paw flinches and (c) time spent licking the paw were evaluated 0–30 min after ipl injection of KO_2 . Results are mean \pm SEM of 6 mice per group per experiment, and are representative of 2 independent experiments. * $p < 0.05$ vs. saline group, # $p < 0.05$ vs. KO_2 group. One-way ANOVA followed Tukey's post hoc.

doi:10.1371/journal.pone.0153015.g001




Fig 2. Naringenin inhibits KO₂-induced mechanical hyperalgesia, thermal hyperalgesia and myeloperoxidase (MPO) activity. (a-c) Mice received naringenin (50 mg/kg, po) treatment 1 h before ipl injection of 30 μg of KO₂. Mechanical (a) and thermal (b) hyperalgesia were evaluated between 0.5–7 h after ipl injection of KO₂. (c) At the 7th h after KO₂ injection, paw skin samples were collected for MPO activity assay. Results are mean ± SEM of 6 mice per group per experiment, and are representative of 2 independent experiments. **p* < 0.05 vs. saline group, #*p* < 0.05 vs. KO₂ group. Repeated measures two-way ANOVA for hyperalgesia data and One-way ANOVA for MPO activity followed Tukey's post hoc.

doi:10.1371/journal.pone.0153015.g002




Fig 3. Naringenin inhibits KO₂-induced mechanical and thermal hyperalgesia by activating NO/cGMP/PKG/K_{ATP} channel signaling pathway. Mice received (a,b) L-NAME (NOS inhibitor, 90 mg/kg, ip, 1 h), (c,d) ODQ (guanylate cyclase inhibitor, 0.3 mg/kg, ip, 30 min), (e,f) KT5823 (PKG inhibitor, 0.5 μg/mice, ip, 5 min), or (g,h) glibenclamide (K_{ATP} channel inhibitor, 0.3 mg/kg, po, 45 min) treatment before administration of naringenin (50 mg/kg, po). (a-h) 1 h after naringenin administration, mice received an ipl injection of 30 μg of KO₂. Mechanical (a,c,e,g) and thermal (b,d,f,h) hyperalgesia were evaluated between 0.5–7 h after the ipl

injection of KO_2 . Results are mean \pm SEM of 6 mice per group per experiment, and are representative of 2 independent experiments. * $p < 0.05$ vs. saline group, # $p < 0.05$ vs. KO_2 group. Repeated measures two-way ANOVA followed Tukey's post hoc.

doi:10.1371/journal.pone.0153015.g003

Naringenin inhibits KO_2 -induced oxidative stress

Mice received naringenin (50 mg/kg, po) treatment 1h before KO_2 injection (30 μg , ipl). Samples of paw skin were collected after 3 h for colorimetric assays and RT-qPCR. KO_2 depleted paw skin GSH (Fig 4a) and antioxidant capacity (Fig 4b); and increased lipid peroxidation (Fig 4c) and superoxide anion production (Fig 4d) in the paw skin. Naringenin treatment inhibited these deleterious effects of KO_2 by restoring GSH levels (Fig 4a), and antioxidant capacity (Fig 4b), and inhibiting lipid peroxidation (Fig 4c) and O_2^- production (Fig 4d). Corroborating the results on oxidative stress (Fig 4a–4d), naringenin also inhibited KO_2 -induced gp91^{phox} mRNA expression, a component of NADPH oxidase, an important source of O_2^- production (Fig 4e).

Naringenin inhibits KO_2 -induced cytokine production and mRNA expression

The protocol of this section was the same as for Fig 4, with samples analyzed by ELISA and RT-qPCR. Naringenin treatment inhibited KO_2 -induced TNF α (Fig 5a) and IL-10 (Fig 5b) production, and IL-33 mRNA expression (Fig 5c).

Naringenin inhibits KO_2 -induced COX-2 mRNA expression

The protocol of this section was the same as for Fig 4, with samples analyzed by RT-qPCR. Naringenin treatment inhibited KO_2 -induced COX-2 mRNA expression (Fig 6).

Naringenin inhibits KO_2 -induced preproET-1 mRNA expression

The protocol of this section was the same as for Fig 4, and samples analyzed by RT-qPCR. Naringenin treatment inhibited KO_2 -induced preproET-1 mRNA expression (Fig 7).

Naringenin increases Nrf2/HO-1 mRNA expression

The protocol of this section was the same as for Fig 4, with samples analyzed by RT-qPCR. Naringenin inhibited the KO_2 -induced decrease in Nrf2 mRNA expression (Fig 8a), whilst enhancing KO_2 -induced HO-1 mRNA expression (Fig 8b).

Discussion

Pain is a multisensory experience, which occurs in response to stimulation of the channels/receptors that depolarizes nociceptor terminals, thereby generating an action potential, which activates synaptic transmission in the dorsal horn of spinal cord. Such processes induce a characteristic set of responses, including hyperalgesia (an increased response to a stimulus that normally provoke pain) and allodynia (pain due to a stimulus that does not normally provoke pain) [1].

Free radicals in biological systems are natural products during interactions between cells, as well as between tissues and organs. However, an imbalance between endogenous oxidants and antioxidants can alter cell homeostasis. Oxidants target proteins, thereby inducing cell damage and contributing to cell death [35]. Free radicals also cause inflammation [36]. Consequently, treatment with antioxidants is a promising approach to control a wide array of pathophysiological events. KO_2 is a O_2^- donor, and its injection resulting in nociceptive responses, such as




Fig 4. Naringenin inhibits KO_2 -induced oxidative stress and gp91^{phox} mRNA expression. (a-e) Mice received naringenin (50 mg/kg, po) treatment 1 h before ipl injection of 30 μ g of KO_2 . Paw skin samples were collected 3 h after ipl KO_2 injection. Sample analyses were (a) GSH levels, (b) total antioxidant capacity (FRAP assay), (c) lipid peroxidation (TBARS assay), (d) O_2^- production (NBT assay), and (e) gp91^{phox} mRNA expression by RT-qPCR. β -actin was a reference gene to normalize mRNA expression data. Results are mean \pm SEM of 6 mice per group per experiment, and are representative of 2 independent experiments. * p < 0.05 vs. saline group, # p < 0.05 vs. KO_2 group. One-way ANOVA followed Tukey's post hoc.

doi:10.1371/journal.pone.0153015.g004




Fig 5. Naringenin inhibits KO_2 -induced cytokine production and mRNA expression. (a-c) Mice received naringenin (50 mg/kg, po) treatment 1 h before ipl injection of 30 μ g of KO_2 . Paw skin samples were collected 3 h after ipl KO_2 injection. Sample analyses were (a) TNF α , and (b) IL-10 production by ELISA, and (c) IL-33 mRNA expression by RT-qPCR. β -actin was a reference gene to normalize mRNA expression data. Results are mean \pm SEM of 6 mice per group per experiment, and are representative of 2 independent experiments. * p < 0.05 vs. saline group, # p < 0.05 vs. KO_2 group. One-way ANOVA followed Tukey's post hoc.

doi:10.1371/journal.pone.0153015.g005




Fig 6. Naringenin inhibits KO_2 -induced COX-2 mRNA expression. Mice received naringenin (50 mg/kg, po) treatment 1 h before ipl injection of 30 μ g KO_2 . Paw skin samples were collected 3 h after ipl KO_2 injection and analyzed for COX-2 mRNA expression by RT-qPCR. β -actin was a reference gene to normalize mRNA expression data. Results are mean \pm SEM of 6 mice per group per experiment, and are representative of 2 independent experiments. * p < 0.05 vs. saline group, # p < 0.05 vs. KO_2 group. One-way ANOVA followed Tukey's post hoc.

doi:10.1371/journal.pone.0153015.g006

overt pain-like behavior, as well as mechanical and thermal hyperalgesia, which can be inhibited by analgesics [6]. Data indicates that O_2^- -induced pain depends on both direct superoxide anion effects, and indirect effects, partly *via* the production of peroxynitrite [37]. O_2^- injection triggers inflammatory pain by mechanisms involving cytokine, COX-2 and ET-1 synthesis [6,38,39]. Oxidative stress is involved in the pain of varied inflammatory diseases such as rheumatoid arthritis [40,41], gout [42], delayed onset muscle soreness [43] and diabetes [44].




Fig 7. Naringenin inhibits KO_2 -induced preproET-1 mRNA expression. Mice received naringenin (50 mg/kg, po) treatment 1 h before ipl injection of 30 μ g of KO_2 . Paw skin samples were collected 3 h after ipl KO_2 injection for preproET-1 mRNA expression analysis by RT-qPCR. β -actin was a reference gene to normalize mRNA expression data. Results are mean \pm SEM of 6 mice per group per experiment, and are representative of 2 independent experiments. * p < 0.05 vs. saline group, # p < 0.05 vs. KO_2 group. One-way ANOVA followed Tukey's post hoc.

doi:10.1371/journal.pone.0153015.g007




Fig 8. Naringenin increases Nrf2 and HO-1 mRNA expression. (a-b) Mice received naringenin (50 mg/kg, po) treatment 1 h before ipl injection of 30 μ g KO₂. Paw skin samples were collected 3 h after ipl KO₂ injection for (a) Nrf2, and (b) HO-1 mRNA expression analysis by RT-qPCR. β -actin was a reference gene to normalize mRNA expression data. Results are mean \pm SEM of 6 mice per group per experiment, and are representative of 2 independent experiments. * p < 0.05 vs. saline group, # p < 0.05 vs. KO₂ group. One-way ANOVA followed Tukey's post hoc.

doi:10.1371/journal.pone.0153015.g008

Therefore, KO₂-induced inflammatory pain is a useful model to study the action of analgesics that target oxidative stress-dependent events [12].

The therapeutic properties of naringenin include antinociception [15–17], as well as antioxidant [22,23,44–48] and anti-inflammatory activities [18,19,21,22]. However, the analgesic effect and mechanisms of action of naringenin in a model of oxidative stress-triggered inflammatory pain remained to be determined. Herein, we provide the first evidence that naringenin inhibits KO₂-induced overt-pain like behavior, MPO activity, mechanical hyperalgesia and thermal hyperalgesia via reduction of cytokine production and oxidative stress.

The activation of the NO–cGMP–PKG–K_{ATP} channel signaling pathway leads to nociceptor hyperpolarization, thereby reducing nociceptor neuronal transmission [25–28]. The present results show that naringenin inhibits the KO₂-induced mechanical and thermal hyperalgesia, at least in part, by activating the NO–cGMP–PKG–K_{ATP} channel signaling pathway. Free radical damage to membrane components leads to cellular dysfunction [49], which, by exposing neoepitopes, will contribute to an autoimmune response, which is receiving increasing interest as to its role in longer-term nociception. GSH is an important redox cellular system, with efficacy partly driven by the direct interactions of sulfhydryl groups (-SH) with ROS, thereby promoting a direct detox reaction [50]. Naringenin was able to restore GSH levels, as well as inhibiting other indicators of oxidative stress, such as FRAP, TBARS, and NBT in the paw skin. The KO₂ solution releases O₂⁻ for up to 10 min [6], however, NBT reduction (O₂⁻ production) occurred in paw skin samples collected 3 h after KO₂ injection. Therefore, O₂⁻ injection induces further O₂⁻. This is corroborated by the results showing KO₂-induced the gp91^{phox} mRNA expression, which was inhibited by naringenin. The gp91^{phox} subunit of NADPH oxidase participates in the electron transfer to oxygen, thereby generating O₂⁻ [51]. O₂⁻-induced pain also depends on hyperalgesic cytokines, with TNFR1 deficiency reducing O₂⁻-induced pain and oxidative stress. TNF α -induced hyperalgesia depends on NADPH oxidase activation [38]. Not all cytokines are nociceptive, with IL-10 having anti-hyperalgesic effects [43]. IL-33 is a hyperalgesic cytokine that regulates the production of TNF α and IL-10 [52–55]. Naringenin inhibited O₂⁻-induced production of TNF α and IL-10, as well as IL-33 mRNA expression.

Cytokine-induced inflammatory hyperalgesia in turn, depends, at least in part, on COX-2-dependent production of prostanoids such as prostaglandin E₂ and prostacyclin [1]. O₂⁻ also induces COX-2 mRNA expression, with celecoxib (a selective COX-2 inhibitor) diminishing O₂⁻-induced pain [6]. Naringenin inhibited O₂⁻-induced COX-2 mRNA expression. There is also a close relationship between cytokines and ET-1. Endothelin receptor antagonists inhibit cytokine-induced hyperalgesia and cytokines induce preproET-1 mRNA expression and ET-1

production [56,57]. Bosentan, an endothelin receptor antagonist, inhibits O_2^- -induced hyperalgesia, and O_2^- induces preproET-1 mRNA expression [39]. Naringenin inhibited O_2^- -induced prepro-ET-1 mRNA expression. Furthermore, activation of the ET receptors promotes oxidative stress and reduces the free radical scavenging ability [58]. Therefore, it is likely that ET-1 also contributes to the regulation of oxidative stress. In fact, bosentan inhibited O_2^- -induced oxidative stress [39].

Consistent with the naringenin inhibition of cytokine production as well as COX-2 and preproET-1 mRNA expression, naringenin inhibits the activation of the pro-inflammatory transcription factor, NF κ B, in several models of inflammation. For instance, naringenin inhibits NF κ B activation in dextran sulphate sodium-induced colitis [59], ethanol-induced liver injury [18], streptozotocin-induced diabetes in mice [60] and rats [61], and experimental stroke [23]. Furthermore, naringenin also inhibits NF κ B DNA-binding activity in ovalbumin-induced asthma [62] as well as in *in vitro* studies [63,64]. As such, it is not unlikely that naringenin inhibits KO_2 -induced cytokines production, as well as COX-2 and preproET-1, by inhibiting NF κ B induction and activity. Nrf2 is an important transcriptional regulator of the antioxidant response. In fact, Nrf2 activation is essential to the production of endogenous antioxidants such as GSH, thioredoxin system, HO-1, and NQO1 (NAD(P)H dehydrogenase, quinone 1) [65]. Keap-1 detects cellular environment changes such as oxidative stress, resulting in the activation of Nrf2, which readily translocates to the nucleus and upregulates downstream targets such as HO-1 [66]. Furthermore, the activation of the Nrf2/HO-1 pathway inhibits the production of inflammatory molecules such as TNF α [67–69], IL-6 [68,69] and IL-1 β [67,68]. Nrf2 indirectly modulates NF κ B activity [70], given that Nrf2 and NF κ B compete for binding to the nuclear complex, coactivator p300/CBP (E1A binding protein p300/CREB-binding protein) [71]. Importantly, naringenin activates Nrf2 and is an agonist of the aryl hydrocarbon receptor, contributing to reducing the production of reactive oxygen species and inflammatory mediators, as well as modulating specific immune cell activity [47,72]. In the present study, naringenin inhibited the KO_2 -induced decrease in Nrf2 mRNA expression and increased HO-1 mRNA expression. This mechanism might account for the analgesic effect of naringenin in KO_2 -induced inflammation, by inhibiting pro-hyperalgesic cytokine production, and gp91^{phox} as well as inducing antioxidant molecules.

The data presented here suggest that naringenin inhibits O_2^- -induced inflammatory overt pain-like behavior, hyperalgesia, and neutrophil recruitment by activating the NO–cGMP–PKG–K_{ATP} channel signaling pathway, as well as inhibiting oxidative stress and cytokine production, coupled to decreasing the mRNA of gp91^{phox}, IL-33, COX-2, and preproET-1, and increasing Nrf2 and HO-1 mRNA expression. These data show for the first time that naringenin inhibits inflammatory pain triggered by O_2^- and its mechanisms. Therefore, suggesting that naringenin is a promising therapeutic approach as an analgesic, antioxidant and anti-inflammatory compound, requiring further investigation in nociception, as well as in the array of other medical conditions where these pathophysiological changes are also evident [73].

Acknowledgments

Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Ministério da Ciências Tecnologia e Inovação (MCTI), Secretaria da Ciência, Tecnologia e Ensino Superior (SETI), Fundação Araucária and Parana State Government grants supported this study (Brazil). A.C.Z. received a postdoctoral fellowship from Fundação Araucária/CAPES (Brazil).

Author Contributions

Conceived and designed the experiments: RC WAVJ. Performed the experiments: MFM CCC LCS ACZ FAPR. Analyzed the data: MFM CCC LCS ACZ FAPR SRG MMB RC WAVJ. Contributed reagents/materials/analysis tools: SRG MMB RC WAVJ. Wrote the paper: MFM CCC LCS ACZ FAPR SRG MMB RC WAVJ. Drafted the article or revised it critically for important intellectual content: MFM CCC LCS ACZ FAPR SRG MMB RC WAVJ. Final approval of the version to be published: MFM CCC LCS ACZ FAPR SRG MMB RC WAVJ. Agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved: MFM CCC LCS ACZ FAPR SRG MMB RC WAVJ.

References

1. Verri WA, Cunha TM, Parada CA, Poole S, Cunha FQ, Ferreira SH. Hypernociceptive role of cytokines and chemokines: Targets for analgesic drug development? *Pharmacol Ther.* 2006; 112: 116–138. doi: [10.1016/j.pharmthera.2006.04.001](https://doi.org/10.1016/j.pharmthera.2006.04.001) PMID: [16730375](https://pubmed.ncbi.nlm.nih.gov/16730375/)
2. Keeble JE, Bodkin JV, Liang L, Wodarski R, Davies M, Fernandes ES, et al. Hydrogen peroxide is a novel mediator of inflammatory hyperalgesia, acting via transient receptor potential vanilloid 1-dependent and independent mechanisms. *Pain.* International Association for the Study of Pain; 2009; 141: 135–142. doi: [10.1016/j.pain.2008.10.025](https://doi.org/10.1016/j.pain.2008.10.025)
3. Ndengele MM, Cuzzocrea S, Esposito E, Mazzon E, Di Paola R, Matuschak GM, et al. Cyclooxygenases 1 and 2 contribute to peroxynitrite-mediated inflammatory pain hypersensitivity. *FASEB J.* 2008; 22: 3154–3164. doi: [10.1096/fj.08-108159](https://doi.org/10.1096/fj.08-108159) PMID: [18497304](https://pubmed.ncbi.nlm.nih.gov/18497304/)
4. Salvemini D, Little JW, Doyle T, Neumann WL. Roles of reactive oxygen and nitrogen species in pain. *Free Radic Biol Med.* 2011; 51: 951–966. doi: [10.1016/j.freeradbiomed.2011.01.026](https://doi.org/10.1016/j.freeradbiomed.2011.01.026) PMID: [21277369](https://pubmed.ncbi.nlm.nih.gov/21277369/)
5. Wang Z-Q. A Newly Identified Role for Superoxide in Inflammatory Pain. *J Pharmacol Exp Ther.* 2004; 309: 869–878. doi: [10.1124/jpet.103.064154](https://doi.org/10.1124/jpet.103.064154) PMID: [14988418](https://pubmed.ncbi.nlm.nih.gov/14988418/)
6. Maioli NA, Zarpelon AC, Mizokami SS, Calixto-Campos C, Guazelli CFS, Hohmann MSN, et al. The superoxide anion donor, potassium superoxide, induces pain and inflammation in mice through production of reactive oxygen species and cyclooxygenase-2. *Brazilian J Med Biol Res.* 2015; 48: 321–31. doi: [10.1590/1414-431X20144187](https://doi.org/10.1590/1414-431X20144187)
7. Droy-Lefaix MT, Drouet Y, Geraud G, Hosford D, Braquet P. Superoxide dismutase (SOD) and the PAF-antagonist (BN 52021) reduce small intestinal damage induced by ischemia-reperfusion. *Free Radic Res Commun.* 1991; 12–13 Pt 2: 725–35. PMID: [2060844](https://pubmed.ncbi.nlm.nih.gov/2060844/)
8. Matata BM, Galiñanes M. Peroxynitrite is an essential component of cytokines production mechanism in human monocytes through modulation of nuclear factor-kappa B DNA binding activity. *J Biol Chem.* 2002; 277: 2330–5. doi: [10.1074/jbc.M106393200](https://doi.org/10.1074/jbc.M106393200) PMID: [11706022](https://pubmed.ncbi.nlm.nih.gov/11706022/)
9. Salvemini D, Wang ZQ, Zweier JL, Samouilov A, Macarthur H, Misko TP, et al. A nonpeptidyl mimic of superoxide dismutase with therapeutic activity in rats. *Science.* 1999; 286: 304–6. PMID: [10514375](https://pubmed.ncbi.nlm.nih.gov/10514375/)
10. Boughton-Smith NK, Evans SM, Laszlo F, Whittle BJ, Moncada S. The induction of nitric oxide synthase and intestinal vascular permeability by endotoxin in the rat. *Br J Pharmacol.* 1993; 110: 1189–95 PMID: [7507778](https://pubmed.ncbi.nlm.nih.gov/7507778/)
11. Navarro SA, Serafim KGG, Mizokami SS, Hohmann MSN, Casagrande R, Verri WA. Analgesic activity of piracetam: Effect on cytokine production and oxidative stress. *Pharmacol Biochem Behav.* 2013; 105: 183–192. doi: [10.1016/j.pbb.2013.02.018](https://doi.org/10.1016/j.pbb.2013.02.018) PMID: [23474372](https://pubmed.ncbi.nlm.nih.gov/23474372/)
12. Fattori V, Pinho-Ribeiro FA, Borghi SM, Alves-Filho JC, Cunha TM, Cunha FQ, et al. Curcumin inhibits superoxide anion-induced pain-like behavior and leukocyte recruitment by increasing Nrf2 expression and reducing NF-κB activation. *Inflamm Res.* 2015; 64: 993–1003. doi: [10.1007/s00011-015-0885-y](https://doi.org/10.1007/s00011-015-0885-y) PMID: [26456836](https://pubmed.ncbi.nlm.nih.gov/26456836/)
13. Calixto-Campos C, Carvalho TT, Hohmann MSN, Pinho-Ribeiro FA, Fattori V, Manchope MF, et al. Vanillic Acid Inhibits Inflammatory Pain by Inhibiting Neutrophil Recruitment, Oxidative Stress, Cytokine Production, and NFκB Activation in Mice. *J Nat Prod.* 2015; 78: 1799–808. doi: [10.1021/acs.jnatprod.5b00246](https://doi.org/10.1021/acs.jnatprod.5b00246) PMID: [26192250](https://pubmed.ncbi.nlm.nih.gov/26192250/)
14. Lee C, Jeong T, Choi Y, Hyun B, Oh G. Anti-Atherogenic Effect of Citrus Flavonoids, Naringin and Naringenin, Associated with Hepatic ACAT and Aortic VCAM-1 and MCP-1 in High Cholesterol-Fed Rabbits. 2001; 688: 681–688. doi: [10.1006/bbrc.2001.5001](https://doi.org/10.1006/bbrc.2001.5001)

15. Anuja GI, Latha PG, Shine VJ, Suja SR, Shikha P, Kumar KS, et al. Antioedematous and Analgesic Properties of Fertile Fronds of *Drynaria quercifolia*. Hindawi Publishing Corporation; 2014; 2014: 2–8. doi: [10.1155/2014/302089](https://doi.org/10.1155/2014/302089)
16. Kaulaskar S, Bhutada P, Rahigude A, Jain D, Harle U. Effects of naringenin on allodynia and hyperalgesia in rats with chronic constriction injury-induced neuropathic pain. *Journal of Chinese integrative medicine*. 2012. pp. 1482–9. doi: [10.3736/jcim20121223](https://doi.org/10.3736/jcim20121223) PMID: [23257145](https://pubmed.ncbi.nlm.nih.gov/23257145/)
17. Hasanein P, Fazeli F. Role of naringenin in protection against diabetic hyperalgesia and tactile allodynia in male Wistar rats. *J Physiol Biochem*. 2014; 70: 997–1006. doi: [10.1007/s13105-014-0369-5](https://doi.org/10.1007/s13105-014-0369-5) PMID: [25407136](https://pubmed.ncbi.nlm.nih.gov/25407136/)
18. Jayaraman J, Jesudoss VA, Menon VP, Namasivayam N. Anti-inflammatory role of naringenin in rats with ethanol induced liver injury. *Toxicol Mech Methods*. 2012; 22: 568–576. doi: [10.3109/15376516.2012.707255](https://doi.org/10.3109/15376516.2012.707255) PMID: [22900548](https://pubmed.ncbi.nlm.nih.gov/22900548/)
19. Chao C-L, Weng C-S, Chang N-C, Lin J-S, Kao S-T, Ho F-M. Naringenin more effectively inhibits inducible nitric oxide synthase and cyclooxygenase-2 expression in macrophages than in microglia. *Nutr Res*. 2010; 30: 858–64. doi: [10.1016/j.nutres.2010.10.011](https://doi.org/10.1016/j.nutres.2010.10.011) PMID: [21147369](https://pubmed.ncbi.nlm.nih.gov/21147369/)
20. Karuppagounder V, Arumugam S, Thandavarayan RA, Pitchaimani V, Sreedhar R, Afrin R, et al. Naringenin ameliorates daunorubicin induced nephrotoxicity by mitigating AT1R, ERK1/2-NFκB p65 mediated inflammation. *Int Immunopharmacol*. 2015; 28: 154–159. doi: [10.1016/j.intimp.2015.05.050](https://doi.org/10.1016/j.intimp.2015.05.050) PMID: [26072060](https://pubmed.ncbi.nlm.nih.gov/26072060/)
21. Yang J, Li Q, Zhou XD, Kolosov VP, Perelman JM. Naringenin attenuates mucous hypersecretion by modulating reactive oxygen species production and inhibiting NF-κB activity via EGFR-PI3K-Akt/ERK MAPKinase signaling in human airway epithelial cells. *Mol Cell Biochem*. 2011; 351: 29–40. doi: [10.1007/s11010-010-0708-y](https://doi.org/10.1007/s11010-010-0708-y) PMID: [21229383](https://pubmed.ncbi.nlm.nih.gov/21229383/)
22. Martinez RM, Pinho-Ribeiro FA, Steffen VS, Caviglione CV, Vignoli JA, Barbosa DS, et al. Naringenin Inhibits UVB Irradiation-Induced Inflammation and Oxidative Stress in the Skin of Hairless Mice. *J Nat Prod*. 2015; 78: 1647–1655. doi: [10.1021/acs.jnatprod.5b00198](https://doi.org/10.1021/acs.jnatprod.5b00198) PMID: [26154512](https://pubmed.ncbi.nlm.nih.gov/26154512/)
23. Raza SS, Khan MM, Ahmad A, Ashafaq M, Islam F, Wagner A. P, et al. Neuroprotective effect of naringenin is mediated through suppression of NF-κB signaling pathway in experimental stroke. *Neuroscience*. 2013; 230: 157–171. doi: [10.1016/j.neuroscience.2012.10.041](https://doi.org/10.1016/j.neuroscience.2012.10.041) PMID: [23103795](https://pubmed.ncbi.nlm.nih.gov/23103795/)
24. Esmaeili MA, Alilou M. Naringenin attenuates CCl4-induced hepatic inflammation by the activation of an Nrf2-mediated pathway in rats. *Clin Exp Pharmacol Physiol*. 2014; 41: 416–22. doi: [10.1111/1440-1681.12230](https://doi.org/10.1111/1440-1681.12230) PMID: [24684352](https://pubmed.ncbi.nlm.nih.gov/24684352/)
25. Cunha TM, Roman-Campos D, Lotufo CM, Duarte HL, Souza GR, Verri WA, et al. Morphine peripheral analgesia depends on activation of the PI3Kγ/AKT/nNOS/NO/KATP signaling pathway. *Proc Natl Acad Sci U S A*. 2010; 107: 4442–7. doi: [10.1073/pnas.0914733107](https://doi.org/10.1073/pnas.0914733107) PMID: [20147620](https://pubmed.ncbi.nlm.nih.gov/20147620/)
26. Duarte ID, dos Santos IR, Lorenzetti BB, Ferreira SH. Analgesia by direct antagonism of nociceptor sensitization involves the arginine-nitric oxide-cGMP pathway. *Eur J Pharmacol*. 1992; 217: 225–227. doi: [10.1016/0014-2999\(92\)90881-4](https://doi.org/10.1016/0014-2999(92)90881-4) PMID: [1330593](https://pubmed.ncbi.nlm.nih.gov/1330593/)
27. Sachs D, Cunha FQ, Ferreira SH. Peripheral analgesic blockade of hypernociception: activation of arginine/NO/cGMP/protein kinase G/ATP-sensitive K⁺ channel pathway. *Proc Natl Acad Sci U S A*. 2004; 101: 3680–5. doi: [10.1073/pnas.0308382101](https://doi.org/10.1073/pnas.0308382101) PMID: [14990791](https://pubmed.ncbi.nlm.nih.gov/14990791/)
28. Ventura-Martínez R, Déciga-Campos M, Díaz-Reval MI, González-Trujano ME, López-Muñoz FJ. Peripheral involvement of the nitric oxide-cGMP pathway in the indomethacin-induced antinociception in rat. *Eur J Pharmacol*. 2004; 503: 43–8. doi: [10.1016/j.ejphar.2004.09.018](https://doi.org/10.1016/j.ejphar.2004.09.018) PMID: [15496294](https://pubmed.ncbi.nlm.nih.gov/15496294/)
29. Cunha TM, Verri WA, Vivancos GG, Moreira IF, Reis S, Parada CA, et al. An electronic pressure-meter nociception paw test for mice. *Braz J Med Biol Res*. 2004; 37: 401–7. PMID: [15060710](https://pubmed.ncbi.nlm.nih.gov/15060710/)
30. Lavich TR, Cordeiro RSB, Silva PMR, Martins MA. A novel hot-plate test sensitive to hyperalgesic stimuli and non-opioid analgesics. *Brazilian J Med Biol Res*. 2005; 38: 445–51. /S0100-879X2005000300016
31. Sedlak J, Lindsay RH. Estimation of total, protein-bound, and nonprotein sulfhydryl groups in tissue with Ellman's reagent. *Anal Biochem*. 1968; 25: 192–205. PMID: [4973948](https://pubmed.ncbi.nlm.nih.gov/4973948/)
32. Katalinic V, Modun D, Music I, Boban M. Gender differences in antioxidant capacity of rat tissues determined by 2,2'-azinobis (3-ethylbenzothiazoline 6-sulfonate; ABTS) and ferric reducing antioxidant power (FRAP) assays. *Comp Biochem Physiol C Toxicol Pharmacol*. 2005; 140: 47–52. doi: [10.1016/j.cca.2005.01.005](https://doi.org/10.1016/j.cca.2005.01.005) PMID: [15792622](https://pubmed.ncbi.nlm.nih.gov/15792622/)
33. Hohmann MSN, Cardoso RDR, Pinho-Ribeiro FA, Crespigio J, Cunha TM, Alves-Filho JC, et al. 5-lipoxygenase deficiency reduces acetaminophen-induced hepatotoxicity and lethality. *Biomed Res Int*. 2013; 2013: 627046. doi: [10.1155/2013/627046](https://doi.org/10.1155/2013/627046) PMID: [24288682](https://pubmed.ncbi.nlm.nih.gov/24288682/)

34. Guedes RP, Bosco LD, Teixeira CM, Araújo ASR, Llesuy S, Belló-Klein A, et al. Neuropathic pain modifies antioxidant activity in rat spinal cord. *Neurochem Res.* 2006; 31: 603–9. doi: [10.1007/s11064-006-9058-2](https://doi.org/10.1007/s11064-006-9058-2) PMID: [16770731](https://pubmed.ncbi.nlm.nih.gov/16770731/)
35. Dean RT, Fu S, Stocker R, Davies MJ. Biochemistry and pathology of radical-mediated protein oxidation. *Biochem J.* 1997; 324
36. Closa D, Folch-Puy E. Oxygen free radicals and the systemic inflammatory response. *IUBMB Life.* 2004; 56: 185–91. doi: [10.1080/15216540410001701642](https://doi.org/10.1080/15216540410001701642) PMID: [15230345](https://pubmed.ncbi.nlm.nih.gov/15230345/)
37. Khattab MM. TEMPOL, a membrane-permeable radical scavenger, attenuates peroxynitrite- and superoxide anion-enhanced carrageenan-induced paw edema and hyperalgesia: a key role for superoxide anion. *Eur J Pharmacol.* 2006; 548: 167–73. doi: [10.1016/j.ejphar.2006.08.007](https://doi.org/10.1016/j.ejphar.2006.08.007) PMID: [16973155](https://pubmed.ncbi.nlm.nih.gov/16973155/)
38. Yamacita-Borin FY, Zarpelon AC, Pinho-Ribeiro FA, Fattori V, Alves-Filho JC, Cunha FQ, et al. Superoxide anion-induced pain and inflammation depends on TNF α /TNFR1 signaling in mice. *Neurosci Lett.* 2015; 605: 53–58. doi: [10.1016/j.neulet.2015.08.015](https://doi.org/10.1016/j.neulet.2015.08.015) PMID: [26291484](https://pubmed.ncbi.nlm.nih.gov/26291484/)
39. Serafim KGG, Navarro SA, Zarpelon AC, Pinho-Ribeiro FA, Fattori V, Cunha TM, et al. Bosentan, a mixed endothelin receptor antagonist, inhibits superoxide anion-induced pain and inflammation in mice. *Naunyn Schmiedeberg Arch Pharmacol.* 2015; 388: 1211–1221. doi: [10.1007/s00210-015-1160-z](https://doi.org/10.1007/s00210-015-1160-z) PMID: [26246053](https://pubmed.ncbi.nlm.nih.gov/26246053/)
40. Moon S-J, Woo Y-J, Jeong J-H, Park M-K, Oh H-J, Park J-S, et al. Rebamipide attenuates pain severity and cartilage degeneration in a rat model of osteoarthritis by downregulating oxidative damage and catabolic activity in chondrocytes. *Osteoarthritis Cartilage.* 2012; 20: 1426–38. doi: [10.1016/j.joca.2012.08.002](https://doi.org/10.1016/j.joca.2012.08.002) PMID: [22890185](https://pubmed.ncbi.nlm.nih.gov/22890185/)
41. Fattori V, Amaral FA, Verri WA. Neutrophils and arthritis: Role in disease and pharmacological perspectives. *Pharmacol Res.* 2016; doi: [10.1016/j.phrs.2016.01.027](https://doi.org/10.1016/j.phrs.2016.01.027)
42. Zamudio-Cuevas Y, Hernández-Díaz C, Pineda C, Reginato AM, Cerna-Cortés JF, Ventura-Ríos L, et al. Molecular basis of oxidative stress in gouty arthropathy. *Clin Rheumatol.* 2015; 34: 1667–72. doi: [10.1007/s10067-015-2933-y](https://doi.org/10.1007/s10067-015-2933-y) PMID: [25854697](https://pubmed.ncbi.nlm.nih.gov/25854697/)
43. Borghi SM, Pinho-Ribeiro FA, Zarpelon AC, Cunha TM, Alves-Filho JC, Ferreira SH, et al. Interleukin-10 limits intense acute swimming-induced muscle mechanical hyperalgesia in mice. *Exp Physiol.* 2015; 100: 531–544. doi: [10.1113/EP085026](https://doi.org/10.1113/EP085026) PMID: [25711612](https://pubmed.ncbi.nlm.nih.gov/25711612/)
44. Al-Rejaie SS, Aleisa AM, Abuhashish HM, Parmar MY, Ola MS, Al-Hosaini AA, et al. Naringenin neutralises oxidative stress and nerve growth factor discrepancy in experimental diabetic neuropathy. *Neurol Res.* 2015; 37: 924–933. doi: [10.1179/1743132815Y.0000000079](https://doi.org/10.1179/1743132815Y.0000000079) PMID: [26187552](https://pubmed.ncbi.nlm.nih.gov/26187552/)
45. Chen S, Ding Y, Tao W, Zhang W, Liang T, Liu C. Naringenin inhibits TNF- α induced VSMC proliferation and migration via induction of HO-1. *Food Chem Toxicol.* 2012; 50: 3025–31. doi: [10.1016/j.fct.2012.06.006](https://doi.org/10.1016/j.fct.2012.06.006) PMID: [22709785](https://pubmed.ncbi.nlm.nih.gov/22709785/)
46. Cavia-Saiz M, Busto MD, Pilar-Izquierdo MC, Ortega N, Perez-Mateos M, Muñoz P. Antioxidant properties, radical scavenging activity and biomolecule protection capacity of flavonoid naringenin and its glycoside naringin: A comparative study. *J Sci Food Agric.* 2010; 90: 1238–1244. doi: [10.1002/jsfa.3959](https://doi.org/10.1002/jsfa.3959) PMID: [20394007](https://pubmed.ncbi.nlm.nih.gov/20394007/)
47. Lou H, Jing X, Wei X, Shi H, Ren D, Zhang X. Naringenin protects against 6-OHDA-induced neurotoxicity via activation of the Nrf2/ARE signaling pathway. *Neuropharmacology.* 2014; 79: 380–388. doi: [10.1016/j.neuropharm.2013.11.026](https://doi.org/10.1016/j.neuropharm.2013.11.026) PMID: [24333330](https://pubmed.ncbi.nlm.nih.gov/24333330/)
48. Ramprasath T, Senthamizharasi M, Vasudevan V, Sasikumar S, Yuvaraj S, Selvam GS. Naringenin confers protection against oxidative stress through upregulation of Nrf2 target genes in cardiomyoblast cells. *J Physiol Biochem.* 2014; 70: 407–15. doi: [10.1007/s13105-014-0318-3](https://doi.org/10.1007/s13105-014-0318-3) PMID: [24526395](https://pubmed.ncbi.nlm.nih.gov/24526395/)
49. Chtourou Y, Aouey B, Kebieche M, Fetoui H. Protective role of naringin against cisplatin induced oxidative stress, inflammatory response and apoptosis in rat striatum via suppressing ROS-mediated NF- κ B and P53 signaling pathways. *Chem Biol Interact.* 2015; 239: 76–86. doi: [10.1016/j.cbi.2015.06.036](https://doi.org/10.1016/j.cbi.2015.06.036) PMID: [26120027](https://pubmed.ncbi.nlm.nih.gov/26120027/)
50. Lushchak VI. Glutathione homeostasis and functions: potential targets for medical interventions. *J Amino Acids.* 2012; 2012: 736837. doi: [10.1155/2012/736837](https://doi.org/10.1155/2012/736837) PMID: [22500213](https://pubmed.ncbi.nlm.nih.gov/22500213/)
51. Yu L, Quinn MT, Cross AR, Dinauer MC. Gp91phox is the heme binding subunit of the superoxide-generating NADPH oxidase. *Proc Natl Acad Sci.* 1998; 95: 7993–7998. doi: [10.1073/pnas.95.14.7993](https://doi.org/10.1073/pnas.95.14.7993) PMID: [9653128](https://pubmed.ncbi.nlm.nih.gov/9653128/)
52. Verri WA, Guerrero ATG, Fukada SY, Valerio DA, Cunha TM, Xu D, et al. IL-33 mediates antigen-induced cutaneous and articular hypernociception in mice. *Proc Natl Acad Sci U S A.* 2008; 105: 2723–8. doi: [10.1073/pnas.0712116105](https://doi.org/10.1073/pnas.0712116105) PMID: [18250323](https://pubmed.ncbi.nlm.nih.gov/18250323/)

53. Magro DAC, Hohmann MSN, Mizokami SS, Cunha TM, Alves-Filho JC, Casagrande R, et al. An interleukin-33/ST2 signaling deficiency reduces overt pain-like behaviors in mice. *Braz J Med Biol Res.* 2013; 46: 601–6. doi: [10.1590/1414-431X20132894](https://doi.org/10.1590/1414-431X20132894) PMID: [23903682](https://pubmed.ncbi.nlm.nih.gov/23903682/)
54. Zarpelon AC, Cunha TM, Alves-Filho JC, Pinto LG, Ferreira SH, McInnes IB, et al. IL-33/ST2 signalling contributes to carrageenin-induced innate inflammation and inflammatory pain: role of cytokines, endothelin-1 and prostaglandin E2. *Br J Pharmacol.* 2013; 169: 90–101. doi: [10.1111/bph.12110](https://doi.org/10.1111/bph.12110) PMID: [23347081](https://pubmed.ncbi.nlm.nih.gov/23347081/)
55. Zarpelon AC, Rodrigues FC, Lopes AH, Souza GR, Carvalho TT, Pinto LG, et al. Spinal cord oligodendrocyte-derived alarmin IL-33 mediates neuropathic pain. *FASEB J.* 2016; 30: 54–65. doi: [10.1096/fj.14-267146](https://doi.org/10.1096/fj.14-267146) PMID: [26310268](https://pubmed.ncbi.nlm.nih.gov/26310268/)
56. Verri WA, Cunha TM, Parada CA, Wei X, Ferreira SH, Liew FY, et al. IL-15 mediates immune inflammatory hypernociception by triggering a sequential release of IFN-gamma, endothelin, and prostaglandin. *Proc Natl Acad Sci U S A.* 2006; 103: 9721–5. doi: [10.1073/pnas.0603286103](https://doi.org/10.1073/pnas.0603286103) PMID: [16766656](https://pubmed.ncbi.nlm.nih.gov/16766656/)
57. Verri WA, Cunha TM, Parada CA, Poole S, Liew FY, Ferreira SH, et al. Antigen-induced inflammatory mechanical hypernociception in mice is mediated by IL-18. *Brain Behav Immun.* 2007; 21: 535–43. doi: [10.1016/j.bbi.2006.11.005](https://doi.org/10.1016/j.bbi.2006.11.005) PMID: [17194566](https://pubmed.ncbi.nlm.nih.gov/17194566/)
58. Goto T, Hussein MH, Kato S, Daoud GA-H, Kato T, Sugiura T, et al. Endothelin receptor antagonist attenuates oxidative stress in a neonatal sepsis piglet model. *Pediatr Res.* 2012; 72: 600–5. doi: [10.1038/pr.2012.134](https://doi.org/10.1038/pr.2012.134) PMID: [23041664](https://pubmed.ncbi.nlm.nih.gov/23041664/)
59. Dou W, Zhang J, Sun A, Zhang E, Ding L, Mukherjee S, et al. Protective effect of naringenin against experimental colitis via suppression of Toll-like receptor 4/NF- κ B signalling. *Br J Nutr.* 2013; 110: 599–608. doi: [10.1017/S0007114512005594](https://doi.org/10.1017/S0007114512005594) PMID: [23506745](https://pubmed.ncbi.nlm.nih.gov/23506745/)
60. Tsai SJ, Huang CS, Mong MC, Kam WY, Huang HY, Yin MC. Anti-inflammatory and antifibrotic effects of naringenin in diabetic mice. *J Agric Food Chem.* 2012; 60: 514–521. doi: [10.1021/jf203259h](https://doi.org/10.1021/jf203259h) PMID: [22117528](https://pubmed.ncbi.nlm.nih.gov/22117528/)
61. Chen F, Zhang N, Ma X, Huang T, Shao Y, Wu C, et al. Naringin Alleviates Diabetic Kidney Disease through Inhibiting Oxidative Stress and Inflammatory Reaction. *PLoS One.* 2015; 10: e0143868. doi: [10.1371/journal.pone.0143868](https://doi.org/10.1371/journal.pone.0143868) PMID: [26619044](https://pubmed.ncbi.nlm.nih.gov/26619044/)
62. Shi Y, Dai J, Liu H, Li R-R, Sun P-L, Du Q, et al. Naringenin inhibits allergen-induced airway inflammation and airway responsiveness and inhibits NF- κ B activity in a murine model of asthma. *Can J Physiol Pharmacol.* 2009; 87: 729–35. doi: [10.1139/y09-065](https://doi.org/10.1139/y09-065) PMID: [19794524](https://pubmed.ncbi.nlm.nih.gov/19794524/)
63. Lim R, Barker G, Wall CA, Lappas M. Dietary phytophenols curcumin, naringenin and apigenin reduce infection-induced inflammatory and contractile pathways in human placenta, foetal membranes and myometrium. *Mol Hum Reprod.* 2013; 19: 451–62. doi: [10.1093/molehr/gat015](https://doi.org/10.1093/molehr/gat015) PMID: [23475986](https://pubmed.ncbi.nlm.nih.gov/23475986/)
64. Tsai SH, Lin-Shiau SY, Lin JK. Suppression of nitric oxide synthase and the down-regulation of the activation of NF κ B in macrophages by resveratrol. *Br J Pharmacol.* 1999; 126: 673–80. doi: [10.1038/sj.bjp.0702357](https://doi.org/10.1038/sj.bjp.0702357) PMID: [10188978](https://pubmed.ncbi.nlm.nih.gov/10188978/)
65. Rangasamy T, Cho CY, Thimmulappa RK, Zhen L, Srisuma SS, Kensler TW, et al. Genetic ablation of Nrf2 enhances susceptibility to cigarette smoke-induced emphysema in mice. *J Clin Invest.* 2004; 114: 1248–59. doi: [10.1172/JCI21146](https://doi.org/10.1172/JCI21146) PMID: [15520857](https://pubmed.ncbi.nlm.nih.gov/15520857/)
66. Li N, Alam J, Venkatesan MI, Eiguren-Fernandez A, Schmitz D, Di Stefano E, et al. Nrf2 Is a Key Transcription Factor That Regulates Antioxidant Defense in Macrophages and Epithelial Cells: Protecting against the Proinflammatory and Oxidizing Effects of Diesel Exhaust Chemicals. *J Immunol.* American Association of Immunologists; 2004; 173: 3467–3481. doi: [10.4049/jimmunol.173.5.3467](https://doi.org/10.4049/jimmunol.173.5.3467)
67. Yeligar SM, Machida K, Kalra VK. Ethanol-induced HO-1 and NQO1 are differentially regulated by HIF-1 α and Nrf2 to attenuate inflammatory cytokine expression. *J Biol Chem.* 2010; 285: 35359–73. doi: [10.1074/jbc.M110.138636](https://doi.org/10.1074/jbc.M110.138636) PMID: [20833713](https://pubmed.ncbi.nlm.nih.gov/20833713/)
68. So H, Kim H, Kim Y, Kim E, Pae H-O, Chung H-T, et al. Evidence that cisplatin-induced auditory damage is attenuated by downregulation of pro-inflammatory cytokines via Nrf2/HO-1. *J Assoc Res Otolaryngol.* 2008; 9: 290–306. doi: [10.1007/s10162-008-0126-y](https://doi.org/10.1007/s10162-008-0126-y) PMID: [18584244](https://pubmed.ncbi.nlm.nih.gov/18584244/)
69. Innamorato NG, Rojo AI, García-Yagüe AJ, Yamamoto M, de Ceballos ML, Cuadrado A. The transcription factor Nrf2 is a therapeutic target against brain inflammation. *J Immunol.* 2008; 181: 680–9. PMID: [18566435](https://pubmed.ncbi.nlm.nih.gov/18566435/)
70. Pan H, Wang H, Wang X, Zhu L, Mao L. The absence of Nrf2 enhances NF- κ B-dependent inflammation following scratch injury in mouse primary cultured astrocytes. *Mediators Inflamm.* 2012; 2012: 217580. doi: [10.1155/2012/217580](https://doi.org/10.1155/2012/217580) PMID: [22529521](https://pubmed.ncbi.nlm.nih.gov/22529521/)
71. Kim S-W, Lee H-K, Shin J-H, Lee J-K. Up-down regulation of HO-1 and iNOS gene expressions by ethyl pyruvate via recruiting p300 to Nrf2 and depriving It from p65. *Free Radic Biol Med.* 2013; 65: 468–76. doi: [10.1016/j.freeradbiomed.2013.07.028](https://doi.org/10.1016/j.freeradbiomed.2013.07.028) PMID: [23891677](https://pubmed.ncbi.nlm.nih.gov/23891677/)

72. Wang H-K, Yeh C-H, Iwamoto T, Satsu H, Shimizu M, Totsuka M. Dietary flavonoid naringenin induces regulatory T cells via an aryl hydrocarbon receptor mediated pathway. *J Agric Food Chem.* 2012; 60: 2171–8. doi: [10.1021/jf204625y](https://doi.org/10.1021/jf204625y) PMID: [22324845](https://pubmed.ncbi.nlm.nih.gov/22324845/)
73. Anderson G, Maes M. Oxidative/nitrosative stress and immuno-inflammatory pathways in depression: treatment implications. *Curr Pharm Des.* 2014; 20: 3812–47. doi: [10.2174/13816128113196660738](https://doi.org/10.2174/13816128113196660738) PMID: [24180395](https://pubmed.ncbi.nlm.nih.gov/24180395/)