CORRECTION

Correction: Detection of a Serum Siderophore by LC-MS/ MS as a Potential Biomarker of Invasive Aspergillosis

Cassandra S. Carroll, Lawrence N. Amankwa, Linda J. Pinto, Jeffrey D. Fuller, Margo M. Moore

In [Fig 3](#), the TAFC value for the SLE patient data is incorrect. The correct TAFC value is <5 ng/ml. Please see the corrected [Fig 3](#) here.

OPEN ACCESS

Citation: Carroll CS, Amankwa LN, Pinto LJ, Fuller JD, Moore MM (2016) Correction: Detection of a Serum Siderophore by LC-MS/ MS as a Potential Biomarker of Invasive Aspergillosis. PLoS ONE 11 (5): e0155451. doi:10.1371/journal.pone.0155451

Published: May 10, 2016

Copyright: © 2016 Carroll et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Fig 3. Representative TAFC chromatograms from various patient serum samples. Note the difference in scale of the Y-axis. All samples were concentrated 4-fold prior to TAFC measurement.

doi:10.1371/journal.pone.0155451.g001

Reference

1. Carroll CS, Amankwa LN, Pinto LJ, Fuller JD, Moore MM (2016) Detection of a Serum Siderophore by LC-MS/MS as a Potential Biomarker of Invasive Aspergillosis. PLoS ONE 11(3): e0151260. doi:[10.1371/journal.pone.0151260](https://doi.org/10.1371/journal.pone.0151260) PMID: [26974544](https://pubmed.ncbi.nlm.nih.gov/26974544/)