CORRECTION

Correction: Early Treatment Critical: Bexarotene Reduces Amyloid-Beta Burden In Silico

Joseph Rosenthal, Georges Belfort, David Isaacson

[S1 Text](#), [S1](#) and [S2](#) Tables are incorrect. Please view the corrected [S1 Text](#), [S1](#) and [S2](#) Tables here.

Supporting Information

S1 Text. Note on *APP/PS1* mice.
(PDF)

S1 Table. Solver run times and percent changes in final diseased neuron concentration and $A\beta_{42}$ load. Each percent change is given in absolute value when compared to results from a run solved with ode45 for a 90 day simulation of a nine month-old *APP/PS1* transgenic mouse with $\text{mg} \cdot \text{kg}^{-1}$ bexarotene treatment.
(PDF)

S2 Table. Percent change of the concentration of healthy brain cells in a 90 day simulation of nine-month-old *APP/PS1* mouse with $100 \text{ mg} \cdot \text{kg}^{-1}$ bexarotene treatment. Parameters are increased by 10%, and the approximate corresponding percent change of the system is given for $0 \text{ mg} \cdot \text{kg}^{-1}$ and $100 \text{ mg} \cdot \text{kg}^{-1}$ of bexarotene.
(PDF)

Reference

1. Rosenthal J, Belfort G, Isaacson D (2016) Early Treatment Critical: Bexarotene Reduces Amyloid-Beta Burden In Silico. PLoS ONE 11(4): e0153150. doi: [10.1371/journal.pone.0153150](https://doi.org/10.1371/journal.pone.0153150) PMID: [27073866](https://pubmed.ncbi.nlm.nih.gov/27073866/)

OPEN ACCESS

Citation: Rosenthal J, Belfort G, Isaacson D (2016) Correction: Early Treatment Critical: Bexarotene Reduces Amyloid-Beta Burden In Silico. PLoS ONE 11(5): e0156474. doi: [10.1371/journal.pone.0156474](https://doi.org/10.1371/journal.pone.0156474)

Published: May 24, 2016

Copyright: © 2016 Rosenthal et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.