CORRECTION

Correction: Neglecting the fallow season can significantly underestimate annual methane emissions in Mediterranean rice fields

Maite Martínez-Eixarch, Carles Alcaraz, Marc Viñas, Joan Noguerol, Xavier Aranda, Francesc Xavier Prenafeta-Boldú, Jesús Antonio Saldaña-De la Vega, Maria del Mar Català, Carles Ibáñez

Notice of Republication

An incorrect version of S1 Table was published in error. This article was republished on July 25, 2018 to correct for this error. Please download this article again to view the correct version.

Reference

1. Martínez-Eixarch M, Alcaraz C, Viñas M, Noguerol J, Aranda X, Prenafeta-Boldú FX, et al. (2018) Neglecting the fallow season can significantly underestimate annual methane emissions in Mediterranean rice fields. PLoS ONE 13(5): e0198081. <https://doi.org/10.1371/journal.pone.0198081> PMID: 29852016

OPEN ACCESS

Citation: Martínez-Eixarch M, Alcaraz C, Viñas M, Noguerol J, Aranda X, Prenafeta-Boldú FX, et al. (2018) Correction: Neglecting the fallow season can significantly underestimate annual methane emissions in Mediterranean rice fields. PLoS ONE 13(8): e0202159. <https://doi.org/10.1371/journal.pone.0202159>

Published: August 8, 2018

Copyright: © 2018 Martínez-Eixarch et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.