CORRECTION

Correction: Preferring more e-cigarette flavors is associated with e-cigarette use frequency among adolescents but not adults

Meghan E. Morean, Ellyn R. Butler, Krysten W. Bold, Grace Kong, Deepa R. Camenga, Dana A. Cavallo, Patricia Simon, Stephanie S. O'Malley, Suchitra Krishnan-Sarin

There are errors in the Funding statement. The correct funding information is as follows: Research reported in this publication was supported by the National Institutes of Health/ NIDA and FDA Center for Tobacco Products (CTP) (P50DA036151; Yale Tobacco Center of Regulatory Science, TCORS); the National Institute on Drug Abuse (T32DA019426); and the National Center on Addiction and Substance Abuse at Columbia University (to G.K.). The content is solely the responsibility of the authors and does not necessarily represent the official views of the funding agencies, including the NIH or the Food and Drug Administration. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Reference

1. Morean ME, Butler ER, Bold KW, Kong G, Camenga DR, Cavallo DA, et al. (2018) Preferring more e-cigarette flavors is associated with e-cigarette use frequency among adolescents but not adults. PLoS ONE 13(1): e0189015. <https://doi.org/10.1371/journal.pone.0189015> PMID: 29300749

OPEN ACCESS

Citation: Morean ME, Butler ER, Bold KW, Kong G, Camenga DR, Cavallo DA, et al. (2018) Correction: Preferring more e-cigarette flavors is associated with e-cigarette use frequency among adolescents but not adults. PLoS ONE 13(9): e0204349. <https://doi.org/10.1371/journal.pone.0204349>

Published: September 13, 2018

Copyright: © 2018 Morean et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.