

RESEARCH ARTICLE

# Human coronavirus alone or in co-infection with rhinovirus C is a risk factor for severe respiratory disease and admission to the pediatric intensive care unit: A one-year study in Southeast Brazil

Alessandra K. Matsuno<sup>1</sup>\*, Talita B. Gagliardi<sup>2</sup>, Flavia E. Paula<sup>2</sup>, Luciano K. S. Luna<sup>2</sup>, Bruna L. S. Jesus<sup>2</sup>, Renato T. Stein<sup>3</sup>, Davi C. Aragon<sup>1</sup>, Ana P. C. P. Carlotti<sup>1</sup>‡, Eurico Arruda<sup>2</sup>‡

**1** Department of Pediatrics, Ribeirão Preto Medical School, University of São Paulo, Ribeirão Preto, Brazil, **2** Department of Cell Biology and Virology Research Center, Ribeirão Preto Medical School, University of Sao Paulo, Ribeirão Preto, Brazil, **3** Department of Pediatrics, Pontifícia Universidade Católica do Rio Grande do Sul School of Medicine, Porto Alegre, Brazil

\* These authors contributed equally to this work.

‡ These authors also contributed equally to this work.

\* [matsuno@fmrp.usp.br](mailto:matsuno@fmrp.usp.br)



**OPEN ACCESS**

**Citation:** Matsuno AK, Gagliardi TB, Paula FE, Luna LKS, Jesus BLS, Stein RT, et al. (2019) Human coronavirus alone or in co-infection with rhinovirus C is a risk factor for severe respiratory disease and admission to the pediatric intensive care unit: A one-year study in Southeast Brazil. PLoS ONE 14 (6): e0217744. <https://doi.org/10.1371/journal.pone.0217744>

**Editor:** Raymond J. Pickles, University of North Carolina at Chapel Hill, UNITED STATES

**Received:** October 2, 2018

**Accepted:** May 19, 2019

**Published:** June 3, 2019

**Copyright:** © 2019 Matsuno et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the manuscript and its Supporting Information files.

**Funding:** AKM and EAN supported mainly by an academic grant from Abbott Laboratórios do Brasil Ltda based on an unrestricted investigator-generated proposal. EAN Additional funding was provided by a grant from the Sao Paulo State Research Foundation (FAPESP) and by FAEPA -

## Abstract

### Objective

We aimed to assess the profile of respiratory viruses in young children hospitalized for acute lower respiratory tract infection (ALRI) and its association with disease severity, defined as need for pediatric intensive care unit (PICU) admission.

### Design

Prospective observational cohort study.

### Setting

A tertiary-care university hospital in Brazil.

### Patients

Children younger than three years attending the pediatric emergency room with ALRI who were admitted to the hospital.

### Interventions

None.

“Fundação de Apoio ao Ensino Pesquisa e Assistência” (in English would be “Foundation of Support to Teaching Research and Assistance) of the University of Sao Paulo Hospital of Ribeirão Preto. EA has long standing scholarship support from the Brazilian Research Council (CNPq). The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** AKM and EAN were supported by an academic grant from Abbott Laboratórios do Brasil Ltda, based on an unrestricted investigator-generated proposal. This does not alter our adherence to PLOS ONE policies on sharing data and materials, as detailed online in the guide for authors: <https://journals.plos.org/plosone/s/competing-interests>.

## Measurements and main results

Nasopharyngeal aspirates were collected from patients from June 1<sup>st</sup>, 2008 to May 31<sup>st</sup>, 2009 within the first 48 hours of hospitalization. Nasopharyngeal aspirates were tested for 17 human respiratory viruses by molecular and immunofluorescence based assays. Simple and multiple log-binomial regression models were constructed to assess associations of virus type with a need for PICU admission. Age, prematurity, the presence of an underlying disease and congenital heart disease were covariates. Nasopharyngeal aspirates were positive for at least one virus in 236 patients. Rhinoviruses were detected in 85.6% of samples, with a preponderance of rhinovirus C (RV-C) (61.9%). Respiratory syncytial virus was detected in 59.8% and human coronavirus (HCoV) in 11% of the samples. Co-detections of two to five viruses were found in 78% of the patients. The detection of HCoV alone (adjusted relative risk (RR) 2.18; 95% CI 1.15–4.15) or in co-infection with RV-C (adjusted RR 2.37; 95% CI 1.23–4.58) was independently associated with PICU admission.

## Conclusions

The detection of HCoV alone or in co-infection with RV-C was independently associated with PICU admission in young children hospitalized for ALRI.

## Introduction

Acute lower respiratory tract infections (ALRI) are responsible for the death of approximately 160,000 neonates and over 760,000 infants annually [1]. Respiratory syncytial virus (RSV) and rhinovirus (RV) are the most frequent causes of acute respiratory infections in children [2–4]. Primary RSV infections may lead to severe bronchiolitis and pneumonia [5]. RV usually causes rhinopharyngitis [6], but may also cause illness of the lower respiratory tract and asthma exacerbations, and has been associated with severe ALRI in young children [7,8]. A few studies have investigated the viral profile in children attending the pediatric emergency room (ER) with ALRI and its association with disease severity [9–12], but findings vary. Moreover, the role of viral co-infections in illness severity has been controversial [10, 12, 13–15]. In young children who presented to an outpatient clinic or ER in Japan with acute respiratory illness, RSV, RV, parainfluenza viruses (PIV) and human metapneumovirus (HMPV) were the most prevalent viruses. Similarly to what was found in a previous study in Brazil [4], detection of RSV, alone or in co-detections, was associated with increased disease severity in children in Japan [9]. In Seattle, USA, RV was the most frequent virus detected in children less than three years old presenting to the ER with ALRI, and 52% of them required hospital admission [10]. In addition, RV viral load and co-detection with RSV was associated with more severe disease in that study [10]. In children younger than two years old who presented to a pediatric ER with ALRI in Malaysia, RSV and RV were the most frequently detected viruses and, although RSV was associated with a history of wheezing, virus detection was not associated with need for hospitalization [11]. In The Netherlands, the most frequent viruses detected in children presenting to the ER or outpatient clinic with ALRI were RSV, RV and human coronavirus (HCoV), and RSV detection correlated with longer duration of oxygen therapy [12]. However, there was no association between the virus species and hospital length of stay, and virus co-detections were not associated with disease outcome [12]. In the present study, a comprehensive polymerase-chain-reaction (PCR) panel of primers and probes was used to detect respiratory viruses, enabling virus

identification to the species level, in respiratory samples collected from children younger than three years with ALRI seen at the pediatric ER and subsequently admitted to hospital. Association of different respiratory viruses with disease severity, defined as need for admission to the pediatric intensive care unit (PICU), was assessed.

## Material and methods

### Patients and samples

This was a prospective cohort study conducted in a tertiary-care university hospital in Brazil. The study was approved by the Research Ethics Committee of the Clinical Hospital, School of Medicine of Ribeirão Preto, University of São Paulo, a tertiary-care facility in the city of Ribeirão Preto, state of Sao Paulo, Brazil. A written informed consent was obtained from parents/guardians (protocol number HCRP A07-020). All consecutive children younger than three years attending the pediatric emergency room with ALRI who were admitted to hospital from June 1<sup>st</sup>, 2008 to May 31<sup>st</sup>, 2009 were eligible for the study. ALRI was defined by the presence of cough, tachypnea, respiratory distress with prolonged expiratory time, and wheezing or crackles on auscultation. Patients with a diagnosis of bacterial pneumonia as indicated by clinical presentation and chest X-ray findings or a positive blood culture were excluded from analysis.

Demographic, clinical and outcome data were collected from patients' health records. Need for PICU admission was considered the main indicator of disease severity. Nasopharyngeal aspirates were collected from patients within the first 48 hours of hospitalization, as previously described [16] and they were split into aliquots: two 250 $\mu$ L aliquots mixed with 750 $\mu$ L of TRIzol (Invitrogen, Thermo Fisher Scientific, MA, USA) to DNA/RNA extraction and two 500 $\mu$ L backup aliquots mixed 1:1 with 500 $\mu$ L of viral transport medium, which consists of minimal essential medium with Eagle's salts plus 20% fetal bovine serum, 15% glycerol and 1% antibiotic-antimycotic solution (GIBco, Thermo). All backup aliquots were stored at -70°C until analysis.

### Rapid screening for RSV

Nasopharyngeal aspirates were routinely screened for RSV, by either a rapid chromatographic immunoassay for RSV antigen (Directigen EZ RSV Test, Becton Dickinson and Company, Franklin Lakes, NJ, USA) or indirect immunofluorescence (IF) assay. Directigen EZ RSV was performed on 250 $\mu$ L of nasopharyngeal aspirates following the manufacturer's protocol. Indirect IF assay was performed with RSV-specific monoclonal antibody (MAb 858-4; Millipore, MA, USA) diluted 1:100 in phosphate-buffered saline (GIBco, Thermo), revealed with Alexa Fluor 488-labeled donkey anti-mouse IgG (Life Technologies, Carlsbad, CA, USA) diluted 1:200 in phosphate-buffered saline. Slide preparation and IF protocols are available at <http://dx.doi.org/10.17504/protocols.io.w8rfhv6> and <http://dx.doi.org/10.17504/protocols.io.w8ufhww>.

### Detection of RNA and DNA respiratory viruses by real-time PCR

TRIzol aliquots from each sample were used to do DNA and RNA extractions. Total RNA was extracted following manufacturer's protocol with some adaptations (<http://dx.doi.org/10.17504/protocols.io.w8vfhw6>), and DNA-enriched fractions were used for DNA purification using the Wizard Genomic DNA purification kit (Promega, Madison, WI, USA) as per manufacturer's protocol. One microgram of total RNA was used in a reverse transcription reaction carried out with "Multiscribe reverse transcriptase" (Applied Biosystems, Thermo), primed

with random hexamers and following manufacturer's protocol. This random cDNA was used in TaqMan real-time PCR assays to detect RSV-A and B; HMPV A and B; human influenza viruses (FLU) A and B; and PIV 1 and 3. Human beta-actin housekeeping gene was used as internal controls in all assays. All PCR assays were performed using only one set of primers-probe per reaction, except for RSV and HMPV detection, which were tested in duplex format. Human bocavirus (HBoV) and human adenovirus (HAdV) were detected by qPCR in a single format assay (<http://dx.doi.org/10.17504/protocols.io.w8yfhw>). HCoV detection was performed by a nested RT-PCR method that uses primers targeting the RNA-dependent RNA polymerase gene [17]. Rhinovirus (RV) detection was performed by a two-step PCR method (<http://dx.doi.org/10.17504/protocols.io.w83fhyn>) [18, 19].

All PCR reactions were performed on a Thermocycler 7300 (Applied Biosystems, Thermo), using published primers and probes sequences [20]. All real-time PCR plates included appropriate negative controls matched to every step of the testing of nasopharyngeal aspirates. Negative controls were total RNA or DNA extracted from uninfected HeLa cells and ultrapure water treated in the same way as clinical samples.

### Statistical analysis

The analysis was made using SAS 9.4 (SAS/STAT User's Guide, Version 9.4, Cary, NC: SAS Institute Inc., 2013). Data were expressed as median (range) or number (%). Patients were grouped according to the need for PICU admission. Continuous variables between groups were compared by Mann-Whitney U test and categorical variables, by Fisher's exact test. Simple and multiple log-binomial regression models were constructed to assess associations of virus type with need for PICU admission. Relative risks (RR) and 95% confidence intervals (95%CI) were obtained after adjusting log-binomial regression models. Initially, simple log-binomial regression models were fitted, resulting in crude relative risks. Subsequently, the adjustment of multiple log-binomial regression models considering age, prematurity, the presence of an underlying disease and congenital heart disease as covariates, resulted in adjusted relative risks [21]. A 5% significance level was considered in all analysis.

### Results

Over the study period, nasopharyngeal aspirates were collected from 279 children seen at the pediatric emergency room with ALRI. The study population comprised 236 patients who had at least one respiratory virus detected in respiratory specimens. All patients were hospitalized. Twenty-five percent of patients ( $n = 60$ ) had comorbidities. The most common underlying diseases were neurological impairment ( $n = 17$ ), congenital anomalies ( $n = 13$ ) and chronic lung disease ( $n = 11$ ). Forty-seven patients (19.9%) were admitted to the PICU; 25 of them (53.2%) received invasive mechanical ventilation for a median time of 7 days (range 1 to 99 days). Length of PICU stay ranged from 1 to 254 days (median 9.5 days). Demographic data were not significantly different between patients admitted to the PICU compared with those who did not need PICU admission. However, use of systemic antibiotics and the presence of underlying diseases and congenital heart disease were more frequent in patients admitted to the PICU, and they also had a longer hospital length of stay, [Table 1](#).

The most frequently detected virus was RV (85.6%), followed by RSV (59.8%), HBoV (23.7%), HMPV (17.8%), HCoV (11.4%), HAdV (10.6%), PIV (10.2%) and FLU (8.5%). Co-detections were found in 182 (78%) patients, [S1 Table](#).

The results of multiple log-binomial regression analyses showed that the detection of HCoV alone (adjusted relative risk (RR) 2.18; 95% CI 1.15–4.15) or in co-infection with RV-C (adjusted RR 2.37; 95% CI 1.23–4.58) was independently associated with PICU admission, [S2 Table](#).

Table 1. Demographic and clinical data.

Characteristic	All (n = 236)	PICU admission (n = 47)	No PICU admission (n = 189)
Age (months)	5.2 (0.2–35)	3.6 (0.2–35)	5.9 (0.3–34)
Age < 6 months	125 (53)	29 (61.7)	96 (50.8)
Weight (kg)	6 (2–21)	5.2 (2.1–14)	6.2 (2–21)
Male gender	89 (38)	18 (38.3)	71 (37.6)
Prematurity	43 (18.2)	7 (14.9)	36 (19)
Underlying disease	60 (25)	21 (44.7)	39 (20.6)*
Congenital heart disease	23 (9.7)	9 (19)	14 (7.4)*
Use of systemic antibiotics	153 (64.8)	37 (78.7)	116 (61.4)*
Length of hospital stay (days)	8 (1–254)	18 (3–254)	7 (1–210)*

Data are expressed as median (range) or n (%). PICU, pediatric intensive care unit.

\*P < 0.05 for comparison between PICU admission and No PICU admission groups

<https://doi.org/10.1371/journal.pone.0217744.t001>

Eight patients (3.4%) died. Their median age was 3.5 months (range 2.3–9.3 months); six (75%) were female. Six patients (75%) who died had comorbidities: hydrocephalus (n = 3), Pompe disease (n = 1), Down syndrome (n = 1) and pulmonary hypoplasia (n = 1). Three patients had a single virus type detected in their respiratory samples (RV-C in two patients, and RV-A in one patient), two patients had dual viral co-detection (RV-C + RSV-B and RV-C + HCoV OC43), and three patients had triple viral co-detection (RSV-A + FLU-A + HBoV; RV-C + RSV-A + HMPV-A; and RV-C + HMPV-A + FLU-B). The causes of death were respiratory insufficiency (n = 4), cardiogenic shock (n = 2) and septic shock (n = 2). Of note, the patient who had the longest hospital stay (254 days) and also the longest duration of mechanical ventilation (99 days) had pulmonary hypoplasia, a severe underlying condition, and he ultimately died of sepsis.

## Discussion

In this study, viruses were detected in 85% of young children attending the emergency room with ALRI and subsequently admitted to the hospital. RV, especially RV-C, was the most frequently detected virus, followed by RSV. Twenty percent of patients were admitted to the PICU. Comorbidities and congenital heart disease were more frequent in patients who were admitted to the PICU. In addition, the detection of HCoV alone or in co-infection with RV-C was independently associated with PICU admission.

Similarly to our data, RV was also the most commonly detected virus in children younger than 3 years presenting to the Seattle Children's Hospital pediatric ER with a symptomatic respiratory tract infection, and the majority of them had lower respiratory tract infection and required hospitalization [10]. RV was also the most prevalent virus detected in children aged two weeks to 5 years admitted to a hospital with ALRI in Germany, exceeding the frequency of RSV [22]. Moreover, RV-C has been reported as the most frequent RV species and has been associated with severe disease in children less than 3 years old [8, 22].

Viral co-detection was found in approximately three-quarters of our study population, with RV/RSV as the most frequent duo. However, we observed that viral co-detection per se was not a risk factor for PICU admission. In keeping with this, a recent systematic review and meta-analysis showed that respiratory viral coinfection was not associated with need for hospitalization, intensive care admission or length of stay in children [15]. In addition, the number of detected viruses has not been associated with illness severity [14]. Indeed, surprisingly, an

inverse correlation between the clinical severity score and the number of viruses detected has been observed [23]. Nevertheless, a previous study performed in Southeast Brazil showed that coinfections, especially involving RSV, were associated with increased severity [13]. In the present study, RSV in co-detection with other viruses was not associated with increased disease severity.

HCoV was detected in 11% of children admitted to hospital with ALRI in our study, with a predominance of OC43 (40.7%) and 229E (33%) types. Although human coronaviruses most frequently cause common colds, they may cause severe respiratory diseases, such as severe acute respiratory syndrome [24]. We observed that one-third of children with HCoV infection and 41% of patients with HCoV in co-infection with RV-C were admitted to the PICU. Furthermore, we found that the detection of HCoV alone or in co-infection with RV-C was an independent risk factor for PICU admission. Similar to our results, HCoV was detected in 8.2% of hospitalized children aged  $3.2 \pm 3.9$  years with respiratory tract infection in New York, OC43 was the most prevalent type (40.1%), and 11% of patients with HCoV infection needed PICU admission. Additionally, the presence of chronic complex underlying conditions, including cardiovascular, genetic and respiratory diseases was associated with increased disease severity [25], which is corroborated by our data.

Almost two-thirds of our children were treated with systemic antibiotics. This high frequency of antibiotic use is similar to that reported in developed countries [26, 27]. Empiric antimicrobial therapy, driven by unfavorable clinical conditions in the face of probable bacterial infections, is likely to be initiated in the absence of immediate laboratory confirmation of virus detection. In the present study, the only rapid diagnostic test for respiratory virus available early at hospital admission was the RSV rapid antigen detection test. Therefore, the development of clinically relevant rapid tests for respiratory viruses is needed and shall help antimicrobial stewardship programs.

Since the present study was done in a single health center, this raises concern that epidemiologic data may not be generalizable. While this limitation is acknowledged, the results obtained are similar to those reported from Europe and North America [8, 22]. Another concern could be the assumption of disease etiology based on virus detection at one-time point. However, this is a practical diagnostic approach worldwide and all patients enrolled in the study had signs and symptoms of ALRI requiring hospitalization concomitantly with virus detection. It should also be noted that, although the diagnosis of bacterial pneumonia, as indicated by clinical presentation, chest X-ray findings or positive blood culture, was an exclusion criterion, it is difficult to discriminate viral and bacterial pneumonia in cases with negative bacterial cultures, because clinical presentation and chest radiography findings may overlap.

In conclusion, the detection of HCoV alone or in co-infection with RV-C was independently associated with PICU admission in young children hospitalized for ALRI. Rapid and reliable diagnostic tests for respiratory viruses associated with disease severity should be widely available to improve patient management and optimize healthcare resources.

## Supporting information

**S1 Table. Viruses detected in respiratory specimens of all patients (n = 236).**  
(DOCX)

**S2 Table. Viruses detected in respiratory samples and risk for admission to the pediatric intensive care unit (PICU).**  
(DOCX)

## Acknowledgments

We thank Marisa Akiko Iwamoto and Maria Lúcia Silva for expert technical assistance.

## Author Contributions

**Conceptualization:** Alessandra K. Matsuno, Talita B. Gagliardi, Renato T. Stein, Ana P. C. P. Carlotti.

**Data curation:** Talita B. Gagliardi, Flavia E. Paula, Luciano K. S. Luna.

**Formal analysis:** Alessandra K. Matsuno, Talita B. Gagliardi, Flavia E. Paula, Bruna L. S. Jesus, Renato T. Stein, Eurico Arruda.

**Funding acquisition:** Alessandra K. Matsuno, Eurico Arruda.

**Investigation:** Alessandra K. Matsuno, Talita B. Gagliardi, Flavia E. Paula, Eurico Arruda.

**Methodology:** Alessandra K. Matsuno, Talita B. Gagliardi, Flavia E. Paula, Luciano K. S. Luna, Bruna L. S. Jesus, Renato T. Stein, Davi C. Aragon, Ana P. C. P. Carlotti, Eurico Arruda.

**Project administration:** Alessandra K. Matsuno.

**Resources:** Renato T. Stein.

**Software:** Davi C. Aragon.

**Supervision:** Alessandra K. Matsuno, Eurico Arruda.

**Validation:** Luciano K. S. Luna, Davi C. Aragon.

**Visualization:** Luciano K. S. Luna.

**Writing – original draft:** Alessandra K. Matsuno, Talita B. Gagliardi, Flavia E. Paula, Ana P. C. P. Carlotti, Eurico Arruda.

**Writing – review & editing:** Alessandra K. Matsuno, Talita B. Gagliardi, Ana P. C. P. Carlotti, Eurico Arruda.

## References

1. Global Healthy Observatory Data Repository: World Acute Respiratory Infections. 2016 [accessed on 05 Dec 2017]. In: World Health Organization [internet]. Available from: <http://apps.who.int/gho/data/view.main.CM1002015WORLD-CH9?lang=en>.
2. Rakes GP, Arruda E, Ingram JM, Hoover GE, Zambrano JC, Hayden FG et al: Rhinovirus and respiratory syncytial virus in wheezing children requiring emergency care. *Am J Respir Crit Care Med*. 1999; 159:785–790 <https://doi.org/10.1164/ajrccm.159.3.9801052> PMID: 10051251
3. Sonogo M, Pellegrin MC, Becker G, Lazzerini M: Risk factors for mortality from acute lower respiratory infections (ALRI) in children under five years of age in low and middle-income countries: a systematic review and meta-analysis of observational studies. *PLoS ONE*. 2015; 10:e0116380. <https://doi.org/10.1371/journal.pone.0116380> PMID: 25635911
4. Arruda E, Jones MH, Escremim de Paula F, Chong D, Bugarin G, Notario G et al: The burden of single virus and viral coinfections on severe lower respiratory tract infections among preterm infants: a prospective birth cohort study in Brazil. *Pediatr Infect Dis J*. 2014; 33:997–1003. <https://doi.org/10.1097/INF.0000000000000349> PMID: 25361184
5. Walsh EE, Breese-Hall C. Respiratory syncytial virus. In: Bennett JE, Dolin R, Blaser MJ, editors. *Principles and Practice of Infectious Diseases*. 8th Ed. Philadelphia, MS: Elsevier Saunders; 2015. pp. 1948–1960.
6. Legg JP, Warner JA, Johnston SL, Warner JO: Frequency of detection of picornaviruses and seven other respiratory pathogens in infants. *Pediatr Infect Dis J*. 2005; 24:611–616. <https://doi.org/10.1186/1471-2334-11-41> PMID: 15999002

7. Fawcner-Corbett DW, Khoo SK, Duarte CM, Bezerra PG, Bochkov YA, Gern JE et al: Rhinovirus-C detection in children presenting with acute respiratory infection to hospital in Brazil. *J of Med Virol*. 2016; 88:58–63. <https://doi.org/10.1002/jmv.24300>
8. Lauinger IL, Bible JM, Halligan EP, Bangalore H, Tosas O, Aaron EJ et al: Patient characteristics and severity of human rhinovirus infections in children. *J Clin Virol*. 2013; 58:216–220. <https://doi.org/10.1016/j.jcv.2013.06.042> PMID: 23886500
9. Miyaji Y, Kobayashi M, Sugai K, Tsukagoshi H, Niwa S, Fujitsuka-Nozawa A, et al: Severity of respiratory signs and symptoms and virus profiles in Japanese children with acute respiratory illness. *Microbiol Immunol*. 2013; 57:811–821. <https://doi.org/10.1111/1348-0421.12102> PMID: 24117766
10. Chu HY, Englund JA, Strelitz B, Lacombe K, Jones C, Follmer K et al: Rhinovirus disease in children seeking care in a tertiary pediatric emergency department. *J Pediatric Infect Dis Soc*. 2016; 5:29–38. <https://doi.org/10.1093/jpids/piu099> PMID: 26908489
11. Nathan AM, Qiao YL, Jafar FL, Chan YF, Eg KP, Thauagnanam S, et al: Viruses and hospitalization for childhood lower respiratory tract infection in Malaysia: a prospective study. *PediatrRespirolCrit Care Med*. 2017; 1:46–51. [https://doi.org/10.4103/prcm.prcm\\_2\\_17](https://doi.org/10.4103/prcm.prcm_2_17)
12. Wishaupt JO, van der Ploeg T, de Groot R, Versteegh FG, Hartwig NG: Single- and multiple viral respiratory infections in children: disease and management cannot be related to a specific pathogen. *BMC Infect Dis*. 2017; 17:62. <https://doi.org/10.1186/s12879-016-2118-6> PMID: 28077074
13. Costa LF, Queiróz DAO, Silveira HL, Bernardino Neto M, de Paula NT, Oliveira TF et al: Human rhinovirus and disease severity in children. *Pediatrics*. 2014; 133:e312–e321. <https://doi.org/10.1542/peds.2013-2216> PMID: 24420808
14. Cebe-López M, Herberg J, Pardo-Seco J, Gómez-Carballa A, Martínón-Torres N, Salas A et al: Does viral co-infection influence the severity of acute respiratory infection in children? *PLoS One*. 2016; 11: e0152481. <https://doi.org/10.1371/journal.pone.0152481> PMID: 27096199
15. Scotta MC, Chakr VCBG, de Moura A, Becker RG, de Souza Ap, Jones MH et al: Respiratory viral coinfection and disease severity in children: a systematic review and meta-analysis. *J Clin Virol*. 2016; 80:45–56. <https://doi.org/10.1016/j.jcv.2016.04.019> PMID: 27155055
16. Cintra OA, Owa MA, Machado AA, Cervi MC, Figueiredo LT, Rocha GM et al: Occurrence and severity of infections caused by subgroup A and B respiratory syncytial virus in children in Southeast Brazil. *J Med Virol*. 2001; 65:408–412. PMID: 11536252
17. de Souza Luna LK, Heiser V, Regamey N, Panning M, Drexler JF, Mulangu S et al: Generic detection of coronaviruses and differentiation at the prototype strain level by reverse transcription-PCR and non-fluorescent low-density microarray. *J Clin Microbiol*. 2007; 45:1049–1052. <https://doi.org/10.1128/JCM.02426-06> PMID: 17229859
18. Lee WM, Lemanske RF, Evans M, Vang F, Pappas T, Gangnon R et al: Human rhinovirus species and season of infection determine illness severity. *Am J Respir Crit Care Med*. 2012; 186:886–891. <https://doi.org/10.1164/rccm.201202-0330OC> PMID: 22923659
19. Lee WM, Kiesner C, Pappas T, Lee L, Grindle K, Jartti T et al: A diverse group of previously unrecognized human rhinoviruses are common causes of respiratory illnesses in infants. *PLoS ONE*. 2007; 2: e966. <https://doi.org/10.1371/journal.pone.0000966> PMID: 17912345
20. Proenca-Modena JL, Pereira Valera FC, Jacob MG, Buzatto GP, Saturno TH, Lopes L et al: High rates of detection of respiratory viruses in tonsillar tissues from children with chronic adenotonsillar disease. *PLoS One*. 2012; 7:e42136. <https://doi.org/10.1371/journal.pone.0042136> PMID: 22870291
21. Blizzard L, Hosmer DW: Parameter estimation and goodness-of-fit in log binomial regression. *Biom J*. 2006; 48:5–22. PMID: 16544809
22. Renwick N, Schweiger B, Kapoor V, Liu Z, Villari J, Bullmann R et al: A recently identified rhinovirus genotype is associated with severe respiratory-tract infection in children in Germany. *J Infect Dis*. 2007; 196:1754–1760. <https://doi.org/10.1086/524312> PMID: 18190255
23. Do LAH, Bryant JE, Tran AT, Nguyen BH, Tran TTL, Quynh HT et al: Respiratory syncytial virus and other viral infections among children under two years old in Southern Vietnam 2009–2010: clinical characteristics and disease severity. *PLoS One*. 2016; 11:e0160606. <https://doi.org/10.1371/journal.pone.0160606> PMID: 27500954
24. Zhang S, Tuo J, Huang X, Zhu X, Zhang DM, Zhou K et al: Epidemiology characteristics of human coronaviruses in patients with respiratory infection symptoms and phylogenetic analysis of HCoV-OC43 during 2010–2015 in Guangzhou. *PLoS One*. 2018; 13:e0191789. <https://doi.org/10.1371/journal.pone.0191789> PMID: 29377913
25. Varghese L, Zachariah P, Vargas C, La Russa P, Demmer RT, Furuya YE et al: Epidemiology and clinical features of human coronaviruses in the pediatric population. *J Pediatric Infect Dis Soc*. 2017. <https://doi.org/10.1093/jpids/pix027>

26. De Luca M, Donà D, Montagnani C, Lo Vecchio A, Romanengo M, Tagliabue C et al: Antibiotic prescriptions and prophylaxis in Italian children. Is it time to change? Data from the ARPEC project. *PLoS One*. 2016; 11:e0154662. <https://doi.org/10.1371/journal.pone.0154662> PMID: 27182926
27. Kelesidis T, Braykov N, Uslan DZ, Morgan DJ, Gandra S, Johannsson B, et al: Indications and types of antibiotic agents used in 6 acute care hospitals, 2009–2010: A pragmatic retrospective observational study. *Infect Control Hosp Epidemiol*. 2016; 37:70–79. <https://doi.org/10.1017/ice.2015.226> PMID: 26456803