RESEARCH ARTICLE

Post grafting time significantly influences royal jelly yield and content of macro and trace elements

Saad N. Al-Kahtani^{1*}, El-Kazafy A. Taha^{1,2}

1 Arid Land Agriculture Department, College of Agricultural Sciences and Foods, King Faisal University, Al-Ahsa, Saudi Arabia, **2** Economic Entomology Department, Faculty of Agriculture, Kafrelsheikh University, Kafrelsheikh, Egypt

* salkahtani@kfu.edu.sa

OPEN ACCESS

Citation: Al-Kahtani SN, Taha E-KA (2020) Post grafting time significantly influences royal jelly yield and content of macro and trace elements. PLoS ONE 15(9): e0238751. <https://doi.org/10.1371/journal.pone.0238751>

Editor: Shahid Farooq, Harran University, TURKEY

Received: July 24, 2020

Accepted: August 21, 2020

Published: September 8, 2020

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0238751>

Copyright: © 2020 Al-Kahtani, Taha. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript.

Funding: The current study was financially supported by the Deanship of Scientific Research, King Faisal University with a project number 160075.

Abstract

Royal jelly (RJ) is commercially harvested after the 4th day of queen larval age. In the current study, it was harvested after 24, 48, 72, and 96 hours after grafting of 1-day larval age queens to investigate changes in macro and trace elements associated with harvesting time. The RJ yields were significantly affected by harvest time, and the highest yield was obtained 72 hours after grafting. The highest phosphorus (P) and zinc (Zn) contents were obtained from RJ harvested 24 hours after grafting. Royal jelly harvested 48 hours after grafting had the highest concentrations of magnesium (Mg), calcium (Ca), potassium (K), sodium (Na), iron (Fe), and manganese (Mn). Likewise, RJ harvested 96 hours after grafting had higher concentrations of copper (Cu). Royal jelly harvested 72 hours after grafting showed the second rank for P, Mg, Ca, K, Na, Fe, Cu, and Mn concentrations. In descending order, P, Mg, Ca, and K were the most dominant elements in RJ harvested at different times after grafting. The Mg, Ca, K, Na, Cu, and Mn concentrations in RJ were all positively correlated, and P, Fe, and Zn were positively correlated. The P and Zn were negatively correlated with Ca, Cu, and Mn. It was concluded that macro and trace element contents in RJ can differ depending on the harvest time after grafting. We recommend harvesting RJ at 72 hours after grafting for possible use as healthy nutritional human food supplement.

Introduction

Royal jelly (RJ) is a valuable bee product released by the hypopharyngeal and mandibular glands located in the heads of worker honeybees. These glands become more active by the time the workers reach an age of 5–15 days [1–3]. Royal jelly is the diet of young honeybee larvae and plays the main role in caste differentiation of honeybees. The queen larvae receive RJ for 5 days, while the worker larvae receive RJ only for 3 days, and then receive a mixture of honey, pollen and water for 2 days. Thereafter, a queen of honeybee lives for several years, and the worker bees live for a few months [4].

Royal jelly is a milky substance composed of proteins, sugars, lipids, vitamins, and individual elements [5–9]. It can be harvested in relatively small quantities from normally built queen

Competing interests: The authors have declared that no competing interests exist.

cells [10], or commercially in comparatively larger amounts by using an artificial queen-rearing method [11–15].

The elements content in RJ have determined as ash [5, 6, 8, 9, 16–22]. The standard amount of ash in RJ is reportedly approximately 0.8–3% [23]. The element concentration of RJ produced in Saudi Arabia ranges from 2.16% in RJ harvested 24 hours after grafting to 3.03% in RJ harvested 72 hours after grafting [8]. However, the elements content in RJ produced have differed affected by the botanical and geographical origins and other biological factors related to the worker bees [24, 25].

Royal jelly is an important functional food parameter that possesses several health-promoting characteristics. It has been widely used in commercial healthy foods and medical products in many countries. Royal jelly has been approved to possess several functional properties such as antioxidant, antibacterial, antihypercholesterolemic, anti-inflammatory, antitumor, disinfectant, vasodilative and hypotensive activities [18, 26]. In addition, it has used to supplement many diseases, including cardiovascular, diabetes, cancer and Alzheimer's disease [26]. The daily requirements of the individual elements in human nutrition have determined as 800–1200 mg P, 800 mg K, 800–900 mg Ca, 300–400 mg Mg, 6–22 mg Zn, 10–20 mg Fe, 4–5 mg Mn and 1–3 mg Cu [27, 28]. Trace elements play a vital role in the biomedical activities of RJ, as these have a variety of biological functions [18].

The nutritional and therapeutic characteristics of RJ have led to its inclusion in an increasing number of food products [7, 9, 18, 29–34]. Royal jelly is commercially harvested at the end of the 4th day of queen larval age [2, 8, 35], but there are few reports on the element content of RJ harvested before this time. In the present study the effects of harvesting time on the macro and trace element contents of RJ were investigated, with a particular focus on elements that are essential for human nutrition [phosphorus (P), potassium (K), sodium (Na), magnesium (Mg), calcium (Ca), iron (Fe), zinc (Zn), copper (Cu), and manganese (Mn)].

Materials and methods

Royal jelly sampling

The RJ used in the study was produced at the apiary of the Training and Research Station, King Faisal University, Al-Ahsa (25°25'46"N, 49°37'19"E; 121 m above sea level), Saudi Arabia during February and March in 2019. One colony of the hybrid Carniolan honeybee (*Apis mellifera carnica* Pollmann) was selected as a breeder colony to provide all larvae used for grafting. Ten colonies of 10 frames each were selected for RJ production. Fifty wax cups were grafted with larvae that were 24 hours of age, and the cups were introduced into each colony. Royal jelly was harvested from the accepted queen cell cups 24, 48, 72, and 96 hours after grafting in previously weighed small bottles. The bottles with RJ were weighed using an electronic balance. The mean yield of RJ (mg/queen cell) for every harvest time was calculated by subtraction [8].

Macro and trace elements extraction and analysis

Wet digestion with nitric acid [36] was used for element extraction. One gram of RJ from each harvest was digested in a Kjeldahl flask with 10 ml of 75% nitric acid (HNO₃) for the oxidation of carbonaceous matter. The contents of the flask were heated 100–120 °C, to evaporate the acid. Drops of perchloric acid (HClO₄) were added until all the organic matter was oxidized. This point was reached when no further darkening of the solution occurred on continuous heating and a clear solution was obtained. It was cooled and gauged to 50 ml with distilled water. A blank experiment was carried out by adding the same amount of nitric acid to 1 ml distilled water. An atomic absorption spectrophotometer (Avanta E, GBC, Australia) was used

to detect calcium (Ca), copper (Cu), iron (Fe), magnesium (Mg), manganese (Mn), potassium (K), sodium (Na) and zinc (Zn). A UV-VIS spectrophotometer (UV-2550 Shimadzu, Kyoto, Japan) was used to detect the phosphorus (P) concentrations.

Statistical analysis

The difference between harvesting times was tested by one-way analysis of variance (ANOVA), which indicated significant difference for harvesting times. The normality in data was tested by Shapiro-Wilk normality test, which indicated data were normally distributed. Therefore, the analyses were performed on original data. The ANOVA and Pearson's correlational coefficients were used to assess differences and associations between the elements investigated via the PROC GLM function in SAS version 9.1 [37]. Duncan's [38] multiple range test was used to compare means.

Results

Royal jelly yields were significantly ($p < 0.01$) affected by harvest time (Table 1). The highest yield (318.5 mg/queen cell) was obtained 72 hours after grafting. The lowest yield (70.4 mg/queen cell) was obtained 24 hours after grafting (Fig 1).

The mean concentrations (mg/kg) of P (8636.7–10113.3), Mg (6920.5–9545.2), Ca (2806.2–4334.5), K (1630.0–4334.5), Na (193.0–675.0), Fe (43.4–76.3), Zn (32.8–78.7), Cu (34.4–39.9) and Mn (0.5–1.9) differed significantly ($p < 0.01$) at different harvesting times (Table 1). Royal jelly harvested 48 hours after grafting had the highest mean contents (mg/kg) of Mg (9545.2), Ca (4334.5), K (2126.2), Na (675.0), Fe (76.3) and Mn (1.9). The highest concentrations of P (10113.3) and Zn (78.7) were obtained from RJ harvested 24 hours after grafting. Royal jelly harvested 96 hours after grafting had higher concentrations of Cu (39.9). The lowest concentrations of Mg (6920.5), Ca (2806.2), K (1630.3), Na (193.2), Cu (34.4) and Mn (0.5) were present in RJ harvested 24 hours after grafting, and the lowest concentrations of P (8636.7), Fe (52.4), and Zn (32.8) were present in RJ harvested 96 hours after grafting (Table 2).

There were relative abundances of P (46.3, 35.4, 37.3, and 40.5% of the total elements quantified), Mg (31.7, 36.6, 36.9, 64'and 33.1%), Ca (12.8, 16.6, 14.7 and 16.6%), K (7.5, 8.2, 8.0 and 8.2%) and Na (0.9, 2.6, 2.6, and 1.0%) for RJ harvested 24, 48, 72 and 96 hours after grafting, respectively (Table 3).

Concentrations of Mg, Ca, K, Na, Cu, and Mn in RJ were positively correlated ($r = 0.54$ – 0.98 , $p < 0.01$) (Table 4). Concentrations of P, Fe, and Zn were positively correlated ($r = 0.61$ –

Table 1. Analysis of variance of yield and nutrient concentration in royal jelly.

Variable	SS	MS	F value	P value
RJ yield	125893.4	41964.5	1523.2	< 0.0001*
P	6198224.3	2066074.7	679959.0	< 0.0001*
Mg	32852748.7	10950916.2	2879108.0	< 0.0001*
Ca	7045039.8	2348346.6	697253.0	< 0.0001*
K	882608.6	294202.8	77788.1	< 0.0001*
Na	1212430.9	404143.6	150918.0	< 0.0001*
Fe	1876.9	625.6	312.8	< 0.0001*
Zn	6788.3	2262.7	1216.2	< 0.0001*
Cu	121.2	40.4	35.7	< 0.0001*
Mn	6.7	2.2	1393.5	< 0.0001*

Source of variation = post-grafting time, degree of freedom = 4, SS = sum of squares, MS = mean squares, * = significant.

<https://doi.org/10.1371/journal.pone.0238751.t001>

Fig 1. Effect of harvesting time on royal jelly yield (mg/queen cell).

<https://doi.org/10.1371/journal.pone.0238751.g001>

0.95, $P < 0.01$). Fe was positively correlated with K ($r = 0.47$, $p < 0.05$) and Mg ($r = 0.61$, $p < 0.01$). P and Zn were negatively correlated with Ca, Cu, and Mn ($r = -0.51$ to -0.85 , $p < 0.01$).

Discussion

All larvae used for grafting in the current study were obtained from the same queen, and reared for RJ production under the same colony conditions, so differences between the amounts of RJ/queen cell can reasonably be summarized to be due to differences in harvesting times [8]. Royal jelly yields in queen cells were related to the amounts of RJ required by each larval instar, which increases with larval age. The amount of RJ in a queen cell initially increased gradually with larva age, then it peaks on the 4th day of larval age, then it decreased on the 5th day of larval age. In commercial systems RJ is harvested on the 4th day of larval age because that is when the yields are maximal [2, 8, 12, 14, 35].

In the current study, all RJ samples were produced in the same colonies, and all larvae used for grafting were derived from the same queen, so the observed differences in concentrations

Table 2. Effects of different harvesting times on concentrations (mg/kg) of macro and trace elements in royal jelly.

Element	Time (hours) after grafting				Mean
	24	48	72	96	
Macro elements					
P	10113.3 ± 2.3 ^a	9222.4 ± 1.8 ^c	9245.3 ± 1.7 ^b	8636.7 ± 1.4 ^d	9304.4
Mg	6920.5 ± 1.8 ^d	9545.2 ± 2.4 ^a	9166.7 ± 1.7 ^b	7053.3 ± 1.6 ^c	8171.4
Ca	2806.2 ± 2.1 ^d	4334.5 ± 1.4 ^a	3652.4 ± 1.8 ^b	3544.7 ± 1.5 ^c	3584.5
K	1630.3 ± 2.2 ^d	2126.2 ± 2.2 ^a	1980.4 ± 1.3 ^b	1750.2 ± 1.8 ^c	1871.8
Na	193.2 ± 2.0 ^d	675.1 ± 1.5 ^a	640.0 ± 1.4 ^b	205.3 ± 1.5 ^c	428.4
Trace elements					
Fe	72.3 ± 1.4 ^b	76.3 ± 1.7 ^a	73.4 ± 1.3 ^b	52.4 ± 1.2 ^c	74.0
Zn	78.7 ± 1.6 ^a	55.1 ± 1.5 ^b	41.8 ± 1.2 ^c	32.8 ± 0.9 ^d	52.1
Cu	34.4 ± 1.3 ^b	39.5 ± 1.0 ^a	39.1 ± 0.4 ^a	39.9 ± 1.2 ^a	38.2
Mn	0.5 ± 0.0 ^c	1.9 ± 0.1 ^a	1.6 ± 0.0 ^a	1.1 ± 0.0 ^b	1.3

The values shown are means ± standard deviation. The means of each row followed by a different letter differed significantly.

<https://doi.org/10.1371/journal.pone.0238751.t002>

Table 3. Relative abundance (%) of macro elements in royal jelly at different harvesting times.

Elements	Time (hrs) after grafting				Average
	24	48	72	96	
P	46.3	35.4	37.2	40.5	39.8
Mg	31.7	36.6	36.9	33.1	34.6
Ca	12.8	16.6	14.7	16.6	15.2
K	7.5	8.2	8.0	8.2	8.0
Na	0.9	2.6	2.6	1.0	1.7

<https://doi.org/10.1371/journal.pone.0238751.t003>

of macro and trace elements can reasonably be summarized to be due to the type of RJ introduced to the larvae at the different larval instars. There were numerous significant differences in element concentrations in RJ harvested at different times after grafting. Wang et al. [5] also reported significant differences in the concentrations of elements in RJ harvested at different times.

The concentrations detected in the current study were within the normal limits for most elements [23], and within the range of Mn in RJ produced in China [5]. Notably however, the Ca, Na, Mg, Fe, Zn, and Cu concentrations were higher than the ranges in RJ produced in China [5], Lithuania [34], and Bingol [9], and the concentrations of Ca, Fe, and Mn were higher than the concentrations in RJ produced in Jordan [20]. Conversely, K concentrations in the RJ samples in the current study were lower than those in RJ produced in China [5], Lithuania [34], and Bingol [9].

Concentrations of Mg, Ca, K, Na, and Mn in RJ harvested 48 hours after grafting were higher than those in RJ harvested 72 hours after grafting and 96 hours after grafting, and the lowest concentrations were detected in RJ harvested 24 hours after grafting. The high concentrations of these elements in RJ harvested at 48 and 72 hours after grafting were related to the requirements of larvae at these larval ages [8]. The respective concentrations of Mg, Ca, K, Na, Cu, and Mn in RJ harvested 48 hours after grafting were 138%, 154%, 130%, 349%, 115%, and 380% of the corresponding concentrations in RJ harvested 24 hours after grafting. In RJ harvested 72 hours after grafting the relative amounts were 132%, 130%, 121%, 331%, 114%, and 320%.

Generally, the macro elements could be arranged in a descending order: P > Mg > Ca > K > Na, and they constituted 39.8%, 34.6%, 15.2%, 8.0%, and 1.7% of the total elements quantified. Wang et al. [5] reported that K, Mg, Ca, Na, and Fe were the most abundant elements in RJ produced in China, Adaškevičiūtė et al. [34] reported that K, P, Mg, Na, and Ca were the

Table 4. Pearson's correlation coefficients for macro and trace elements in royal jelly.

	P	Mg	Ca	K	Na	Fe	Zn	Cu	Mn
P	1								
Mg	-0.24	1							
Ca	-0.61**	0.83**	1						
K	-0.40*	0.97**	0.94**	1					
Na	-0.22	0.99**	0.80**	0.96**	1				
Fe	0.61**	0.61**	0.19	0.47*	0.62**	1			
Zn	0.95**	-0.27	-0.51**	-0.37	-0.27	0.73**	1		
Cu	-0.85**	0.55**	0.75**	0.65**	0.54**	-0.41*	-0.81**	1	
Mn	-0.56**	0.93**	0.95**	0.98**	0.92**	-0.12	-0.54**	0.78**	1

* $p < 0.05$ (two-tailed)

** $p < 0.01$ (two-tailed).

<https://doi.org/10.1371/journal.pone.0238751.t004>

most abundant elements in RJ from Lithuania, and Bengü et al. [9] reported that K, Mn, Mg, Ca, and Na were the most abundant elements in RJ from Bingol. Variations in macro element concentrations in RJ produced in different regions are reportedly affected by floral resources, differences in climate and geography [31, 39], and the elements in pollen consumed by nurse worker bees [40, 41].

Trace elements play a critical role in the biological characteristics of RJ due to their biological activities [9]. Fe, Zn, Cu, and Mn constituted less than 0.8% of the total estimated elements in RJ harvested at different times after grafting. Relatively similar Fe, Zn, and Cu concentrations were reportedly detected in RJ produced in China [5], Lithuania [34], and Bingol [9]. Conversely, a high concentration of Mn was reportedly detected in RJ from Bingol [9].

Conclusions

In the current study, the concentrations of macro and trace elements in RJ differed significantly at different times after grafting. Royal jelly harvested at 48 and 72 hours after grafting had high concentrations of Mg, Ca, K, Na, Fe, Cu, and Mn and it can be used as a healthy nutritional human food supplement.

Author Contributions

Conceptualization: Saad N. Al-Kahtani.

Data curation: El-Kazafy A. Taha.

Formal analysis: Saad N. Al-Kahtani, El-Kazafy A. Taha.

Funding acquisition: Saad N. Al-Kahtani.

Investigation: El-Kazafy A. Taha.

Methodology: El-Kazafy A. Taha.

Project administration: Saad N. Al-Kahtani.

Resources: Saad N. Al-Kahtani.

Software: El-Kazafy A. Taha.

Supervision: Saad N. Al-Kahtani.

Validation: El-Kazafy A. Taha.

Visualization: El-Kazafy A. Taha.

Writing – original draft: El-Kazafy A. Taha.

References

1. Deseyn J, Billen J. Age-dependent morphology and ultrastructure of the hypopharyngeal gland of *Apis mellifera* workers (Hymenoptera, Apidae). *Apidologie*. 2005; 36: 49–57.
2. Taha EKA. Studies on honeybee (*Apis mellifera* L.). Unpublished Ph.D. Thesis, Tanta University. 2005.
3. Balkanska R, Al L, Mărghițaș, Pavel CI, Ignatova M, Tomoș LI. Comparison of physicochemical parameters in royal jelly from Romania and Bulgaria. *Bull UASVM Anim Sci Biotechnol*. 2013; 70: 117–121.
4. Moselhy WA, Fawzy AM, Kamel AA. An evaluation of the potent antimicrobial effects and unsaponifiable matter analysis of the royal jelly. *Life Sci J*. 2013; 10: 290–296.
5. Wang Y, Ma L, Zhang W, Cui X, Wang H, Xu B. Comparison of the nutrient composition of royal jelly and worker jelly of honey bees (*Apis mellifera*). *Apidologie*. 2016; 47: 48–56.

6. Margaoan R, Marghitas LA, Dezmirean DS, Bobis O, Bonta V, Catana C, et al. Comparative study on quality parameters of royal jelly, apilarnil and queen bee larvae triturate. *Bull UASVM Anim Sci Biotechnol*. 2017; 74: 51–58.
7. Maghsoudlou A, Mahoonak AS, Mohebodini H, Toldra F. Royal jelly: chemistry, storage and bioactivities. *J Apic Sci*. 2019; 63: 17–40.
8. Taha EKA, Al-Kahtani S. Effect of harvest time on royal jelly yield and chemical composition. *J Kans Entomol Soc*. 2019.
9. Bengü AŞ, Ayna A, Özbolat S, Tunç A, Çiftçi M, Darendelloğlu E. Content and antimicrobial activities of bingol royal jelly. *Turk J Agric Nat Sci*. 2020; 7: 480–489.
10. Taha EKA. Effect of transferring the apiaries on activity of honeybee colonies. Unpublished M. Sc. Thesis, Tanta University. 2000.
11. Dedej S. Effect of double grafting in queen rearing. *Ape Nostra Amica*. 1994; 16: 11–14.
12. Al-Fattah MAA. Factors increasing the acceptance of transplanted larvae in artificial queen cells and royal jelly production of the honeybee colonies. *J Agric Sci Mansoura Univ*. 1996; 21: 4555–4563.
13. Chen S, Su S, Lin X. An introduction to high-yielding royal jelly production methods in China. *Bee World*. 2002; 83: 69–77.
14. Taha EKA, Shawer MB, El-Dakhkhni TN, Helal RM. Effect of nectar and pollen flora on royal jelly and honey production. In *Proceedings of the 6th international conference of Arab Apiculture*. Saudi Arabia. 2009. p. 73.
15. Sharaf El-Din HA. Studies on royal jelly production in honeybee colonies. Unpublished M.Sc. Thesis, Cairo University. 2010.
16. Wongcha V. Seasonal variation of chemical composition of royal jelly produced in Thailand. *Thammasat International Journal of Science and Technology*. 2002; 7: 1–8.
17. Zheng HQ, Hu FL, Dietemann V. Changes in composition of royal jelly harvested at different times: consequences for quality standards. *Apidologie*. 2011; 42: 39–47.
18. Ramadan MF, Al-Ghamdi A. Bioactive compounds and health-promoting properties of royal jelly: a review. *J Funct Foods*. 2012; 4: 39–52.
19. Melliou E, Chinou I. Chemistry and bioactivities of royal jelly. *Stud Nat Prod Chem*. 2014; 43: 261–290.
20. Nabas Z, Haddadin M, Haddadin J, Nazer I. Chemical composition of royal jelly and effects of synbiotic with two different locally isolated probiotic strains on antioxidant activities. *Pol J Food Nutr Sci*. 2014; 64: 171–180.
21. Hamid NA, Bakar ABA, Mohamed M. Phytochemical analysis and GC–MS profile of royal jelly from selected areas in Malaysia. *Malays Appl Biol*. 2018; 47: 101–107.
22. Kazemi V, Eskafi M, Saeedi M, Manayi A, Hadjiakhoondi A. Physicochemical properties of royal jelly and comparison of commercial with raw specimens. *Jundishapur J Nat Pharm Prod*. 2019; 14: 1–3.
23. Sabatini AG, Marcazzan GL, Caboni MF, Bogdanov S, de Almeida-Muradian LB. Quality and standardisation of royal jelly. *J ApiProduct ApiMed Sci*. 2009; 1: 1–6.
24. Stocker A, Schramel P, Kettrup A, Bengsch E. Trace and mineral elements in royal jelly and homeostatic effects. *J Trace Elem Med Biol*. 2005; 19: 183–189. <https://doi.org/10.1016/j.jtemb.2005.08.004> PMID: 16325534
25. Garcia-Amoedo LH, Almeida-Muradian LBd. Physicochemical composition of pure and adulterated royal jelly. *Quim Nova*. 2007; 30: 257–259.
26. Ahmad S, Campos MG, Fratini F, Altaye SZ, Li J. New Insights into the Biological and Pharmaceutical Properties of Royal Jelly. *Int. J. Mol. Sci*. 2020; 21: 382–407.
27. Tayar M, Cibik R. Food chemical. (2nd ed., 2011; pp. 157–178). Bursa, Dora.
28. Demirci M. Food chemical. (7th ed., 2014; pp. 148–156). Istanbul, Hat.
29. Balkanska R, Karadjova I, Ignatova M. Comparative analyses of chemical composition of royal jelly and drone brood. *Bulg Chem Commun*. 2014; 46: 412–416.
30. Cornara L, Biagi M, Xiao J, Burlando B. Therapeutic properties of bioactive compounds from different honeybee products. *Front pharmacol*. 2017; 8: 412. <https://doi.org/10.3389/fphar.2017.00412> PMID: 28701955
31. Kocot J, Kielczykowska M, Luchowska-Kocot D, Kurzepa J, Musik I. Antioxidant potential of propolis, bee pollen, and royal jelly: possible medical application. *Oxidative Med Cell Longev*. 2018; 7074209.
32. Nath KGRA Krishna SBA, Smrithi S Nair AJ, Sugunan VS. Biochemical characterization and biological evaluation of royal jelly from Apis cerana. *J Food Technol Food Chem*. 2019; 2: 101–107.

33. Kunugi H, Ali AM. Royal jelly and its components promote healthy aging and longevity: from animal models to humans. *Int J Mol Sci.* 2019; 20: 4662–4687.
34. Adaškevičiūtė V, Kaškonienė V, Kaškonas P, Barčauskaitė K, Maruška A. Comparison of physico-chemical properties of bee pollen with other bee products. *Biomolecules.* 2019; 9: 819–841.
35. Lercker G, Caboni M, Vecchi M, Sabatini A, Nanetti A, Piana L. Composizione della frazione glucidica della gelatina reale e della gelatina delle api operaie in relazione all'età larvale. *Apicoltura.* 1985; 1: 123–139.
36. AOAC. Official methods of analysis. 17th ed. Washington, DC, USA: Association of Official Analytical Chemists; 2000.
37. Institute SAS. SAS/STAT user's guide, release 9.1. Cary, NC: SAS Institute Inc.; 2003.
38. Duncan DB. Multiple range and multiple F tests. *Biometrics.* 1955; 11: 1–42.
39. Siğ AK, Öz-Siğ Ö, Güney M. Royal jelly: a natural therapeutic? *Ortadogu Tıp Derg.* 2019; 11: 333–341.
40. Taha EKA. Chemical composition and amounts of mineral elements in honeybee-collected pollen in relation to botanical origin. *J Apic Sci.* 2015; 59: 75–81.
41. Taha EKA, Al-Kahtani S. Macro- and trace elements content in honeybee pollen loads in relation to the harvest season. *Saudi J Biol Sci.* 2020; 27: 1797–1800. <https://doi.org/10.1016/j.sjbs.2020.05.019> PMID: 32565698