CORRECTION

Correction: Biogeographical patterns and speciation of the genus *Pinguicula* (Lentibulariaceae) inferred by phylogenetic analyses

Hiro Shimai, Hiroaki Setoguchi, David L. Roberts, Miao Sun

[Fig 5](#) is incorrect. The authors have provided a corrected version here.

OPEN ACCESS

Citation: Shimai H, Setoguchi H, Roberts DL, Sun M (2021) Correction: Biogeographical patterns and speciation of the genus *Pinguicula* (Lentibulariaceae) inferred by phylogenetic analyses. PLoS ONE 16(12): e0261600. <https://doi.org/10.1371/journal.pone.0261600>

Published: December 14, 2021

Copyright: © 2021 Shimai et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Fig 5. Phylogenetic comparison of nrDNA (ITS) and concatenated cpDNA. The figure shows topological incongruence between the ITS and combined cpDNA (*matK* + *rpl32-trnL*) trees. Vertical bars and connected lines are coloured based on major clades in the ITS tree; red for Clade I (the southeastern USA), green for Clade III (South America), blue for Clades IV and V (the temperate Northern Hemisphere), gold for Clades VI, VII, and VIII (Mexico and Central America), purple for Clade IX (Cuba), and black for others and the outgroup.

<https://doi.org/10.1371/journal.pone.0261600.g001>

Reference

1. Shimai H, Setoguchi H, Roberts DL, Sun M (2021) Biogeographical patterns and speciation of the genus *Pinguicula* (Lentibulariaceae) inferred by phylogenetic analyses. PLoS ONE 16(6): e0252581. <https://doi.org/10.1371/journal.pone.0252581> PMID: 34097720