

CORRECTION

Correction: Feelings of Hopelessness in Midlife and Cognitive Health in Later Life: A Prospective Population-Based Cohort Study

Krister Håkansson, Hilikka Soininen, Bengt Winblad, Miia Kivipelto

The images for Figs 2 and 3 are incorrectly switched. The image that appears as Fig 2 should be Fig 3, and the image that appears as Fig 3 should be Fig 2. The figure captions appear in the correct order. Please view the correct Figs 2 and 3 here.



 OPEN ACCESS

Citation: Håkansson K, Soininen H, Winblad B, Kivipelto M (2015) Correction: Feelings of Hopelessness in Midlife and Cognitive Health in Later Life: A Prospective Population-Based Cohort Study. PLoS ONE 10(11): e0142465. doi:10.1371/journal.pone.0142465

Published: November 4, 2015

Copyright: © 2015 Håkansson et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Fig 2. Levels of hopelessness at baseline and follow-up for the different outcome groups. Changes in hopelessness scores from baseline until follow up for those who at follow-up were either without cognitive impairment or were diagnosed with any cognitive impairment, mild cognitive impairment or Alzheimer's disease. No changes from baseline to follow-up were statistically significant within any of the outcome categories (Student t-test for independent samples). All differences between the non-impaired group and any of the cognitive impairment groups were statistically significant, both at baseline and at follow-up (* = $p \leq 0.05$, ** = $p \leq 0.01$, *** = $p \leq 0.001$ as indicated by Student's t-test for paired samples). The graph is based on scores from participants with measurements of hopelessness both at baseline and follow-up (N = 1246).

doi:10.1371/journal.pone.0142465.g001




Fig 3. Hopelessness and cognitive impairment for ApoE4 carriers and non-carriers. Odds ratios from logistic regressions after adjustments for age, education, gender, BMI, blood pressure, cholesterol, residence area, occupation, physical activity, smoking, marital status, and depression. The risk is in comparison to ApoE4 non-carriers with low levels of hopelessness at midlife (OR = 1). Stars indicate the level of statistical significance (three levels <0.05, <0.01, <0.001).

doi:10.1371/journal.pone.0142465.g002

Reference

1. Håkansson K, Soininen H, Winblad B, Kivipelto M (2015) Feelings of Hopelessness in Midlife and Cognitive Health in Later Life: A Prospective Population-Based Cohort Study. PLoS ONE 10(10): e0140261. doi: [10.1371/journal.pone.0140261](https://doi.org/10.1371/journal.pone.0140261) PMID: [26460971](https://pubmed.ncbi.nlm.nih.gov/26460971/)