

CORRECTION

Correction: Offshore Habitat Preference of Overwintering Juvenile and Adult Black Sea Bass, *Centropristis striata*, and the Relationship to Year-Class Success

Alicia S. Miller, Gary R. Shepherd, Paula S. Fratantoni

The images for Figs 4–6 are incorrectly switched. The image that appears as [Fig 6](#) should be [Fig 5](#), the image that appears as [Fig 5](#) should be [Fig 4](#) and the image that appears as [Fig 4](#) should be [Fig 6](#).



 OPEN ACCESS

Citation: Miller AS, Shepherd GR, Fratantoni PS (2016) Correction: Offshore Habitat Preference of Overwintering Juvenile and Adult Black Sea Bass, *Centropristis striata*, and the Relationship to Year-Class Success. PLoS ONE 11(5): e0156355. doi:10.1371/journal.pone.0156355

Published: May 19, 2016

Copyright: © 2016 Miller et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Fig 4. Generalized additive model (GAM) plots identifying estimated effects of smoothed covariates on NEFSC spring survey catch of juvenile black sea bass. The magnitude of the y axis indicates the relative importance of each covariate with the effective degrees of freedom for the smoother denoted in the y axis label. Shaded areas represent 2 SE confidence intervals. Rugplots along the x-axis reflect the relative density of data points.

doi:10.1371/journal.pone.0156355.g001




Fig 5. Generalized additive model (GAM) plots identifying estimated effects of smoothed covariates on NEFSC spring survey catch of adult black sea bass. The magnitude of the y axis indicates the relative importance of each covariate with the effective degrees of freedom for the smoother denoted in the y axis label. Shaded areas represent 2 SE confidence intervals. Rugplots along the x-axis reflect the relative density of data points.

doi:10.1371/journal.pone.0156355.g002




Fig 6. Annual mean catch (n) per tow of black sea bass on the NEFSC spring survey. Data is constrained to survey strata used in the stock assessment. Solid black lines represent model predicted values with 2 standard error confidence intervals shown by the grey surrounding areas. Empirical mean catch of the raw data is denoted by a dotted line.

doi:10.1371/journal.pone.0156355.g003

Reference

1. Miller AS, Shepherd GR, Fratantoni PS (2016) Offshore Habitat Preference of Overwintering Juvenile and Adult Black Sea Bass, *Centropristis striata*, and the Relationship to Year-Class Success. PLoS ONE 11(1): e0147627. doi:10.1371/journal.pone.0147627 PMID: 26824350