CORRECTION

Correction: Risk factors for post-induction hypotension in patients with end-stage renal disease undergoing general anesthesia

Supachai Trikisayaveach, Laortip Rattanapittayaporn, Jatuporn Pakpirom, Chutida Sungworawongpana, Suttasinee Petsakul

There are errors in the author affiliations. The correct affiliations are as follows,

Supachai Trikisayaveach[1], Laortip Rattanapittayaporn[1], Jatuporn Pakpirom[1], Chutida Sungworawongpana[1], Suttasinee Petsakul[1]

1 Department of Anesthesiology, Faculty of Medicine, Prince of Songkla University, Hat Yai, Songkhla, Thailand.

Reference

1. Trikisayaveach S, Rattanapittayaporn L, Pakpirom J, Sungworawongpana C, Petsakul S. Risk factors for post-induction hypotension in patients with end-stage renal disease undergoing general anesthesia. PLoS One. 2026;21(5):e0350083. https://doi.org/10.1371/journal.pone.0350083 PMID: 42213723

Check for updates

OPEN ACCESS

Citation: Trikisayaveach S, Rattanapittayaporn L, Pakpirom J, Sungworawongpana C, Petsakul S (2026) Correction: Risk factors for post-induction hypotension in patients with end-stage renal disease undergoing general anesthesia. PLoS One 21(8): e0356297. https://doi.org/10.1371/journal.pone.0356297

Published: August 17, 2026

Copyright: © 2026 Trikisayaveach et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.