

Table S3. Results from autocorrelation analyses.

Network	Autocorrelation in persistence among functionally close species				Autocorrelation in persistence among phylogenetically close plant species				Autocorrelation in functional originality among phylogenetically close plant species			
	I	I_{max}	n	Class center	I	I_{max}	n	Class center	I	I_{max}	n	Class center
Albrecht [1]	-0.153	0.561	52	0.665	0.195	1.16	50	106.1	-0.054	0.694	50	106.1
Devoto [2]	0.121	0.73	34	0.377	-0.048	0.765	34	135.6	-0.082	0.703	34	135.6
Dicks [3]	-0.142	0.784	48	0.604	-0.032	0.83	48	48.5	0.029	1.576	48	48.5
Hegland [4]	0.018	0.472	188	0.484	-0.006	0.254	224	101.7	-0.034	0.564	186	97.7
Junker [5]	-0.046	0.553	154	2.348	0.066	0.577	188	113.1	-0.007	0.216	154	113.1
Memmott [6]	0.245	1.166	102	0.493	0.087	0.722	146	98.65	0.11	0.67	102	98.65
Weiner [7]	-0.068	0.536	774	0.533	-0.001	0.245	1062	95.2	0.124	0.646	760	90.55

References

1. Albrecht M, Riesen M, Schmid B (2010) Plant-pollinator network assembly along the chronosequence of a glacier foreland. *Oikos* 119: 1610–1624.
2. Devoto M, Bailey S, Craze P, Memmott J (2012) Understanding and planning ecological restoration of plant-pollinator networks. *Ecology Letters*: 319–328.
3. Dicks L V, Corbet SA, Pywell RF (2002) Compartmentalization in plant – insect flower visitor webs. *Journal of Animal Ecology* 71: 32–43.

4. Hegland SJ, Dunne J, Nielsen A, Memmott J (2010) How to monitor ecological communities cost-efficiently : The example of plant – pollinator networks.
5. Junker RR, Höcherl N, Blüthgen N (2010) Responses to olfactory signals reflect network structure of flower-visitor interactions. *Journal of Animal Ecology* 79: 818–823.
6. Memmott J (1999) The structure of a plant-pollinator food web. *Ecology Letters* 2: 276–280.
7. Weiner CN, Werner M, Linsenmair KE, Blüthgen N (2011) Land use intensity in grasslands : Changes in biodiversity , species composition and specialisation in flower visitor networks. *Basic and Applied Ecology* 12: 292–299.

