


S2 Figure

S2 Fig. Sub-analysis of DBPC RCTs applying a few-foods diet (FFD) intervention, subgrouped per rater (parent or other). Forest plot of FFD effects and homogeneity statistics.



DBPC = double-blind placebo-controlled
SMD = standardized mean difference

*inpatient study: ward ratings are commensurable to parent ratings

** after adjusting for multiple observations per study: SMD = 0.59, 95% CI: (0.06, 1.13), p = 0.029

References

1. Egger J, Carter CM, Graham PJ, Gumley D, Soothill JF. Controlled trial of oligoantigenic treatment in the hyperkinetic syndrome. *Lancet*. 1985;1(8428):540-5.
2. Kaplan BJ, McNicol J, Conte RA, Moghadam HK. Dietary replacement in preschool-aged hyperactive boys. *Pediatrics*. 1989;83(1):7-17.
3. Carter CM, Urbanowicz M, Hemsley R, Mantilla L, Strobel S, Graham PJ, et al. Effects of a few food diet in attention deficit disorder. *Arch Dis Child*. 1993;69(5):564-8. Epub 1993/11/01. PubMed PMID: 8257176; PubMed Central PMCID: PMC1029619.
4. Boris M, Mandel FS. Foods and additives are common causes of the attention deficit hyperactive disorder in children. *Ann Allergy*. 1994;72(5):462-8.
5. Schmidt MH, Mocks P, Lay B, Eisert HG, Fojkar R, Fritz-Sigmund D, et al. Does oligoantigenic diet influence hyperactive/conduct-disordered children--a controlled trial. *Eur Child Adolesc Psychiatry*. 1997;6(2):88-95.