

**S2 Table. Idealized Q-sorts for People Living With HIV and HIV Healthcare Provider**

<b>Summarized statements</b>	<b>PLWH-VP1</b> Conditional participation	<b>PLWH-VP2</b> Most motivated	<b>PLWH-VP3</b> Moderately motivated	<b>PLWH-VP4</b> Reticent	<b>Nb</b>	<b>HHP-VP1</b> Moderately motivated	<b>HHP-VP2</b> Most motivated	<b>HHP-VP3</b> Benefit-centered	<b>HHP-VP4</b> Reticent
trt-free phase monitoring must be adapted to each patient	1	2	0	1	1	-2	0	-4	1
trt-free phase follow-up must be regular and closely monitored	2	1	-1	0	2	0	1	2	-3
I would not participate if the medical follow-up of the trt-free phase were too restrictive	-2	-4	1	2	4	-1	-1	-2	2
I would refuse to participate if the innovative trt lasted more than 6 months	-2	-2	-2	-2	5	-1	-2	-2	-4
I would participate only if the innovative trt were administered on an outpatient basis	2	-2	-1	1	6	-3	0	-1	3
I would not participate because of the increased risk of HIV transmission during the trt-free phase	-1	-3	-2	-3	7	1	-1	-1	2
I would not participate if adverse effects of the innovative trt were too acute	-1	-2	2	1	8	3	3	0	4
I would not participate if the innovative trt affected vital organs	4	-1	4	0	9	4	0	1	1
The innovative trt must not entail irreversible adverse effects	3	0	4	1	10	4	-1	4	3
I would not participate if adverse effects lasted more than 5 days after the innovative trt	-1	-1	0	-3	11	1	-2	-2	-2

It is fundamental that my lifestyle does not change	1	-3	1	2	12	0	1	-3	3
It is fundamental that my current physician for HIV would supervise the three phases of the trial	0	2	0	2	3	2	-1	1	0
<i>It is fundamental that my current physician for HIV would believe in the trial</i>	4	3	2	3	13	2	2	2	2
A medical contact must always be available by phone	0	1	-1	-1	14	1	0	1	-1
I don't believe in this type of trial.	-4	-4	-4	-1	15	-3	-4	-3	-1
I believe that participating in a clinical trial guarantees better care	-2	1	0	0	17	0	-1	-1	-1
I think that it would be better to invest in access to ART for everyone	0	0	-4	0	18	-2	-3	2	0
I think that it would be better to invest in prevention	-1	-2	-3	2	19	-1	-4	0	-2
Financial compensation could entice me to participate	-3	-1	-2	-4	20	0	0	0	-4
I would participate in this trial even if the trt-free phase did not last more than 6 months	0	2	1	1	21	1	2	-3	0
<i>It is important to participate in HIV research</i>	3	4	3	3	22	1	3	3	1
It is important that medical advances for future generations be made possible	2	3	3	4	23	2	1	2	4
One motivation for participation is that one could forget about the disease	2	0	-3	-4	24	-2	1	-1	-1
Proposing a clinical (research) trial with no direct benefit for the patient has no sense	1	-1	-3	-2	25	-3	-2	4	2

Participating in this trial could be a way to avoid the long-term consequences of ART	0	4	3	0	26	2	3	3	-1
There is too much uncertainty about the adverse effects for me to participate	-3	-3	1	-2	16	-2	2	1	0
Having regular feedback from participating patients would motivate me	1	1	1	-1	27	-1	2	0	1
Regular information by physicians about the trial's results would motivate me	1	2	2	3	28	0	4	1	0
It is necessary that I have clear information about the trt.	3	3	-1	4	29	3	4	3	1
This trial is more suitable for people diagnosed more than ten years ago.	-4	0	-1	-1	30	-4	1	-2	-3
This trial should focus on those who have participated in only a few trials before	-2	0	-2	-1	31	-4	-3	-4	-2
This trial is more suitable for people who find it difficult to take ART on a daily basis	-3	-1	2	-2	32	3	-3	-1	-2
This trial is more suitable for people who do not work	-1	1	0	-3	33	-1	-2	0	-3

Trt= treatment