

S9 Table. Probability of smoking and smoking propensity by popularity quintiles of peers – probit average marginal effects.

	Tried 1996	1996	2002	2009	by 2009
<i>Smoking propensity:</i>					
Top pop. quintile	0.161*** (0.057)	0.051 (0.040)	0.179*** (0.039)	0.02 (0.064)	0.144** (0.067)
2nd pop. quintile	0.072 (0.058)	0.038 (0.038)	-0.092 (0.058)	-0.024 (0.051)	-0.077 (0.071)
3rd pop. quintile	0.078 (0.051)	0.064 (0.051)	-0.024 (0.057)	0.102* (0.057)	0.067 (0.070)
4th pop. quintile	-0.094* (0.050)	-0.006 (0.034)	-0.026 (0.050)	-0.062 (0.055)	-0.242*** (0.052)
5th pop. quintile	-0.018 (0.053)	0.008 (0.030)	-0.074* (0.045)	-0.033 (0.058)	-0.053 (0.069)

Regressions include school fixed effects. Standard errors clustered at the school level are shown in parenthesis. Peer smokers are those who smoke at least “once or twice a week” in 1995. Peer variables are at the grade level. Includes all covariates from S2 Table. *Significance at the 10% level; **Significance at the 5% level; ***Significance at the 1% level.