

S2 Table. Sensitivity analyses with different cutoffs to define the summary estimate of the TURN (n=112 drug-indication groups).

Cutoff *	TURN: frequency of missing doses (days)						Always acceptable
	1/month	2/month	3/month	1/week	2/week	3/week	
	-n	-n	-n	-n	-n	-n	-n
50% -n	5 ^a	5	10	35	20	13	24
60% -n	10	10	16	31	17	9	19
75% -n	22	16	25	20	11	8	10
80% -n	28	21	21	22	4	8	8
90% -n	56	17	11	17	6	0	5 ^b

n corresponds to the number of drug-indication groups

* the cutoff corresponds to the percentage used to define the summary estimate of the TURN (e.g., for the cutoff highlighted in grey [75%] in the primary analysis, the summary TURN was defined as response category over which 75% of physicians' estimates were located).

^a Drug-indication groups for which the TURN is "1 day/month" whatever the cut-off: antiepileptics in epilepsy, corticosteroids for prevention against graft rejection, glucocorticoids in adrenal insufficiency, hormonal contraceptives for systemic use, immunosuppressants for prevention against graft rejection

^b Drug-indication groups for which the TURN is "always acceptable" whatever the cut-off: anti-dementia drugs, anti-rheumatic products for osteoarthritis, other vitamins, urologics for urinary incontinence, progestogens and estrogens in combination for menopause