

S2 Table. Logistic regression models predicting attrition at follow-up

	Decedents			Lost			Refusal		
	OR	p-value	95% CI	OR	p-value	95% CI	OR	p-value	95% CI
Males	1.49	0.000	[1.28, 1.73]	1.09	0.318	[0.92-1.29]	0.97	0.833	[0.77, 1.23]
Baseline Age	1.08	0.000	[1.07, 1.09]	0.97	0.000	[0.96, 0.98]	0.97	0.000	[0.95, 0.98]
Years of education	1.01	0.392	[0.98, 1.03]	1.09	0.000	[1.07, 1.11]	1.09	0.000	[1.06, 1.12]
Residence (urban)	1.10	0.230	[0.97, 1.33]	2.00	0.000	[1.58, 2.54]	2.43	0.000	[1.70, 3.47]
Baseline Stroke	1.39	0.103	[0.93, 2.07]	2.28	0.002	[1.34, 3.87]	NA	NA	NA
Baseline Diabetes	2.10	0.000	[1.78, 2.43]	0.90	0.355	[0.72, 1.12]	0.78	0.134	[0.57, 1.07]
Baseline Heart Attack	2.18	0.000	[1.63, 2.90]	1.09	0.726	[0.66, 1.78]	0.80	0.605	[0.35, 1.83]
Baseline Hypertension	1.14	0.097	[0.97, 1.33]	0.75	0.003	[0.62, 0.94]	0.75	0.024	[0.58, 0.96]
Baseline Global Cognition	0.64	0.000	[0.57, 0.71]	0.84	0.002	[0.75, 0.94]	0.98	0.818	[0.84, 1.14]