


A) Males: mice group 1 spleen SIRPH 1 and 2 correlations




S1 and S2: First and second SIRPH experiment

1, 2 and A: CpG-1, CpG-2 and average of both

	α-Actin-1	α-Actin-2	MYC-1	MYC-2	PEG3-1	PEG3-2	LIT1-1	LIT1-2	SNRPN-1	SNRPN-2	IAP-1	IAP-2	L1-1	L1-2
Rho	0,60		0,84	0,70	-0,05	-0,01	0,33	0,16	0,46	0,15	0,81	0,19	0,84	0,77
P	7,52E-06		3,62E-14	4,21E-08	7,25E-01	9,19E-01	2,03E-02	2,64E-01	1,32E-03	3,10E-01	4,56E-12	2,02E-01	1,62E-13	1,28E-10

B) Females: mice cohort 1 spleen SIRPH 1 and 2 correlations



S1 and S2: First and second SIRPH experiment

1, 2 and A: CpG-1, CpG-2 and average of both

	α -Actin-1	α -Actin-2	MYC-1	MYC-2	PEG3-1	PEG3-2	LIT1-1	LIT1-2	SNRPN-1	SNRPN-2	IAP-1	IAP-2	L1-1	L1-2
Rho	0,60		0,89	0,77	-0,09	-0,11	-0,13	-0,25	0,34	0,35	0,85	0,45	0,91	0,76
P	8,72E-05		1,61E-16	3,86E-10	5,54E-01	5,02E-01	3,87E-01	1,08E-01	2,37E-02	2,72E-02	4,39E-14	2,05E-03	3,16E-18	8,52E-10