
Figure S7

Α


	parameter	value
1	Cytokine	IFN-gamma
2	Vendor	B ender
3	Samples included in this agreement	7
4	Proportion of both readings in range	18.9
5	Limits of agreement	0.64 to 1.66
6	Constant variance p.value	0.281
7	Constant ratio p.value	0.066
8	Ratio is 1 p.value	0.766

IFN-gamma - Bender




	parameter	value
1	Cytokine	IFN-gamma
2	Vendor	Bio-Rad
3	Samples included in this agreement	26
4	Proportion of both readings in range	70.3
5	Limits of agreement	0.62 to 1.71
6	Constant variance p.value	0.256
7	Constant ratio p.value	0.263
8	Ratio is 1 p.value	0.583

IFN-gamma – Bio-Rad




	parameter	value
1	Cytokine	IF N-gamma
2	Vendor	Invitrogen
3	Samples included in this agreement	4
4	Proportion of both readings in range	10.8
5	Limits of agreement	0.25 to 5.05
6	Constant variance p.value	0.899
7	Constant ratio p.value	0.436
8	Ratio is 1 p.value	0.763

IFN-gamma - Invitrogen



	parameter	value
1	Cytokine	IF N-gamma
2	Vendor	INV_MAG
3	Samples included in this agreement	2
4	Proportion of both readings in range	5.0
5	Limits of agreement	0.98 to 0.99
6	Constant variance p.value	NaN
7	Constant ratio p.value	NaN
8	Ratio is 1 p.value	0.059

IFN-gamma - INV_MAG



-	parameter	value
1	Cytokine	IFN-gamma
2	Vendor	Millipore
3	Samples included in this agreement	9
4	Proportion of both readings in range	24.3
5	Limits of agreement	0.73 to 1.28
6	Constant variance p.value	0.141
7	Constant ratio p.value	0.829
8	Ratio is 1 p.value	0.526

IFN-gamma - Millipore

