

S2 Table. Patient Profile with or without Admission Time after Ischemic Stroke.

Characteristics	Total (n=5705)	With AT (n=4493)	Without AT (n=1212)	P Value
Age, y, Mean ($\pm$ SD)	67.2 $\pm$ 12.2	67.1 $\pm$ 12.2	67.7 $\pm$ 11.9	0.102
Sex (% men)	3399 (59.6)	2688 (59.8)	711 (58.7)	0.464
NIHSS score at admission, median (IQR)	5(2-10)	5(3-11)	5(2-10)	<0.001
Smoking (%)	1478 (25.9)	1186 (26.4)	292 (24.1)	0.104
Hypertension (%)	3721 (65.2)	2905 (64.7)	816 (67.3)	0.083
Diabetes mellitus (%)	1280 (22.4)	987 (22.0)	293 (24.2)	0.102
Atrial fibrillation (%)	559 (9.8)	449 (10.0)	110 (9.1)	0.340
Dyslipidemia (%)	654 (11.5)	521 (11.6)	133 (11.0)	0.546
Heart failure (%)	129 (2.3)	98 (2.2)	31 (2.6)	0.434
Coronary heart disease (%)	958 (16.8)	751 (16.7)	207 (17.1)	0.763
IV rtPA within 3 hours arrived after stroke onset (%)	141 (2.5)	130 (2.9)	11 (0.9)	<0.001

Abbreviations: AT, admission time; SD, standard deviation; IQR, interquartile range;

NIHSS, National Institutes of Health Stroke Scale; IV rtPA, intravenous recombinant tissue-type plasminogen activator.