

S5 Table. Comparison of CKD patients with and without presence of diabetes mellitus

	Non- Diabetes mellitus (n=762)	Diabetes mellitus (n=269)	P value
Age (years)	57(33-75)	59(37-72)	0.30
Gender, male (%)	474 (62)	183 (68)	0.09
CVD, n (%)	224 (29)	146 (54)	<0.0001
Dialysis, n (%)	241(32)	58(22)	0.001
SGA>1, n (%)	221 (29)	99(37)	0.02
%HGS (n=736/249)	89(52-117)	72(47-107)	<0.0001
BMI (kg/m²)	24.2(19.5-30.0)	25.9(21.0-32.9)	<0.0001
LBMI (kg/m²; n=665/225)	17.1 (13.8-20.4)	17.2 (14.1-20.6)	0.10
FBMI (kg/m²; n=665/225)	6.8 (3.9-10.6)	8.4 (4.9-12.1)	<0.0001
S-Albumin (g/L)	35(29-41)	33(25-39)	<0.0001
hs CRP (mg/L)	2.9(0.5-26)	4.7(0.6-28.5)	0.0008

Data presented as median (10th - 90th percentile), number or percentage.

Abbreviations: CVD, cardiovascular disease; SGA, subjective global assessment; % HGS, handgrip strength as percentage of the controls; BMI, body mass index; LBMI, lean body mass index; FBMI, fat body mass index; S-Albumin, serum-albumin; hs CRP, high sensitivity C-reactive protein