

**S2 Fig. Format 2: Alternative package leaflet (intervention): Drug facts box with reading instruction (translation)**

<b>Side effects</b>			
Like all medications, Suffia® can have side effects. Not all undesired symptoms are necessarily due to taking the drug. Undesired symptoms can also arise without taking the drug.			
Occurrence of undesired symptoms over 5 years:			
<b>Undesired symptoms</b>	<b>Of 100 persons who take Suffia®:</b>	<b>Of 100 Persons who do NOT take Suffia®:</b>	<b>Undesired symptoms that are ascribed to Suffia® intake:</b>
Increased blood sugar levels	16 of 100	13 of 100	Taking Suffia® results in increased blood sugar levels in an additional 3 of 100 people.
Slow heart rate	5 of 100	2 of 100	Taking Suffia® results in slow heart rate in an additional 3 of 100 people.
Anemia	4 of 100	4 of 100	Taking Suffia® has no impact on anemia.
Depression	9 of 100	12 of 100	Taking Suffia® prevents depression in 3 of 100 people.
How to read this table: Of 100 people taking Suffia® over 5 years, 16 have increased blood sugar levels. Of 100 comparable people <b>not</b> taking Suffia®, 13 have increased blood sugar levels. Hence, 13 of 100 people taking Suffia® would have increased blood sugar levels even without taking the drug. Suffia® therefore causes an increase in blood sugar levels in 3 of 100 people.			