

Changes in *Emberiza* bunting communities and populations spanning 100 years in Korea

Chang-Yong Choi^{1,2}, Hyun-Young Nam^{1,3*}, Han-Kyu Kim^{4□}, Se-Young Park¹, Jong-Gil Park¹

S3 Table. The number of *Emberiza* buntings captured and banded in the Period II and III in Korea. There was no organized bird banding activities prior to the 1960s.

Species	Period II	Period III			Total
	1964-1970	1993-2017	2017-2019	Subtotal	
<i>Emberiza aureola</i>	266	224	0	224	490
<i>E. chrysophrys</i>	18	1,396	30	1,426	1,444
<i>E. cioides</i>	3,156	73	0	73	3,229
<i>E. elegans</i>	3,074	6,709	75	6,784	9,858
<i>E. fucata</i>	888	274	1	275	1,163
<i>E. jankowskii</i>	0	0	0	0	0
<i>E. leucocephalos</i>	9	5	1	6	15
<i>E. pallasi</i>	0	907	76	983	983
<i>E. pusilla</i>	57	1,336	85	1,421	1,478
<i>E. rustica</i>	61,055	2,539	33	2,572	63,627
<i>E. rutila</i>	46,826	2,409	13	2,422	49,248
<i>E. schoeniclus</i>	11	265	0	265	276
<i>E. spodocephala</i>	1,972	10,897	195	11,092	13,064
<i>E. sulphurata</i>	11	127	0	127	138
<i>E. tristrami</i>	2,079	2,089	33	2,122	4,201
<i>E. variabilis</i>	0	14	0	14	14
<i>E. yessoensis</i>	352	34	1	35	387
Total	119,774	29,298	543	29,841	149,615